

**THIS PAGE IS INSERTED BY OIPE SCANNING  
AND IS NOT PART OF THE OFFICIAL RECORD**

### **Best Available Images**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

**BLACK BORDERS**

**TEXT CUT OFF AT TOP, BOTTOM OR SIDES**

**FADED TEXT**

**BLURRY OR ILLEGIBLE TEXT**

**SKEWED/SLANTED IMAGES**

**COLORED PHOTOS HAVE BEEN RENDERED INTO BLACK AND WHITE**

**VERY DARK BLACK AND WHITE PHOTOS** ✓

**UNDECIPHERABLE GRAY SCALE DOCUMENTS**

**IMAGES ARE THE BEST AVAILABLE  
COPY. AS RESCANNING *WILL NOT*  
CORRECT IMAGES, PLEASE DO NOT  
REPORT THE IMAGES TO THE  
PROBLEM IMAGE BOX.**

10622687 - 10.27.03

Am

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
15 August 2002 (15.08.2002)

PCT

(10) International Publication Number  
WO 02/062423 A1

- (51) International Patent Classification<sup>7</sup>: A61P 31/18, 31/12, A61K 31/496, 31/501, 31/53, 31/506, C07D 401/14, 403/14, 241/02, 253/08, 251/02, 241/36, 471/02
- (21) International Application Number: PCT/US02/00455
- (22) International Filing Date: 2 January 2002 (02.01.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/266,183 2 February 2001 (02.02.2001) US  
60/314,406 23 August 2001 (23.08.2001) US
- (71) Applicant: BRISTOL-MYERS SQUIBB COMPANY  
[US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US).
- (72) Inventors: WANG, Tao; 1312 Town Brooke, Middletown, CT 06457 (US). ZHANG, Zhongxing; 14 Martleshamheath Lane, Madison, CT 06443 (US). MEANWELL, Nicholas, A.; 15 Valli Drive, East Hampton, CT 06424 (US). KADOW, John, F.; 9 Quarry Run, Wallingford, CT 06492 (US). YIN, Zhiwei; 234 Sherman Avenue, Apt. 80, Meriden, CT 06450 (US).
- (74) Agent: DUBOFF, Samuel, J.; Bristol-Myers Squibb Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/062423 A1

(54) Title: COMPOSITION AND ANTIVIRAL ACTIVITY OF SUBSTITUTED AZAINDOLEOXOACETIC PIPERAZINE DERIVATIVES

(57) Abstract: This invention provides compounds having drug and bio-affecting properties, their pharmaceutical compositions and method of use. In particular, the invention is concerned with azaindoleoxoacetyl piperazine derivatives. These compounds possess unique antiviral activity, whether used alone or in combination with other antivirals, anti-infectives, immunomodulators or HIV entry inhibitors. More particularly, the present invention relates to the treatment of HIV and AIDS.

**COMPOSITION AND ANTIVIRAL ACTIVITY OF SUBSTITUTED  
AZAINDOLEOXOACETIC PIPERAZINE DERIVATIVES**

**REFERENCE TO RELATED APPLICATIONS**

5

This application claims the benefit of U.S. Provisional Application Serial Numbers 60/314,406 filed August 23, 2001 and 60/266,183 filed February 2, 2001.

**BACKGROUND OF THE INVENTION**

**Field of the Invention**

10

This invention provides compounds having drug and bio-affecting properties, their pharmaceutical compositions and method of use. In particular, the invention is concerned with azaindole piperazine diamide derivatives that possess unique antiviral activity. More particularly, the present invention relates to compounds useful for the  
15 treatment of HIV and AIDS.

**Background Art**

HIV-1 (human immunodeficiency virus -1) infection remains a major medical  
20 problem, with an estimated 33.6 million people infected worldwide. The number of cases of HIV and AIDS (acquired immunodeficiency syndrome) has risen rapidly. In 1999, 5.6 million new infections were reported, and 2.6 million people died from AIDS. Currently available drugs for the treatment of HIV include six nucleoside reverse transcriptase (RT) inhibitors (zidovudine, didanosine, stavudine, lamivudine,  
25 zalcitabine and abacavir), three non-nucleoside reverse transcriptase inhibitors (nevirapine, delavirdine and efavirenz), and six peptidomimetic protease inhibitors (saquinavir, indinavir, ritonavir, nelfinavir, amprenavir and lopinavir). Each of these drugs can only transiently restrain viral replication if used alone. However, when used in combination, these drugs have a profound effect on viremia and disease  
30 progression. In fact, significant reductions in death rates among AIDS patients have been recently documented as a consequence of the widespread application of combination therapy. However, despite these impressive results, 30 to 50% of patients ultimately fail combination drug therapies. Insufficient drug potency, non-compliance, restricted tissue penetration and drug-specific limitations within certain

cell types (e.g. most nucleoside analogs cannot be phosphorylated in resting cells) may account for the incomplete suppression of sensitive viruses. Furthermore, the high replication rate and rapid turnover of HIV-1 combined with the frequent incorporation of mutations, leads to the appearance of drug-resistant variants and treatment failures when sub-optimal drug concentrations are present (Larder and Kemp; Gulick; Kuritzkes; Morris-Jones *et al*; Schinazi *et al*; Vacca and Condra; Flexner; Berkhout and Ren *et al*; (Ref. 6-14)). Therefore, novel anti-HIV agents exhibiting distinct resistance patterns, and favorable pharmacokinetic as well as safety profiles are needed to provide more treatment options.

10

Currently marketed HIV-1 drugs are dominated by either nucleoside reverse transcriptase inhibitors or peptidomimetic protease inhibitors. Non-nucleoside reverse transcriptase inhibitors (NNRTIs) have recently gained an increasingly important role in the therapy of HIV infections (Pedersen & Pedersen, Ref 15). At least 30 different classes of NNRTI have been described in the literature (De Clercq, Ref. 16) and several NNRTIs have been evaluated in clinical trials. Dipyridodiazepinone (nevirapine), benzoxazinone (efavirenz) and bis(heteroaryl) piperazine derivatives (delavirdine) have been approved for clinical use. However, the major drawback to the development and application of NNRTIs is the propensity for rapid emergence of drug resistant strains, both in tissue cell culture and in treated individuals, particularly those subject to monotherapy. As a consequence, there is considerable interest in the identification of NNRTIs less prone to the development of resistance (Pedersen & Pedersen, Ref 15).

25

Several indole derivatives including indole-3-sulfones, piperazino indoles, pyrazino indoles, and 5H-indolo[3,2-b][1,5]benzothiazepine derivatives have been reported as HIV-1 reverse transcriptase inhibitors (Greenlee *et al*, Ref. 1; Williams *et al*, Ref. 2; Romero *et al*, Ref. 3; Font *et al*, Ref. 17; Romero *et al*, Ref. 18; Young *et al*, Ref. 19; Genin *et al*, Ref. 20; Silvestri *et al*, Ref. 21). Indole 2-carboxamides have also been described as inhibitors of cell adhesion and HIV infection (Boschelli *et al*, US 5,424,329, Ref. 4). Finally, 3-substituted indole natural products (Semicochliodinol A and B, didemethylasterriquinone and isocochliodinol) were disclosed as inhibitors of HIV-1 protease (Fredenhagen *et al*, Ref. 22). Other indole derivatives exhibiting antiviral activity useful for treating HIV are disclosed in PCT WO 00/76521 (Ref. 93). Also, indole derivatives are disclosed in PCT WO 00/71535 (Ref. 94).

35



Structurally related aza-indole amide derivatives have been disclosed previously (Kato et al, Ref. 23; Levacher et al, Ref. 24; Dompe Spa, WO-09504742, Ref. 5(a); SmithKline Beecham PLC, WO-09611929, Ref. 5(b); Schering Corp., US-05023265, Ref. 5(c)). However, these structures differ from those claimed herein in that they are aza-indole mono-amide rather than unsymmetrical aza-indole piperazine diamide derivatives, and there is no mention of the use of these compounds for treating viral infections, particularly HIV. Other azaindoles have been also disclosed by Wang et al, Ref. 95. Nothing in these references can be construed to disclose or suggest the novel compounds of this invention and their use to inhibit HIV infection.

10

### REFERENCES CITED

#### Patent documents

- 15 1. Greenlee, W.J.; Srinivasan, P.C. Indole reverse transcriptase inhibitors. U.S. Patent 5,124,327.
2. Williams, T.M.; Ciccarone, T.M.; Saari, W. S.; Wai, J.S.; Greenlee, W.J.; Balani, S.K.; Goldman, M.E.; Theohrides, A.D. Indoles as inhibitors of HIV reverse transcriptase. European Patent 530907.
- 20 3. Romero, D.L.; Thomas, R.C.; Preparation of substituted indoles as anti-AIDS pharmaceuticals. PCT WO 93 / 01181.
- 25 4. Boschelli, D.H.; Connor, D.T.; Unangst, P.C. Indole-2-carboxamides as inhibitors of cell adhesion. U.S. Patent 5,424,329.
5. (a) Mantovanini, M.; Melillo, G.; Daffonchio, L. Tropyl 7-azaindol-3-ylcarboxyamides as antitussive agents. PCT WO 95/04742 (Dompe Spa). (b) Cassidy, F.; Hughes, I.; Rahman, S.; Hunter, D. J. Bisheteroaryl-carbonyl and carboxamide derivatives with 5HT 2C/2B antagonists activity. PCT WO 96/11929. (c) Scherlock, M. H.; Tom, W. C. Substituted 1H-pyrrolopyridine-3-carboxamides. U. S. Patent 5,023,265.

35

Other Publications

6. Larder, B.A.; Kemp, S.D. Multiple mutations in the HIV-1 reverse transcriptase confer high-level resistance to zidovudine (AZT). *Science*, 1989,  
5 246,1155-1158.
7. Gulick, R.M. Current antiretroviral therapy: An overview. *Quality of Life Research*, 1997, 6, 471-474.
- 10 8. Kuritzkes, D.R. HIV resistance to current therapies. *Antiviral Therapy*, 1997, 2 (Supplement 3), 61-67.
9. Morris-Jones, S.; Moyle, G.; Easterbrook, P.J. Antiretroviral therapies in HIV-1 infection. *Expert Opinion on Investigational Drugs*, 1997, 6(8),1049-1061.  
15
10. Schinazi, R.F.; Larder, B.A.; Mellors, J.W. Mutations in retroviral genes associated with drug resistance. *International Antiviral News*, 1997, 5,129-142,.
11. Vacca, J.P.; Condra, J.H. Clinically effective HIV-1 protease inhibitors.  
20 *Drug Discovery Today*, 1997, 2, 261-272.
12. Flexner, D. HIV-protease inhibitors. *Drug Therapy*, 1998, 338, 1281-1292.
13. Berkhout, B. HIV-1 evolution under pressure of protease inhibitors:  
25 Climbing the stairs of viral fitness. *J. Biomed. Sci.*, 1999, 6, 298-305.
14. Ren, S.; Lien, E. J. Development of HIV protease inhibitors: A survey. *Prog. Drug Res.*, 1998, 51, 1-31.
- 30 15. Pedersen, O.S.; Pedersen, E.B. Non-nucleoside reverse transcriptase inhibitors: the NNRTI boom. *Antiviral Chem. Chemother.* 1999, 10, 285-314.
16. (a) De Clercq, E. The role of non-nucleoside reverse transcriptase inhibitors (NNRTIs) in the therapy of HIV-1 infection. *Antiviral Research*, 1998, 38, 153-179.  
35 (b) De Clercq, E. Perspectives of non-nucleoside reverse transcriptase inhibitors (NNRTIs) in the therapy of HIV infection. *IL. Farmaco*, 1999, 54, 26-45.

17. Font, M.; Monge, A.; Cuartero, A.; Elorriaga, A.; Martinez-Irujo, J.J.; Alberdi, E.; Santiago, E.; Prieto, I.; Lasarte, J.J.; Sarobe, P. and Borrás, F. Indoles and pyrazino[4,5-*b*]indoles as nonnucleoside analog inhibitors of HIV-1 reverse transcriptase. *Eur. J. Med. Chem.*, 1995, 30, 963-971.
- 5
18. Romero, D.L.; Morge, R.A.; Genin, M.J.; Biles, C.; Busso, M.; Resnick, L.; Althaus, I.W.; Reusser, F.; Thomas, R.C and Tarpley, W.G.  
Bis(heteroaryl)piperazine (BHAP) reverse transcriptase inhibitors: structure-activity relationships of novel substituted indole analogues and the identification of 1-[(5-methanesulfonamido-1H-indol-2-yl)-carbonyl]-4-[3-[1-methylethyl)amino]-  
10 pyridinyl]piperazine momomethansulfonate (U-90152S), a second generation clinical candidate. *J. Med. Chem.*, 1993, 36, 1505-1508.
19. Young, S.D.; Amblard, M.C.; Britcher, S.F.; Grey, V.E.; Tran, L.O.; Lumma, W.C.; Huff, J.R.; Schleif, W.A.; Emini, E.E.; O'Brien, J.A.; Pettibone, D.J. 2-Heterocyclic indole-3-sulfones as inhibitors of HIV-reverse transcriptase. *Bioorg. Med. Chem. Lett.*, 1995, 5, 491-496.
- 15
20. Genin, M.J.; Poel, T.J.; Yagi, Y.; Biles, C.; Althaus, I.; Keiser, B.J.; Kopta, L.A.; Friis, J.M.; Reusser, F.; Adams, W.J.; Olmsted, R.A.; Voorman, R.L.; Thomas, R.C. and Romero, D.L. Synthesis and bioactivity of novel bis(heteroaryl)piperazine (BHAP) reverse transcriptase inhibitors: structure-activity relationships and increased metabolic stability of novel substituted pyridine analogs. *J. Med. Chem.*, 1996, 39, 5267-5275.
- 25
21. Silvestri, R.; Artico, M.; Bruno, B.; Massa, S.; Novellino, E.; Greco, G.; Marongiu, M.E.; Pani, A.; De Montis, A and La Colla, P. Synthesis and biological evaluation of 5*H*-indolo[3,2-*b*][1,5]benzothiazepine derivatives, designed as conformationally constrained analogues of the human immunodeficiency virus type 1  
30 reverse transcriptase inhibitor L-737,126. *Antiviral Chem. Chemother.* 1998, 9, 139-148.
22. Fredenhagen, A.; Petersen, F.; Tintelnot-Blomley, M.; Rosel, J.; Mett, H and Hug, P. J. Semicochliodinol A and B: Inhibitors of HIV-1 protease and EGF-R protein Tyrosine Kinase related to Asterriquinones produced by the fungus *Chrysosporium nardarium*. *Antibiotics*, 1997, 50, 395-401.
- 35

23. Kato, M.; Ito, K.; Nishino, S.; Yamakuni, H.; Takasugi, H. New 5-HT<sub>3</sub> (Serotonin-3) receptor antagonists. IV. Synthesis and structure-activity relationships of azabicycloalkaneacetamide derivatives. *Chem. Pharm. Bull.*, **1995**, *43*, 1351-1357.
- 5
24. Levacher, V.; Benoit, R.; Duflos, J.; Dupas, G.; Bourguignon, J.; Queguiner, G. Broadening the scope of NADH models by using chiral and non chiral pyrrolo [2,3-*b*] pyridine derivatives. *Tetrahedron*, **1991**, *47*, 429-440.
- 10
25. Shadrina, L.P.; Dormidontov, Yu.P.; Ponomarev, V.G.; Lapkin, I.I. Reactions of organomagnesium derivatives of 7-aza- and benzoindoles with diethyl oxalate and the reactivity of ethoxalylindoles. *Khim. Geterotsikl. Soedin.*, **1987**, 1206-1209.
- 15
26. Sycheva, T.V.; Rubtsov, N.M.; Sheinker, Yu.N.; Yakhontov, L.N. Some reactions of 5-cyano-6-chloro-7-azaindoles and lactam-lactim tautomerism in 5-cyano-6-hydroxy-7-azaindoles. *Khim. Geterotsikl. Soedin.*, **1987**, 100-106.
27. (a) Desai, M.; Watthey, J.W.H.; Zuckerman, M. A convenient preparation of 1-arylpiperazines. *Org. Prep. Proced. Int.*, **1976**, *8*, 85-86. (b) Adamczyk, M.; Fino, J.R. Synthesis of procainamide metabolites. N-acetyl desethylprocainamide and desethylprocainamide. *Org. Prep. Proced. Int.* **1996**, *28*, 470-474. (c) Rossen, K.; Weissman, S.A.; Sager, J.; Reamer, R.A.; Askin, D.; Volante, R.P.; Reider, P.J. Asymmetric Hydrogenation of tetrahydropyrazines: Synthesis of (*S*)-piperazine 2-tert-butylcarboxamide, an intermediate in the preparation of the HIV protease inhibitor Indinavir. *Tetrahedron Lett.*, **1995**, *36*, 6419-6422. (d) Wang, T.; Zhang, Z.; Meanwell, N.A. Benzoylation of Dianions: Preparation of mono-Benzoylated Symmetric Secondary Diamines. *J. Org. Chem.*, **1999**, *64*, 7661-7662.
- 20
28. Li, H.; Jiang, X.; Ye, Y.-H.; Fan, C.; Romoff, T.; Goodman, M. 3-(Diethoxyphosphoryloxy)-1,2,3-benzotriazin-4(3*H*)-one (DEPBT): A new coupling reagent with remarkable resistance to racemization. *Organic Lett.*, **1999**, *1*, 91-93.
- 25
29. Harada, N.; Kawaguchi, T.; Inoue, I.; Ohashi, M.; Oda, K.; Hashiyama, T.; Tsujihara, K. Synthesis and antitumor activity of quaternary salts of 2-(2'-oxoalkoxy)-9-hydroxyellipticines. *Chem. Pharm. Bull.*, **1997**, *45*, 134-137.
- 30
- 35

30. Schneller, S. W.; Luo, J.-K. Synthesis of 4-amino-1*H*-pyrrolo[2,3-*b*]pyridine (1,7-Dideazaadenine) and 1*H*-pyrrolo[2,3-*b*]pyridin-4-ol (1,7-Dideazahypoxanthine). *J. Org. Chem.*, **1980**, *45*, 4045-4048.
- 5 31. Shiotani, S.; Tanigochi, K. Furopyridines. XXII [1]. Elaboration of the C-substituent *alpha* to the heteronitrogen atom of furo[2,3-*b*]-, -[3,2-*b*]-, -[2,3-*c*]- and -[3,2-*c*]pyridine. *J. Het. Chem.*, **1997**, *34*, 901-907.
- 10 32. Minakata, S.; Komatsu, M.; Ohshiro, Y. Regioselective functionalization of 1*H*-pyrrolo[2,3-*b*]pyridine via its N-oxide. *Synthesis*, **1992**, 661-663.
- 15 33. Klemm, L. H.; Hartling, R. Chemistry of thienopyridines. XXIV. Two transformations of thieno[2,3-*b*]pyridine 7-oxide (1). *J. Het. Chem.*, **1976**, *13*, 1197-1200.
34. Antonini, I.; Claudi, F.; Cristalli, G.; Franchetti, P.; Crifantini, M.; Martelli, S. Synthesis of 4-amino-1- $\beta$ -D-ribofuranosyl-1*H*-pyrrolo[2,3-*b*]pyridine (1-Deazatubercidin) as a potential antitumor agent. *J. Med. Chem.*, **1982**, *25*, 1258-1261.
- 20 35. (a) Regnouf De Vains, J.B.; Papet, A.L.; Marsura, A. New symmetric and unsymmetric polyfunctionalized 2,2'-bipyridines. *J. Het. Chem.*, **1994**, *31*, 1069-1077. (b) Miura, Y.; Yoshida, M.; Hamana, M. Synthesis of 2,3-fused quinolines from 3-substituted quinoline 1-oxides. Part II, *Heterocycles*, **1993**, *36*, 1005-1016.
- 25 (c) Profft, V.E.; Rolle, W. Über 4-merkaptoverbindendes 2-methylpyridins. *J. Prakt. Chem.*, **1960**, *283* (11), 22-34.
- 30 36. Nesi, R.; Giomi, D.; Turchi, S.; Tedeschi, P.; Ponticelli, F. A new one step synthetic approach to the isoxazolo[4,5-*b*]pyridine system. *Synth. Comm.*, **1992**, *22*, 2349-2355.
37. (a) Walser, A.; Zenchoff, G.; Fryer, R.I. Quinazolines and 1,4-benzodiazepines. 75. 7-Hydroxyaminobenzodiazepines and derivatives. *J. Med. Chem.*, **1976**, *19*, 1378-1381. (b) Barker, G.; Ellis, G.P. Benzopyrone. Part I. 6-Amino- and 6-hydroxy-2-substituted chromones. *J. Chem. Soc.*, **1970**, 2230-2233.
- 35 38. Ayyangar, N.R.; Lahoti, R.J.; Daniel, T. An alternate synthesis of 3,4-diaminobenzophenone and mebendazole. *Org. Prep. Proced. Int.*, **1991**, *23*, 627-631.

39. Mahadevan, I.; Rasmussen, M. Ambident heterocyclic reactivity: The alkylation of pyrrolopyridines (azaindoles, diazaindenes). *Tetrahedron*, **1993**, *49*, 7337-7352.
- 5 40. Chen, B.K.; Saksela, K.; Andino, R.; Baltimore, D. Distinct modes of human immunodeficiency type 1 proviral latency revealed by superinfection of nonproductively infected cell lines with recombinant luciferase-encoding viruses. *J. Virol.*, **1994**, *68*, 654-660.
- 10 41. Bodanszky, M.; Bodanszky, A. "The Practice of Peptide Synthesis" 2<sup>nd</sup> Ed., Springer-Verlag: Berlin Heidelberg, Germany, **1994**.
42. Albericio, F. *et al. J. Org. Chem.* **1998**, *63*, 9678.
- 15 43. Knorr, R. *et al. Tetrahedron Lett.* **1989**, *30*, 1927.
44. (a) Jaszay Z. M. *et al. Synth. Commun.*, **1998** *28*, 2761 and references cited therein; (b) Bernasconi, S. *et al. Synthesis*, **1980**, 385.
- 20 45. (a) Jaszay Z. M. *et al. Synthesis*, **1989**, 745 and references cited therein; (b) Nicolaou, K. C. *et al. Angew. Chem. Int. Ed.* **1999**, *38*, 1669.
46. Ooi, T. *et al. Synlett.* **1999**, 729.
- 25 47. Ford, R. E. *et al. J. Med. Chem.* **1986**, *29*, 538.
48. (a) Yeung, K.-S. *et al. Bristol-Myers Squibb Unpublished Results.* (b) Wang, W. *et al. Tetrahedron Lett.* **1999**, *40*, 2501.
- 30 49. Brook, M. A. *et al. Synthesis*, **1983**, 201.

50. Yamazaki, N. *et al. Tetrahedron Lett.* **1972**, 5047.
51. Barry A. Bunin "The Combinatorial Index" 1998 Academic Press, San Diego / London pages 78-82.
52. Richard C. Larock *Comprehensive Organic Transformations* 2nd Ed. 1999, John Wiley and Sons New York.
53. M.D. Mullican *et.al. J.Med. Chem.* **1991**, 34, 2186-2194.
54. Protective groups in organic synthesis 3rd ed. / Theodora W. Greene and Peter G.M. Wuts. New York : Wiley, 1999.
55. Katritzky, Alan R. Lagowski, Jeanne M. The principles of heterocyclic Chemistry New York : Academic Press, 1968
56. Paquette, Leo A. Principles of modern heterocyclic chemistry New York : Benjamin.
57. Katritzky, Alan R.; Rees, Charles W.; *Comprehensive heterocyclic chemistry : the structure, reactions, synthesis, and uses of heterocyclic compounds* 1st ed. Oxford (Oxfordshire) ; New York : Pergamon Press, 1984. 8 v.
58. Katritzky, Alan R. *Handbook of heterocyclic* 1st ed Oxford (Oxfordshire) ; New York : Pergamon Press, 1985.
59. Davies, David I *Aromatic Heterocyclic* Oxford ; New York : Oxford University Press, 1991.

60. Ellis, G. P. Synthesis of fused Chichester [Sussex] ; New York : Wiley,  
c1987-c1992. Chemistry of heterocyclic compounds ; v. 47.
61. Joule, J. A Mills, K. ,Smith, G. F. Heterocyclic Chemistry , 3rd ed London  
5 ;New York Chapman & Hall, 1995.
62. Katritzky, Alan R., Rees, Charles W. , Scriven, Eric F. V. Comprehensive  
heterocyclic chemistry II : a review of the literature 1982-1995.
- 10 63. The structure, reactions, synthesis, and uses of heterocyclic compounds 1st  
ed. Oxford ; New York : Pergamon, 1996. 11 v. in 12 : ill. ; 28 cm.
64. Eicher, Theophil, Hauptmann, Siegfried. The chemistry of heterocycles :  
structure, reactions, syntheses, and applications Stuttgart ; New York : G. Thieme,  
15 1995.
65. Grimmett, M. R. Imidazole and benzimidazole Synthesis London ; San Diego  
: Academic Press, 1997.
- 20 66. Advances in heterocyclic chemistry. Published in New York by Academic  
Press, starting in 1963- present.
67. Gilchrist, T. L. (Thomas Lonsdale) Heterocyclic chemistry 3rd ed. Harlow,  
Essex : Longman, 1997. 414 p. : ill. ; 24 cm.
- 25 68. Farina, Vittorio; Roth, Gregory P. Recent advances in the Stille reaction;  
*Adv. Met.-Org. Chem.* 1996, 5, 1-53.



69. Farina, Vittorio; Krishnamurthy, Venkat; Scott, William J. The Stille reaction ; *Org. React.* (N. Y.) (1997), 50, 1-652.
70. Stille, J. K. *Angew. Chem. Int. Ed. Engl.* 1986, 25, 508-524.
- 5 71. Norio Miyaura and Akiro Suzuki *Chem Rev.* 1995, 95, 2457.
72. Home, D.A. *Heterocycles* 1994, 39, 139.
- 10 73. Kamitori, Y. et.al. *Heterocycles*, 1994, 37(1), 153.
74. Shawali, J. *Heterocyclic Chem.* 1976, 13, 989.
75. a) Kende, A.S. et al. *Org. Photochem. Synth.* 1972, 1, 92. b) Hanks, L.V.;  
15 *Biochem. Prep.* 1966, 11, 63. c) *Synth. Meth.* 22, 837.
76. Hulton et. al. *Synth. Comm.* 1979, 9, 789.
77. Pattanayak, B.K. et.al. *Indian J. Chem.* 1978, 16, 1030.
- 20 78. *Chemische Berichte* 1902, 35, 1545.
79. *Chemische Berichte* Ibid 1911, 44, 493.
- 25 80. Moubarak, I., Vessiere, R. *Synthesis* 1980, Vol. 1, 52-53.
81. *Ind J. Chem.* 1973, 11, 1260.


82. Roomi et.al. *Can J. Chem.* **1970**, *48*, 1689.
83. Sorrel, T.N. *J. Org. Chem.* **1994**, *59*, 1589.
- 5 84. Nitz, T.J. et. al. *J. Org. Chem.* **1994**, *59*, 5828-5832.
85. Bowden, K. et.al. *J. Chem. Soc.* **1946**, 953.
86. Nitz, T.J. et. al. *J. Org. Chem.* **1994**, *59*, 5828-5832.
- 10 87. Scholkopf et. al. *Angew. Int. Ed. Engl.* **1971**, *10(5)*, 333.
88. (a) Behun, J. D.; Levine, R. *J. Org. Chem.* **1961**, *26*, 3379. (b) Rossen, K.; Weissman, S.A.; Sager, J.; Reamer, R.A.; Askin, D.; Volante, R.P.; Reider, P.J.
- 15 Asymmetric Hydrogenation of tetrahydropyrazines: Synthesis of (*S*)-piperazine 2-tert-butylcarboxamide, an intermediate in the preparation of the HIV protease inhibitor Indinavir. *Tetrahedron Lett.*, **1995**, *36*, 6419-6422. (c) Jenneskens, L. W.; Mahy, J.; den Berg, E. M. M. de B.-v.; Van der Hoef, I.; Lugtenburg, J. *Recl. Trav. Chim. Pays-Bas* **1995**, *114*, 97.
- 20 89. Wang, T.; Zhang, Z.; Meanwell, N.A. Benzoylation of Dianions: Preparation of mono-Benzoylated Symmetric Secondary Diamines. *J. Org. Chem.*, **1999**, *64*, 7661-7662.
- 25 90. (a) Adamczyk, M.; Fino, J.R. Synthesis of procainamide metabolites. N-acetyl desethylprocainamide and desethylprocainamide. *Org. Prep. Proced. Int.* **1996**, *28*, 470-474. (b) Wang, T.; Zhang, Z.; Meanwell, N.A. Regioselective mono-Benzoylation of Unsymmetrical Piperazines. *J. Org. Chem.*, in press.

91. Masuzawa, K.; Kitagawa, M.; Uchida, H. *Bull Chem. Soc. Jpn.* **1967**, *40*, 244-245.
- 5 92. Furber, M.; Cooper, M. E.; Donald, D. K. *Tetrahedron Lett.* **1993**, *34*, 1351-1354.
93. Blair, W. S. et al, PCT WO 00/76521 published December 21, 2000.
- 10 94. Mavunkel, B. J. et al, PCT WO 00/71535 published November 30, 2000.
95. Wang, T. et al, PCT WO 01/62255 published August 30, 2001.

### SUMMARY DESCRIPTION OF THE INVENTION

15 The present invention comprises compounds of Formula I, or pharmaceutically acceptable salts thereof, which are effective antiviral agents, particularly as inhibitors of HIV.

20 A first embodiment of a first aspect of the invention are compounds of Formula I, including pharmaceutically acceptable salts thereof,




I

wherein:

25

Q is selected from the group consisting of:



$R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$ , are independently selected from the group consisting of hydrogen, halogen, cyano, nitro,  $\text{COOR}^8$ ,  $\text{XR}^{57}$ ,  $\text{C(O)R}^{57}$ ,  $\text{C(O)NR}^{55}\text{R}^{56}$ , B, D, and E with the proviso that at least one of  $R^1$ - $R^4$  is selected from B or E;

5


m is 1 or 2;

$R^5$  is hydrogen or  $(\text{CH}_2)_n\text{R}^{44}$  wherein n is 0-6;

10  $R^6$  is O or does not exist;

-- may represent a carbon-carbon bond;

A is selected from the group consisting of  $\text{C}_{1-6}$ alkoxy, aryl and heteroaryl; in which  
 15 said aryl is phenyl or naphthyl; said heteroaryl is selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, triazinyl, furanyl, thienyl, pyrrolyl, imidazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, quinolinyl, isoquinolinyl, benzofuranyl, benzothienyl, benzoimidazolyl and benzothiazolyl; and said aryl or heteroaryl is optionally substituted with one or two of the same or different amino, nitro, cyano,  
 20  $\text{C}_{1-6}$ alkoxy,  $-\text{C(O)NH}_2$ ,  $\text{C}_{1-6}$ alkyl,  $-\text{NHC(O)CH}_3$ , halogen or trifluoromethyl;  
 -W- is



B is selected from the group consisting of  $-\text{C(=NR}^{46})(\text{R}^{47})$ ,  $\text{C(O)NR}^{40}\text{R}^{41}$ , aryl, heteroaryl, heteroalicyclic,  $\text{S(O)}_q\text{R}^8$ ,  $\text{P(O)(R}^8)_q(\text{OR}^8)_{2-q}$ ,  $\text{P(S)(R}^8)_q(\text{OR}^8)_{2-q}$ ,  $\text{C(O)R}^7$ ,  
 25  $\text{XR}^8$ ,  $(\text{C}_{1-6})\text{alkylNR}^{40}\text{R}^{41}$ , and  $(\text{C}_{1-6})\text{alkylCOOR}^8$  wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to three same or different substituents selected from the group F;

q is 0, 1, or 2;

30

D is selected from the group consisting of  $(\text{C}_{1-6})\text{alkyl}$ ,  $(\text{C}_{3-7})\text{cycloalkyl}$ ,  $(\text{C}_{2-6})\text{alkenyl}$ ,  $(\text{C}_{3-7})\text{cycloalkenyl}$ ,  $(\text{C}_{2-6})\text{alkynyl}$ , wherein said  $(\text{C}_{1-6})\text{alkyl}$ ,

(C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, and (C<sub>2-6</sub>)alkynyl are optionally substituted with one to three same or different halogens or from one to three same or different substituents selected from the group F;

- 5 E is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, and (C<sub>2-6</sub>)alkynyl are substituted with B;
  
- 10 F is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, (C<sub>1-6</sub>)alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, thiohydroxy, (C<sub>1-6</sub>)thioalkoxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, cyano, halogen, nitro, carbonyl, thiocarbonyl, benzyl, O-thiocarbamyl, N-thiocarbamyl, C-thioamido, -NR<sup>42</sup>C(O)-(C<sub>1-6</sub>)alkyl, -NR<sup>42</sup>C(O)-
- 15 (C<sub>3-6</sub>)cycloalkyl, -NR<sup>42</sup>C(O)-aryl, -NR<sup>42</sup>C(O)-heteroaryl, -NR<sup>42</sup>C(O)-heteroalicyclic, a cyclic N-amido, -NR<sup>42</sup>S(O)<sub>2</sub>-(C<sub>1-6</sub>)alkyl, -NR<sup>42</sup>S(O)<sub>2</sub>-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>42</sup>S(O)<sub>2</sub>-aryl, -NR<sup>42</sup>S(O)<sub>2</sub>-heteroaryl, -NR<sup>42</sup>S(O)<sub>2</sub>-heteroalicyclic, O-carboxy, sulfinyl, sulfonyl, -S(O)<sub>2</sub>NR<sup>42</sup>R<sup>43</sup>, phosphonyl, NR<sup>42</sup>R<sup>43</sup>, (C<sub>1-6</sub>)alkylC(O)NR<sup>42</sup>R<sup>43</sup>, C(O)NR<sup>42</sup>R<sup>43</sup>, NHC(O)NR<sup>42</sup>R<sup>43</sup>, OC(O)NR<sup>42</sup>R<sup>43</sup>,
- 20 NHC(O)OR<sup>54</sup>, (C<sub>1-6</sub>)alkylNR<sup>42</sup>R<sup>43</sup>, COOR<sup>54</sup>, and (C<sub>1-6</sub>)alkylCOOR<sup>54</sup> wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, (C<sub>1-6</sub>)alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, (C<sub>1-6</sub>)thioalkoxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, are optionally substituted with one to nine same or different halogens or from one to five same or different substituents selected from the group G;
  
- 25 G is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, (C<sub>1-6</sub>)alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, thiohydroxy, (C<sub>1-6</sub>)thioalkoxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, cyano, halogen, nitro, carbonyl, thiocarbonyl, benzyl, O-thiocarbamyl, N-thiocarbamyl, C-thioamido, -NR<sup>48</sup>C(O)-(C<sub>1-6</sub>)alkyl, -NR<sup>48</sup>C(O)-
- 30 (C<sub>3-6</sub>)cycloalkyl, -NR<sup>48</sup>C(O)-aryl, -NR<sup>48</sup>C(O)-heteroaryl, -NR<sup>48</sup>C(O)-heteroalicyclic, a cyclic N-amido, -NR<sup>48</sup>S(O)<sub>2</sub>-(C<sub>1-6</sub>)alkyl, -NR<sup>48</sup>S(O)<sub>2</sub>-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>48</sup>S(O)<sub>2</sub>-aryl, -NR<sup>48</sup>S(O)<sub>2</sub>-heteroaryl, -NR<sup>48</sup>S(O)<sub>2</sub>-heteroalicyclic, O-carboxy, sulfinyl, sulfonyl, sulfonamide, phosphonyl, NR<sup>48</sup>R<sup>49</sup>,
- 35 (C<sub>1-6</sub>)alkyl C(O)NR<sup>48</sup>R<sup>49</sup>, C(O)NR<sup>48</sup>R<sup>49</sup>, NHC(O)NR<sup>48</sup>R<sup>49</sup>, OC(O)NR<sup>48</sup>R<sup>49</sup>, NHC(O)OR<sup>54</sup>, (C<sub>1-6</sub>)alkylNR<sup>48</sup>R<sup>49</sup>, COOR<sup>54</sup>, and (C<sub>1-6</sub>)alkylCOOR<sup>54</sup>;

R<sup>7</sup> is selected from the group consisting of aryl, heteroaryl, and heteroalicyclic wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or with from one to three same or different substituents selected from the group F;

5

R<sup>8</sup> is selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to six same or different halogens or from one to five same or different substituents selected from the group F;

10

R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, are each independently selected from the group consisting of hydrogen, or (C<sub>1-6</sub>)alkyl wherein each of said (C<sub>1-6</sub>)alkyl being optionally substituted with one to three same or different halogens;

15

X is selected from the group consisting of NR<sup>5</sup>, O, and S;

R<sup>40</sup> and R<sup>41</sup> are independently selected from the group consisting of

20

Hydrogen;

or (C<sub>1-6</sub>)alkyl or (C<sub>3-7</sub>)cycloalkyl substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F; or (C<sub>1-6</sub>)alkoxy, aryl, heteroaryl, heteroalicyclic or R<sup>40</sup> and R<sup>41</sup> taken together with the nitrogen to which they are attached form a heteroalicyclic ring which may contain up to 5 additional heteroatoms selected from N, O, S(O)<sub>m'</sub>, wherein m' is 0, 1, or 2; and wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

25

with the proviso that only one of R<sup>40</sup> and R<sup>41</sup> may be hydrogen.

30

R<sup>42</sup> and R<sup>43</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, heteroalicyclic or R<sup>42</sup> and R<sup>43</sup> taken together with the nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to 5 additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2; and wherein said (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy,

35

(C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to nine same or different halogens or from one to five same or different substituents selected from the group G;

5 R<sup>44</sup> is selected from the group consisting of:

(1) H, (C<sub>1-6</sub>)alkyl, (C<sub>3-6</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-6</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, halogen, CN, nitro, Ar, COOR<sup>50</sup>, COOAr, -CONR<sub>a</sub>R<sub>b</sub>, TR<sup>50</sup>, NR<sub>a</sub>R<sub>b</sub>, -NC(O)NR<sub>a</sub>R<sub>b</sub>, -OC(O)R<sup>50</sup>, -C[N(R<sub>a</sub>)<sub>2</sub>] = N-T-R<sub>b</sub>, YR<sup>50</sup>, -C(O)R<sup>50</sup>, -C(O)Ar,  
 10 -S(O)R<sub>a</sub> or -S(O)<sub>2</sub>R<sub>a</sub>, provided when R<sup>44</sup> is -S(O)R<sub>a</sub> or -S(O)<sub>2</sub>R<sub>a</sub> then R<sub>a</sub> is not H; and

(2) a 4-7 membered heterocyclic ring, optionally substituted with R<sup>50</sup>, which may contain 1-3 heteroatoms selected from the group consisting of O, S, SO, SO<sub>2</sub>, N, and NR<sup>52</sup>, wherein R<sup>52</sup> is selected from the group consisting of hydrogen,  
 15 (C<sub>1-4</sub>)alkyl, (C<sub>2-4</sub>)alkenyl and (C<sub>2-4</sub>)alkynyl;

T is S or O;

Ar is phenyl or heteroaryl; wherein said phenyl or heteroaryl is optionally substituted  
 20 with one to three of the same or different halogens, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkyl or amino;

R<sub>a</sub> and R<sub>b</sub> are each independently H, (C<sub>1-6</sub>)alkyl or phenyl;

R<sup>46</sup> is selected from the group consisting of H, OR<sup>8</sup>, and NR<sup>40</sup>R<sup>41</sup>;  
 25

R<sup>47</sup> is selected from the group consisting of H, amino, halogen, and (C<sub>1-6</sub>)alkyl;

R<sup>48</sup> and R<sup>49</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, heteroalicyclic or R<sup>48</sup> and R<sup>49</sup> taken together with the  
 30 nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to 5 additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2;

R<sup>50</sup> is selected from the group consisting of H, (C<sub>1-6</sub>)alkyl, (C<sub>3-6</sub>)cycloalkyl, and benzyl, each of said alkyl, cycloalkyl and benzyl being optionally substituted with one to three same or different halogen, amino, OH, CN or NO<sub>2</sub>;

5 R<sup>51</sup> is selected from the group consisting of H, (C<sub>1-6</sub>)alkyl, (C<sub>3-6</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-6</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl or C(O)R<sup>53</sup>, wherein R<sup>53</sup> is H, (C<sub>1-6</sub>)alkyl, or (C<sub>3-6</sub>)cycloalkyl and each of said (C<sub>1-6</sub>)alkyl and (C<sub>3-6</sub>)cycloalkyl being optionally substituted with one to three same or different halogen, amino, OH, CN or NO<sub>2</sub>;

10

Y is O, S or NR<sup>50</sup>R<sup>51</sup>;

R<sup>54</sup> is selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to six same or different halogens or from one to five same or different substituents selected from the group consisting of: amino, OH, CN and NO<sub>2</sub>;

20 R<sup>54'</sup> is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to six same or different halogens or from one to five same or different substituents selected from the group consisting of: amino, OH, CN and NO<sub>2</sub>;

R<sup>55</sup> and R<sup>56</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl; and


30 R<sup>57</sup> is selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl.

With the proviso that in the formulas above the carbon atoms which comprise the carbon-carbon double bond of any C<sub>number</sub>-C<sub>number</sub> alkenyl or the carbon-carbon triple bond of said C<sub>number</sub>-C<sub>number</sub> alkynyl are not the point of attachment to the oxygen, nitrogen, or sulfur to which it is said to be attached;

35



A more preferred embodiment of a first aspect of the invention are compounds of Formula I, including pharmaceutically acceptable salts thereof,



I

wherein:

$R^1$  is hydrogen

$R^2$  and  $R^3$ , are each independently selected from the group (a)-(k) consisting of:

(a) hydrogen,

(b) halogen,

(c) cyano,

(d) nitro,

(e) amino,

(f)  $C_{1-4}$ alkylamino,

(g) di( $C_{1-2}$ alkyl)amino,

(h) hydroxy,

(i)  $C_{1-3}$ alkyl optionally substituted with one to three same or different halogen, hydroxy,  $C_{1-2}$ alkoxy, amino,  $C_{1-4}$ alkylamino, di( $C_{1-4}$ alkyl)amino, cyano,

(j)  $C_{1-6}$ alkoxy,

(k) heteroaryl, said heteroaryl is selected from the group consisting of

pyridinyl, pyrazinyl, pyridazinyl, pyrimidinyl, furanyl, thienyl, benzothienyl, thiazolyl, isothiazolyl, oxazolyl, benzooxazolyl, isoxazolyl, imidazolyl, benzoimidazolyl, 1H-imidazo[4,5-b]pyridin-2-yl, 1H-imidazo[4,5-c]pyridin-2-yl, oxadiazolyl, thiadiazolyl, pyrazolyl, tetrazolyl, tetrazinyl, triazinyl and triazolyl, and said heteroaryl is optionally substituted with  $C_{1-6}$  alkyl groups

(l) phenyl which is independently substituted with one to three same or different halogen, hydroxy,  $C_{1-2}$ alkoxy, amino,  $C_{1-4}$ alkylamino, di( $C_{1-4}$ alkyl)amino, cyano,

$R^4$  is selected from the group consisting of hydrogen, halogen, cyano, nitro,  $\text{COOR}^8$ ,  $\text{XR}^{57}$ ,  $\text{C(O)R}^{57}$ ,  $\text{C(O)NR}^{55}\text{R}^{56}$ , B, D, and E with the proviso that when at least one of  $R^2$  or  $R^3$  is not either heteroaryl or substituted phenyl than  $R_4$  is selected from B or E;

5

m is 2;

$R^5$  is hydrogen;

10  $R^6$  does not exist;

-- represents a carbon-carbon bond or nothing;

A is selected from the group consisting of  $\text{C}_{1-6}$ alkoxy, aryl and heteroaryl; in  
 15 which said aryl is phenyl or said heteroaryl is selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, triazinyl, furanyl, thienyl, pyrrolyl, imidazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, quinolinyl, isoquinolinyl, benzofuranyl, benzothienyl, benzoimidazolyl and benzothiazolyl; and said aryl or heteroaryl is optionally substituted with one or two of the same or different amino, cyano,  
 20  $\text{C}_{1-6}$ alkoxy,  $\text{C}_{1-6}$ alkyl,  $-\text{NHC(O)CH}_3$ , halogen or trifluoromethyl;

B is selected from the group consisting of  $-\text{C(=NR}^{46})(\text{R}^{47})$ ,  $\text{C(O)NR}^{40}\text{R}^{41}$ , aryl, heteroaryl, heteroalicyclic,  $\text{S(O)}_q\text{R}^8$ ,  $\text{P(O)(R}^8)_q(\text{OR}^8)_{2-q}$ ,  $\text{P(S)(R}^8)_q(\text{OR}^8)_{2-q}$ ,  $\text{C(O)R}^8$ ,  $\text{XR}^8$ ,  $(\text{C}_{1-6})\text{alkylNR}^{40}\text{R}^{41}$ , and  $(\text{C}_{1-6})\text{alkylCOOR}^8$  wherein said aryl, heteroaryl, and  
 25 heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

q is 0, 1, or 2;

30 D is selected from the group consisting of  $(\text{C}_{1-6})\text{alkyl}$ ,  $(\text{C}_{3-7})\text{cycloalkyl}$ ,  $(\text{C}_{2-6})\text{alkenyl}$ ,  $(\text{C}_{3-7})\text{cycloalkenyl}$ ,  $(\text{C}_{2-6})\text{alkynyl}$ , wherein said  $(\text{C}_{1-6})\text{alkyl}$ ,  $(\text{C}_{3-7})\text{cycloalkyl}$ ,  $(\text{C}_{2-6})\text{alkenyl}$ ,  $(\text{C}_{3-7})\text{cycloalkenyl}$ , and  $(\text{C}_{2-6})\text{alkynyl}$  are optionally substituted with one to nine same or different halogens or from one to five same or different substituents selected from the group F;

35

E is selected from the group consisting of  $(\text{C}_{1-6})\text{alkyl}$ ,  $(\text{C}_{3-7})\text{cycloalkyl}$ ,  $(\text{C}_{2-6})\text{alkenyl}$ ,  $(\text{C}_{3-7})\text{cycloalkenyl}$ ,  $(\text{C}_{2-6})\text{alkynyl}$ , wherein said  $(\text{C}_{1-6})\text{alkyl}$ ,

(C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, and (C<sub>2-6</sub>)alkynyl are substituted with B;

- F is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl,
- 5 heteroaryl, heteroalicyclic, hydroxy, (C<sub>1-6</sub>)alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, thiohydroxy, (C<sub>1-6</sub>)thioalkoxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, cyano, halogen, nitro, carbonyl, thiocarbonyl, benzyl, O-thiocarbamyl, N-thiocarbamyl, C-thioamido, -NR<sup>42</sup>C(O)-(C<sub>1-6</sub>)alkyl, -NR<sup>42</sup>C(O)-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>42</sup>C(O)-aryl, -NR<sup>42</sup>C(O)-heteroaryl, -NR<sup>42</sup>C(O)-heteroalicyclic, a
- 10 cyclic N-amido, -NR<sup>42</sup>S(O)<sub>2</sub>-(C<sub>1-6</sub>)alkyl, -NR<sup>42</sup>S(O)<sub>2</sub>-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>42</sup>S(O)<sub>2</sub>-aryl, -NR<sup>42</sup>S(O)<sub>2</sub>-heteroaryl, -NR<sup>42</sup>S(O)<sub>2</sub>-heteroalicyclic, sulfinyl, sulfonyl, -S(O)<sub>2</sub>NR<sup>42</sup>R<sup>43</sup>, phosphonyl, NR<sup>42</sup>R<sup>43</sup>, (C<sub>1-6</sub>)alkylC(O)NR<sup>42</sup>R<sup>43</sup>, C(O)NR<sup>42</sup>R<sup>43</sup>, NHC(O)NR<sup>42</sup>R<sup>43</sup>, OC(O)NR<sup>42</sup>R<sup>43</sup>, NHC(O)OR<sup>54</sup>, (C<sub>1-6</sub>)alkylNR<sup>42</sup>R<sup>43</sup>, COOR<sup>54</sup>, and (C<sub>1-6</sub>)alkylCOOR<sup>54</sup> wherein said
- 15 (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, (C<sub>1-6</sub>)alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, (C<sub>1-6</sub>)thioalkoxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;
- 20 G is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, (C<sub>1-6</sub>)alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, thiohydroxy, (C<sub>1-6</sub>)thioalkoxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, cyano, halogen, nitro, carbonyl, thiocarbonyl, benzyl, O-thiocarbamyl, N-thiocarbamyl, C-thioamido, -NR<sup>48</sup>C(O)-(C<sub>1-6</sub>)alkyl, -NR<sup>48</sup>C(O)-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>48</sup>C(O)-aryl, -NR<sup>48</sup>C(O)-heteroaryl, -NR<sup>48</sup>C(O)-heteroalicyclic, a
- 25 cyclic N-amido, -NR<sup>48</sup>S(O)<sub>2</sub>-(C<sub>1-6</sub>)alkyl, -NR<sup>48</sup>S(O)<sub>2</sub>-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>48</sup>S(O)<sub>2</sub>-aryl, -NR<sup>48</sup>S(O)<sub>2</sub>-heteroaryl, -NR<sup>48</sup>S(O)<sub>2</sub>-heteroalicyclic, sulfinyl, sulfonyl, -S(O)<sub>2</sub>NR<sup>48</sup>R<sup>49</sup>, NR<sup>48</sup>R<sup>49</sup>, (C<sub>1-6</sub>)alkylC(O)NR<sup>48</sup>R<sup>49</sup>, C(O)NR<sup>48</sup>R<sup>49</sup>, NHC(O)NR<sup>48</sup>R<sup>49</sup>, OC(O)NR<sup>48</sup>R<sup>49</sup>, NHC(O)OR<sup>54</sup>,
- 30 (C<sub>1-6</sub>)alkylNR<sup>48</sup>R<sup>49</sup>, COOR<sup>54</sup>, and (C<sub>1-6</sub>)alkylCOOR<sup>54</sup>;

- R<sup>7</sup> is selected from the group consisting of aryl, heteroaryl, and heteroalicyclic wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or with from one to two same or different
- 35 substituents selected from the group F;

R<sup>8</sup> is selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, R<sup>18</sup>, R<sup>19</sup> and R<sup>20</sup> are each independently selected from hydrogen or C<sub>1-3</sub>alkyl being optionally substituted with one to three fluorines;

R<sup>23</sup>, R<sup>24</sup>, R<sup>25</sup>, R<sup>26</sup>, R<sup>27</sup>, R<sup>28</sup>, R<sup>29</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, wherein each of said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl being optionally substituted with one to three same or different substituents selected from the group consisting of halogen, hydroxy, cyano, amino and nitro;

X is selected from the group consisting of NR<sup>5</sup>, O, and S;

R<sup>40</sup> and R<sup>41</sup> are independently selected from the group consisting of Hydrogen; or (C<sub>1-6</sub>)alkyl or (C<sub>3-7</sub>)cycloalkyl substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F; or (C<sub>1-6</sub>)alkoxy, aryl, heteroaryl, heteroalicyclic or R<sup>40</sup> and R<sup>41</sup> taken together with the nitrogen to which they are attached form a heteroalicyclic ring which may contain up to 2 additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2; and wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F; with the proviso that only one of R<sup>40</sup> and R<sup>41</sup> may be hydrogen.

R<sup>42</sup> and R<sup>43</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic or R<sup>42</sup> and R<sup>43</sup> taken together with the nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2; and wherein said (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl,

(C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;

5 R<sup>44</sup> is selected from the group consisting of -H

R<sub>a</sub> and R<sub>b</sub> are each independently H, (C<sub>1-6</sub>)alkyl or phenyl;

R<sup>46</sup> is selected from the group consisting of H, OR<sup>8</sup>, and NR<sup>40</sup>R<sup>41</sup>;

10 R<sup>47</sup> is selected from the group consisting of H, amino, halogen, and (C<sub>1-6</sub>)alkyl;

R<sup>48</sup> and R<sup>49</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, allyl, aryl, heteroaryl, heteroalicyclic or R<sup>48</sup> and R<sup>49</sup> taken together with the nitrogen to which they are attached form a heteroaryl  
15 ring or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2;

R<sup>50</sup> is selected from the group consisting of H, (C<sub>1-6</sub>)alkyl, (C<sub>3-6</sub>)cycloalkyl, and benzyl, each of said alkyl, cycloalkyl and benzyl being optionally substituted with  
20 one to three same or different halogen, amino, OH, CN or NO<sub>2</sub>;

R<sup>51</sup> is selected from the group consisting of H, (C<sub>1-6</sub>)alkyl, (C<sub>3-6</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-6</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl or C(O)R<sup>53</sup>, wherein R<sup>53</sup> is H, (C<sub>1-6</sub>)alkyl, or (C<sub>3-6</sub>)cycloalkyl and each of said (C<sub>1-6</sub>)alkyl and (C<sub>3-6</sub>)cycloalkyl being  
25 optionally substituted with one to three same or different halogen, amino, OH, CN or NO<sub>2</sub>;

Y is O, S or NR<sup>50</sup>R<sup>51</sup>;

30 R<sup>54</sup> is selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, allyl, aryl, heteroaryl, and heteroalicyclic wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group consisting of: amino, OH, and NR<sup>55</sup>R<sup>56</sup>;

35 R<sup>54'</sup> is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, allyl, aryl, heteroaryl, and heteroalicyclic wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl,

heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group consisting of: amino, OH, and  $\text{NR}^{55}\text{R}^{56}$ ;

- 5  $\text{R}^{55}$  and  $\text{R}^{56}$  are independently selected from the group consisting of hydrogen,  $(\text{C}_{1-6})$ alkyl, allyl, or  $(\text{C}_{3-7})$ cycloalkyl; and

$\text{R}^{57}$  is selected from the group consisting of hydrogen,  $(\text{C}_{1-6})$ alkyl,  $(\text{C}_{3-7})$ cycloalkyl,  $(\text{C}_{2-6})$ alkenyl,  $(\text{C}_{3-7})$ cycloalkenyl,  $(\text{C}_{2-6})$ alkynyl.

10

An even more preferred embodiment of a first aspect of the invention are compounds of Formula I, including pharmaceutically acceptable salts thereof,

- 15 A is selected from the group consisting of phenyl and heteroaryl in which said heteroaryl is selected from pyridinyl, furanyl and thienyl, and said phenyl or said heteroaryl is optionally substituted with one to two of the same or different amino,  $\text{C}_{1-6}$ alkyl, or halogen;

- - represents a carbon-carbon bond;


20

$\text{R}^9$ ,  $\text{R}^{10}$ ,  $\text{R}^{11}$ ,  $\text{R}^{12}$ ,  $\text{R}^{13}$ , and  $\text{R}^{14}$  are each hydrogen; and

$\text{R}^{15}$  and  $\text{R}^{16}$  are each independently hydrogen or methyl with the proviso that only one is methyl.

25


Q is either



- 30 and then  $\text{R}^2$  is selected from the group consisting of hydrogen, halogen and methoxy; and

$\text{R}_3$  is hydrogen;

Or Q is:



and R<sup>2</sup> is halogen or hydrogen and R<sup>3</sup> is hydrogen;

5 R<sup>4</sup> is selected from the group consisting of B or E

B is selected from the group consisting of -C(O)NR<sup>40</sup>R<sup>41</sup>, substituted phenyl, heteroaryl, and C(O)R<sup>7</sup> wherein said heteroaryl is optionally substituted and phenyl is substituted with one to three same or different halogens or from one to two same or  
10 different substituents selected from the group F;

E is selected from the group consisting of (C<sub>2</sub>)alkenyl, or (C<sub>2</sub>)alkynyl, wherein (C<sub>2-6</sub>)alkenyl or (C<sub>2</sub>)alkynyl are substituted with B;

15 F is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-6</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, (C<sub>1-6</sub>)alkoxy, (C<sub>1-6</sub>)thioalkoxy, cyano, halogen, carbonyl, benzyl, -NR<sup>42</sup>C(O)-(C<sub>1-6</sub>)alkyl, -NR<sup>42</sup>C(O)-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>42</sup>C(O)-aryl, -NR<sup>42</sup>C(O)-heteroaryl, -NR<sup>42</sup>C(O)-heteroalicyclic, a cyclic N-amido, -NR<sup>42</sup>S(O)<sub>2</sub>-(C<sub>1-6</sub>)alkyl, -NR<sup>42</sup>S(O)<sub>2</sub>-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>42</sup>S(O)<sub>2</sub>-aryl, -NR<sup>42</sup>S(O)<sub>2</sub>-heteroaryl, -  
20 NR<sup>42</sup>S(O)<sub>2</sub>-heteroalicyclic, -S(O)<sub>2</sub>NR<sup>42</sup>R<sup>43</sup>, NR<sup>42</sup>R<sup>43</sup>, (C<sub>1-6</sub>)alkylC(O)NR<sup>42</sup>R<sup>43</sup>, C(O)NR<sup>42</sup>R<sup>43</sup>, NHC(O)NR<sup>42</sup>R<sup>43</sup>, OC(O)NR<sup>42</sup>R<sup>43</sup>, NHC(O)OR<sup>54</sup>, (C<sub>1-6</sub>)alkylNR<sup>42</sup>R<sup>43</sup>, COOR<sup>54</sup>, and (C<sub>1-6</sub>)alkylCOOR<sup>54</sup> wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-6</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, (C<sub>1-6</sub>)alkoxy, are optionally substituted with one to three same or different halogens or from one to two  
25 same or different substituents selected from the group G;

G is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, (C<sub>1-6</sub>)alkoxy, (C<sub>1-6</sub>)thioalkoxy, thioaryloxy, cyano, halogen, nitro, carbonyl, thiocarbonyl, benzyl, -NR<sup>48</sup>C(O)-(C<sub>1-6</sub>)alkyl, -NR<sup>48</sup>C(O)-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>48</sup>C(O)-aryl, -NR<sup>48</sup>C(O)-heteroaryl, -NR<sup>48</sup>C(O)-heteroalicyclic, a cyclic N-amido, -NR<sup>48</sup>S(O)<sub>2</sub>-(C<sub>1-6</sub>)alkyl, -NR<sup>48</sup>S(O)<sub>2</sub>-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>48</sup>S(O)<sub>2</sub>-aryl, -NR<sup>48</sup>S(O)<sub>2</sub>-heteroaryl, -NR<sup>48</sup>S(O)<sub>2</sub>-heteroalicyclic, sulfonyl, -S(O)<sub>2</sub>NR<sup>48</sup>R<sup>49</sup>, NR<sup>48</sup>R<sup>49</sup>, (C<sub>1-6</sub>)alkyl C(O)NR<sup>48</sup>R<sup>49</sup>, C(O)NR<sup>48</sup>R<sup>49</sup>, NHC(O)NR<sup>48</sup>R<sup>49</sup>, OC(O)NR<sup>48</sup>R<sup>49</sup>, NHC(O)OR<sup>54</sup>,  
30 (C<sub>1-6</sub>)alkylNR<sup>48</sup>R<sup>49</sup>, COOR<sup>54</sup>, and (C<sub>1-6</sub>)alkylCOOR<sup>54</sup>;

R<sup>7</sup> is selected from the group consisting of aryl, heteroaryl, and heteroalicyclic wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or with from one to two same or different substituents selected from the group F;

R<sup>8</sup> is selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, and (C<sub>3-7</sub>)cycloalkyl, wherein (C<sub>1-6</sub>)alkyl, and (C<sub>3-7</sub>)cycloalkyl are optionally substituted with one to six same or different halogens or from one to two same or different substituents selected from the group F;

R<sup>40</sup> and R<sup>41</sup> are independently selected from the group consisting of Hydrogen; or (C<sub>1-6</sub>)alkyl or (C<sub>3-7</sub>)cycloalkyl substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F; or (C<sub>1-6</sub>)alkoxy, aryl, heteroaryl, heteroalicyclic or R<sup>40</sup> and R<sup>41</sup> taken together with the nitrogen to which they are attached form a heteroalicyclic ring which may contain up to 2 additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2; and wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F; with the proviso that only one of R<sup>40</sup> and R<sup>41</sup> may be hydrogen.

R<sup>42</sup> and R<sup>43</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic or R<sup>42</sup> and R<sup>43</sup> taken together with the nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2; and wherein said (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;

R<sup>44</sup> is selected from the group consisting of -H;

R<sup>48</sup> and R<sup>49</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic or R<sup>48</sup> and R<sup>49</sup> taken together with the nitrogen to which they are attached form a heteroaryl ring



or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, O, S(O)<sub>m'</sub>, wherein m' is 0, 1, or 2;

- R<sup>54</sup> is selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, and heteroalicyclic wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group consisting of: amino, OH, and NR<sup>55</sup>R<sup>56</sup>;
- R<sup>54'</sup> is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, and heteroalicyclic wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group consisting of: amino, OH, and NR<sup>55</sup>R<sup>56</sup>;
- R<sup>55</sup> and R<sup>56</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, or (C<sub>3-7</sub>)cycloalkyl

- Among the preferred compounds of the first embodiment of a first aspect of the invention are compounds of Formula I, including pharmaceutically acceptable salts thereof,

- R<sup>4</sup> is selected from the group consisting of B;
- B is selected from the group consisting of -C(O)NR<sup>40</sup>R<sup>41</sup>, substituted phenyl, or heteroaryl, wherein said phenyl is substituted and heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

- F is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-6</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, (C<sub>1-6</sub>)alkoxy, (C<sub>1-6</sub>)thioalkoxy, cyano, halogen, carbonyl, benzyl, -NR<sup>42</sup>C(O)-(C<sub>1-6</sub>)alkyl, -NR<sup>42</sup>C(O)-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>42</sup>C(O)-aryl, -NR<sup>42</sup>C(O)-heteroaryl, -NR<sup>42</sup>C(O)-heteroalicyclic, a cyclic N-amido, -NR<sup>42</sup>S(O)<sub>2</sub>-(C<sub>1-6</sub>)alkyl, -NR<sup>42</sup>R<sup>43</sup>, C(O)NR<sup>42</sup>R<sup>43</sup>, COOR<sup>54</sup>, and wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-6</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, (C<sub>1-6</sub>)alkoxy, are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;

G is selected from the group consisting of (C<sub>1-6</sub>)alkyl, hydroxy, (C<sub>1-6</sub>)alkoxy, halogen, -NR<sup>48</sup>C(O)-(C<sub>1-6</sub>)alkyl, -NR<sup>48</sup>C(O)-(C<sub>3</sub>)cycloalkyl, a cyclic N-amido, -NR<sup>48</sup>S(O)<sub>2</sub>-(C<sub>1-6</sub>)alkyl, NR<sup>48</sup>R<sup>49</sup>, (C<sub>1-6</sub>)alkyl C(O)NR<sup>48</sup>R<sup>49</sup>, C(O)NR<sup>48</sup>R<sup>49</sup>, (C<sub>1-6</sub>)alkylNR<sup>48</sup>R<sup>49</sup>;

5

R<sup>40</sup> is Hydrogen;

R<sup>41</sup> is (C<sub>1-3</sub>)alkoxy, heteroaryl, or aryl, wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;

10

R<sup>42</sup> and R<sup>43</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic or R<sup>42</sup> and R<sup>43</sup> taken together with the nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2; and wherein said (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;

15

20


R<sup>48</sup> and R<sup>49</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl or R<sup>48</sup> and R<sup>49</sup> taken together with the nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, or O;

25

A second group of preferred compounds of Formula I, including pharmaceutically acceptable salts thereof,

Q is

30



R<sup>4</sup> is B;

A is Phenyl or 2-pyridyl;

- B is selected from the group consisting of  $-C(O)NR^{40}R^{41}$  or heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

Most preferred among this second group of preferred compounds are those where  $R^4$  is B;

- 10 A is Phenyl or 2-pyridyl and B is selected from the group consisting of  $-C(O)NR^{40}R^{41}$  or heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group;
- 15 Compounds where B is heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F claimed;

- Preferred groups for B when B is heteroaryl are selected from the group consisting of thiazole, pyridazine, pyrazine, pyrazole, isoxazole, isothiazole, imidazole, furyl, thienyl, oxazole, oxadiazole, thiadiazole, pyrimidine, pyrazole, triazine, triazole, tetrazole, pyridyl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

- 25 When B is heteroaryl, most preferred is when said heteroaryl is optionally substituted with one to three same or different halogens or a substituent selected from the group  $(C_1-C_6 \text{ alkyl})$ , amino,  $-NHC(O)-(C_1-C_6 \text{ alkyl})$ ,  $-NHS(O)_2-(C_1-C_6 \text{ alkyl})$ , methoxy,  $-C(O)-NH_2$ ,  $C(O)NHMe$ ,  $C(O)NMe_2$ , trifluoromethyl,  $-NHC(C_1-C_6 \text{ alkyl})$ ,  $-N(C_1-C_6 \text{ alkyl})_2$ , -heteroaryl, cyclic N-amido; among the most preferred B is thienyl and when B is thienyl most preferred is when the thienyl is optionally substituted with one to three same or different halogens or a substituent selected from the group  $(C_1-C_6 \text{ alkyl})$ , amino,  $-NHC(O)-(C_1-C_6 \text{ alkyl})$ ,  $-NHS(O)_2-(C_1-C_6 \text{ alkyl})$ , methoxy,  $-C(O)-NH_2$ ,  $C(O)NHMe$ ,  $C(O)NMe_2$ , trifluoromethyl,  $-NHC(C_1-C_6 \text{ alkyl})$ ,  $-N(C_1-C_6 \text{ alkyl})_2$ , -heteroaryl, cyclic N-amido;
- 35 and even more preferred is when the thienyl is optionally substituted with one to three same or different halogens or a substituent selected from the group  $(C_1-C_6$

alkyl), amino,  $-\text{NHC(O)}-(\text{C}_1\text{-C}_6 \text{ alkyl})$ ,  $-\text{NHS(O)}_2-(\text{C}_1\text{-C}_6 \text{ alkyl})$ , methoxy,  $-\text{C(O)}-\text{NH}_2$ ,  $\text{C(O)NHMe}$ ,  $\text{C(O)NMe}_2$ , trifluoromethyl,  $-\text{NHC}(\text{C}_1\text{-C}_6 \text{ alkyl})$ ,  $-\text{N}(\text{C}_1\text{-C}_6 \text{ alkyl})_2$ , -heteroaryl, cyclic N-amido;

- 5 when B is selected from the group consisting of  $-\text{C(O)NR}^{40}\text{R}^{41}$  or  $-\text{C(O)NH}$ -heteroaryl is preferred wherein said heteroaryl is optionally substituted with one to three same or different halogens or a substituent selected from the group  $(\text{C}_1\text{-C}_6 \text{ alkyl})$ , amino,  $-\text{NHC(O)}-(\text{C}_1\text{-C}_6 \text{ alkyl})$ , -methoxy,  $-\text{NHC}(\text{C}_1\text{-C}_6 \text{ alkyl})$ , or  $-\text{N}(\text{C}_1\text{-C}_6 \text{ alkyl})_2$ ;

10

A third group of preferred compounds of Formula I are, including pharmaceutically acceptable salts thereof wherein,

Q is

15



$\text{R}^2$  is selected from the group consisting of hydrogen, halogen, and methoxy;

$\text{R}^4$  is B;

20

B is selected from the group consisting of  $-\text{C(O)NR}^{40}\text{R}^{41}$  or heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

- 25 Most preferred are compounds where A is Phenyl or 2-pyridyl;

Most preferred for B is as described above.

- 30 Preferred groups for B when B is heteroaryl are selected from the group consisting of thiazole, pyridazine, pyrazine, pyrazole, isoxazole, isothiazole, imidazole, furyl, thienyl, oxazole, oxadiazole, thiadiazole, pyrimidine, pyrazole, triazine, triazole, tetrazole, pyridyl, wherein said heteroaryl is optionally substituted with one to three

same or different halogens or from one to two same or different substituents selected from the group F;

- When B is heteroaryl, most preferred is when said heteroaryl is optionally substituted with one to three same or different halogens or a substituent selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NHS(O)<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), methoxy, -C(O)-NH<sub>2</sub>, C(O)NHMe, C(O)NMe<sub>2</sub>, trifluoromethyl, -NHC(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -heteroaryl, cyclic N-amido; among the most preferred B is thienyl, pyrazole, or a six membered heteroaryl containing two ring nitrogens.
- and when B is one of these most preferred groups it is optionally substituted with one to three same or different halogens or a substituent selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NHS(O)<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), methoxy, -C(O)-NH<sub>2</sub>, C(O)NHMe, C(O)NMe<sub>2</sub>, trifluoromethyl, -NHC(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -heteroaryl, cyclic N-amido;
- and even more preferred is when said heteroaryl is optionally substituted with one to three same or different halogens or a substituent selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NHS(O)<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), methoxy, -C(O)-NH<sub>2</sub>, C(O)NHMe, C(O)NMe<sub>2</sub>, trifluoromethyl, -NHC(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -heteroaryl, cyclic N-amido;

Another embodiment of a preferred aspect of the invention are compounds of Formula I, including pharmaceutically acceptable salts thereof,


- A is selected from the group consisting of phenyl and heteroaryl in which said heteroaryl is selected from pyridinyl, furanyl and thienyl, and said phenyl or said heteroaryl is optionally substituted with one to two of the same or different amino, C<sub>1-6</sub>alkyl, or halogen;

- - represents a carbon-carbon bond;


R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup>, and R<sup>14</sup> are each hydrogen; and

- R<sup>15</sup> and R<sup>16</sup> are each independently hydrogen or methyl with the proviso that only one is methyl.

Q is either



or




and then  $R^2$  is selected from the group consisting of hydrogen, halogen and methoxy;

5

And  $R_3$  is hydrogen;

Or Q is:



10

and  $R^2$  is halogen or hydrogen and  $R^3$  is hydrogen;

$R^4$  is B; and

- 15 F is selected from the group consisting of  $(C_{1-6})$ alkyl, hydroxy, heteroaryl, heteroalicyclic, methoxy, methylthioalkoxy, halogen, carbonyl,  $C(O)NR^{42}R^{43}$ ,  $-NR^{42}C(O)-(C_{1-6})$ alkyl,  $-NR^{42}C(O)-(C_{3-6})$ cycloalkyl,  $-NR^{42}C(O)$ -aryl,  $-NR^{42}C(O)$ -heteroaryl,  $-NR^{42}C(O)$ -heteroalicyclic, a cyclic N-amido,  $-NR^{42}S(O)_2-(C_{1-6})$ alkyl,  $-NR^{42}S(O)_2-(C_{3-6})$ cycloalkyl,  $-NR^{42}S(O)_2$ -aryl,  $-NR^{42}S(O)_2$ -heteroaryl,  $-NR^{42}S(O)_2$ -heteroalicyclic,  $NR^{42}R^{43}$ ,  $COOH$
- 20

- G is selected from the group consisting of  $(C_{1-6})$ alkyl,  $(C_{3-7})$ cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy,  $(C_{1-6})$ alkoxy,  $(C_{1-6})$ thioalkoxy, thioaryloxy, cyano, halogen, nitro, carbonyl, thiocarbonyl, benzyl,  $-NR^{48}C(O)-(C_{1-6})$ alkyl,  $-NR^{48}C(O)-(C_{3-6})$ cycloalkyl,  $-NR^{48}C(O)$ -aryl,  $-NR^{48}C(O)$ -heteroaryl,  $-NR^{48}C(O)$ -heteroalicyclic, a cyclic N-amido,  $-NR^{48}S(O)_2-(C_{1-6})$ alkyl,  $-NR^{48}S(O)_2-(C_{3-6})$ cycloalkyl,  $-NR^{48}S(O)_2$ -aryl,  $-NR^{48}S(O)_2$ -heteroaryl,  $-NR^{48}S(O)_2$ -heteroalicyclic, sulfonyl,  $-S(O)_2NR^{48}R^{49}$ ,  $NR^{48}R^{49}$ ,  $(C_{1-6})$ alkyl  $C(O)NR^{48}R^{49}$ ,  $C(O)NR^{48}R^{49}$ ,  $NHC(O)NR^{48}R^{49}$ ,  $OC(O)NR^{48}R^{49}$ ,  $NHC(O)OR^{54'}$ ,  $(C_{1-6})$ alkyl  $NR^{48}R^{49}$ ,  $COOR^{54}$ , and  $(C_{1-6})$ alkyl  $COOR^{54'}$ ;
- 25
- 30

R<sup>7</sup> is selected from the group consisting of aryl, heteroaryl, and heteroalicyclic wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or with from one to two same or different substituents selected from the group F;

5

R<sup>8</sup> is selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, and (C<sub>3-7</sub>)cycloalkyl, wherein (C<sub>1-6</sub>)alkyl, and (C<sub>3-7</sub>)cycloalkyl are optionally substituted with one to six same or different halogens or from one to two same or different substituents selected from the group F;

10

R<sup>40</sup> and R<sup>41</sup> are independently selected from the group consisting of Hydrogen;

or (C<sub>1-6</sub>)alkyl or (C<sub>3-7</sub>)cycloalkyl substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

15

or (C<sub>1-6</sub>)alkoxy, aryl, heteroaryl, heteroalicyclic or R<sup>40</sup> and R<sup>41</sup> taken together with the nitrogen to which they are attached form a heteroalicyclic ring which may contain up to 2 additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2; and wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

20

with the proviso that only one of R<sup>40</sup> and R<sup>41</sup> may be hydrogen.

R<sup>42</sup> and R<sup>43</sup> are independently selected from the group consisting of hydrogen,

(C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic or R<sup>42</sup> and

25

R<sup>43</sup> taken together with the nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2; and wherein said (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl,

(C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;

30

R<sup>44</sup> is selected from the group consisting of -H

35

R<sup>48</sup> and R<sup>49</sup> are independently selected from the group consisting of hydrogen,

(C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic or R<sup>48</sup> and


R<sup>49</sup> taken together with the nitrogen to which they are attached form a heteroaryl ring

or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2;

- 5 R<sup>54</sup> is selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, and heteroalicyclic wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group consisting of: amino, OH, and NR<sup>55</sup>R<sup>56</sup>;
- 10 R<sup>54'</sup> is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, and heteroalicyclic wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group consisting of: amino, OH, and NR<sup>55</sup>R<sup>56</sup>;
- 15 R<sup>55</sup> and R<sup>56</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, or (C<sub>3-7</sub>)cycloalkyl

A fourth group of preferred compounds is those wherein:

20 Q is



R<sup>2</sup> is selected from the group consisting of hydrogen or methoxy;

25 R<sup>3</sup> is hydrogen;

R<sup>4</sup> is B

30 B is selected from the group consisting of -C(O)NR<sup>40</sup>R<sup>41</sup> or heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;



A final preferred aspect of the invention are compounds depicted in Table 2 or Table 4 of the biology section.

5 A second embodiment of the third aspect of the present invention is a method for treating mammals infected with a virus, wherein said virus is HIV, comprising administering to said mammal an antiviral effective amount of a compound of Formula I.

10 A third embodiment of the third aspect of the present invention is a method for treating mammals infected with a virus, such as HIV, comprising administering to said mammal an antiviral effective amount of a compound of Formula I in combination with an antiviral effective amount of an AIDS treatment agent selected from the group consisting of: (a) an AIDS antiviral agent; (b) an anti-infective agent; (c) an immunomodulator; and (d) HIV entry inhibitors.

### DETAILED DESCRIPTION OF THE INVENTION

15

Since the compounds of the present invention, may possess asymmetric centers and therefore occur as mixtures of diastereomers and enantiomers, the present invention includes the individual diastereoisomeric and enantiomeric forms of the compounds of Formula I in addition to the mixtures thereof.

20

### DEFINITIONS

25 The term "C<sub>1-6</sub> alkyl" as used herein and in the claims (unless specified otherwise) mean straight or branched chain alkyl groups such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, amyl, hexyl and the like.

"Halogen" refers to chlorine, bromine, iodine or fluorine.

30 An "aryl" group refers to an all carbon monocyclic or fused-ring polycyclic (i.e., rings which share adjacent pairs of carbon atoms) groups having a completely conjugated pi-electron system. Examples, without limitation, of aryl groups are phenyl, naphthalenyl and anthracenyl. The aryl group may be substituted or unsubstituted. When substituted the substituted group(s) is preferably one or more selected from alkyl, cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, alkoxy, 35 aryloxy, heteroaryloxy, heteroalicycloxy, thiohydroxy, thioaryloxy,

thioheteroaryloxy, thioheteroalicycloxy, cyano, halogen, nitro, carbonyl, O-carbamyl, N-carbamyl, C-amido, N-amido, C-carboxy, O-carboxy, sulfinyl, sulfonyl, sulfonamido, trihalomethyl, ureido, amino and  $-NR^xR^y$ , wherein  $R^x$  and  $R^y$  are independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, aryl, carbonyl, C-carboxy, sulfonyl, trihalomethyl, and, combined, a five- or six-member heteroalicyclic ring.

As used herein, a "heteroaryl" group refers to a monocyclic or fused ring (i.e., rings which share an adjacent pair of atoms) group having in the ring(s) one or more atoms selected from the group consisting of nitrogen, oxygen and sulfur and, in addition, having a completely conjugated pi-electron system. It should be noted that the term heteroaryl is intended to encompass an N-oxide of the parent heteroaryl if such an N-oxide is chemically feasible as is known in the art. Examples, without limitation, of heteroaryl groups are furyl, thienyl, benzothienyl, thiazolyl, imidazolyl, oxazolyl, oxadiazolyl, thiadiazolyl, benzothiazolyl, triazolyl, tetrazolyl, isoxazolyl, isothiazolyl, pyrrolyl, pyranal, tetrahydropyranal, pyrazolyl, pyridyl, pyrimidinyl, quinolinyl, isoquinolinyl, purinyl, carbazolyl, benzoxazolyl, benzimidazolyl, indolyl, isoindolyl, pyrazinyl, diazinyl, pyrazine, triazinyltriazine, tetrazinyl, and tetrazolyl. When substituted the substituted group(s) is preferably one or more selected from alkyl, cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, thiohydroxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, cyano, halogen, nitro, carbonyl, O-carbamyl, N-carbamyl, C-amido, N-amido, C-carboxy, O-carboxy, sulfinyl, sulfonyl, sulfonamido, trihalomethyl, ureido, amino, and  $-NR^xR^y$ , wherein  $R^x$  and  $R^y$  are as defined above.

As used herein, a "heteroalicyclic" group refers to a monocyclic or fused ring group having in the ring(s) one or more atoms selected from the group consisting of nitrogen, oxygen and sulfur. The rings may also have one or more double bonds. However, the rings do not have a completely conjugated pi-electron system. Examples, without limitation, of heteroalicyclic groups are azetidiny, piperidyl, piperazinyl, imidazolinyl, thiazolidinyl, 3-pyrrolidin-1-yl, morpholinyl, thiomorpholinyl and tetrahydropyranal. When substituted the substituted group(s) is preferably one or more selected from alkyl, cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, thiohydroxy, thioalkoxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, cyano, halogen, nitro, carbonyl, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, C-thioamido, N-amido, C-carboxy, O-carboxy, sulfinyl, sulfonyl, sulfonamido, trihalomethanesulfonamido, trihalomethanesulfonyl, silyl,

guanyl, guanidino, ureido, phosphonyl, amino and  $-NR^xR^y$ , wherein  $R^x$  and  $R^y$  are as defined above.

5 An "alkyl" group refers to a saturated aliphatic hydrocarbon including straight chain and branched chain groups. Preferably, the alkyl group has 1 to 20 carbon atoms (whenever a numerical range; e.g., "1-20", is stated herein, it means that the group, in this case the alkyl group may contain 1 carbon atom, 2 carbon atoms, 3 carbon atoms, etc. up to and including 20 carbon atoms). More preferably, it is a medium size alkyl having 1 to 10 carbon atoms. Most preferably, it is a lower alkyl  
10 having 1 to 4 carbon atoms. The alkyl group may be substituted or unsubstituted. When substituted, the substituent group(s) is preferably one or more individually selected from trihaloalkyl, cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, thiohydroxy, thioalkoxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, cyano, halo, nitro, carbonyl, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido,  
15 C-thioamido, N-amido, C-carboxy, O-carboxy, sulfinyl, sulfonyl, sulfonamido, trihalomethanesulfonamido, trihalomethanesulfonyl, and combined, a five- or six-member heteroalicyclic ring.

20 A "cycloalkyl" group refers to an all-carbon monocyclic or fused ring (i.e., rings which share and adjacent pair of carbon atoms) group wherein one or more rings does not have a completely conjugated pi-electron system. Examples, without limitation, of cycloalkyl groups are cyclopropane, cyclobutane, cyclopentane, cyclopentene, cyclohexane, cyclohexadiene, cycloheptane, cycloheptatriene and  
25 adamantane. A cycloalkyl group may be substituted or unsubstituted. When substituted, the substituent group(s) is preferably one or more individually selected from alkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, thiohydroxy, thioalkoxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, cyano, halo, nitro, carbonyl, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, C-thioamido, N-amido, C-carboxy, O-carboxy, sulfinyl, sulfonyl, sulfonamido, trihalo- methanesulfonamido, trihalomethanesulfonyl, silyl, guanyl, guanidino, ureido, phosphonyl, amino and -  
30  $NR^xR^y$  with  $R^x$  and  $R^y$  as defined above.

35 An "alkenyl" group refers to an alkyl group, as defined herein, consisting of at least two carbon atoms and at least one carbon-carbon double bond.

An "alkynyl" group refers to an alkyl group, as defined herein, consisting of at least two carbon atoms and at least one carbon-carbon triple bond.

A "hydroxy" group refers to an -OH group.

5

An "alkoxy" group refers to both an -O-alkyl and an -O-cycloalkyl group as defined herein.

10 An "aryloxy" group refers to both an -O-aryl and an -O-heteroaryl group, as defined herein.

A "heteroaryloxy" group refers to a heteroaryl-O- group with heteroaryl as defined herein.

15 A "heteroalicycloxy" group refers to a heteroalicyclic-O- group with heteroalicyclic as defined herein.

A "thiohydroxy" group refers to an -SH group.

20 A "thioalkoxy" group refers to both an S-alkyl and an -S-cycloalkyl group, as defined herein.

A "thioaryloxy" group refers to both an -S-aryl and an -S-heteroaryl group, as defined herein.

25

A "thioheteroaryloxy" group refers to a heteroaryl-S- group with heteroaryl as defined herein.

30 A "thioheteroalicycloxy" group refers to a heteroalicyclic-S- group with heteroalicyclic as defined herein.

35 A "carbonyl" group refers to a -C(=O)-R" group, where R" is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and heteroalicyclic (bonded through a ring carbon), as each is defined herein.

An "aldehyde" group refers to a carbonyl group where R" is hydrogen.

A "thiocarbonyl" group refers to a  $-C(=S)-R$  group, with R" as defined herein.

5 A "Keto" group refers to a  $-CC(=O)C-$  group wherein the carbon on either or both sides of the  $C=O$  may be alkyl, cycloalkyl, aryl or a carbon of a heteroaryl or heteroaliacyclic group.

10 A "trihalomethanecarbonyl" group refers to a  $Z_3CC(=O)-$  group with said Z being a halogen.

A "C-carboxy" group refers to a  $-C(=O)O-R$  groups, with R" as defined herein.

15 An "O-carboxy" group refers to a  $R"C(-O)O-$  group, with R" as defined herein.

A "carboxylic acid" group refers to a C-carboxy group in which R" is hydrogen.

20 A "trihalomethyl" group refers to a  $-CZ_3$  group wherein Z is a halogen group as defined herein.

25 A "trihalomethanesulfonyl" group refers to an  $Z_3CS(=O)_2-$  groups with Z as defined above.

A "trihalomethanesulfonamido" group refers to a  $Z_3CS(=O)_2NR^x-$  group with Z and  $R^x$  as defined herein.

30 A "sulfinyl" group refers to a  $-S(=O)-R$  group, with R" as defined herein and, in addition, as a bond only; i.e.,  $-S(O)-$ .

A "sulfonyl" group refers to a  $-S(=O)_2R$  group with R" as defined herein and, in addition as a bond only; i.e.,  $-S(O)_2-$ .

35 A "S-sulfonamido" group refers to a  $-S(=O)_2NR^xR^y$ , with  $R^x$  and  $R^y$  as defined herein.

A "N-Sulfonamido" group refers to a  $R''S(=O)_2NR_x-$  group with  $R_x$  as defined herein.

A "O-carbamyl" group refers to a  $-OC(=O)NR^xR^y$  as defined herein.

5

A "N-carbamyl" group refers to a  $R^xOC(=O)NR^y$  group, with  $R^x$  and  $R^y$  as defined herein.

10 A "O-thiocarbamyl" group refers to a  $-OC(=S)NR^xR^y$  group with  $R^x$  and  $R^y$  as defined herein.

A "N-thiocarbamyl" group refers to a  $R^xOC(=S)NR^y-$  group with  $R^x$  and  $R^y$  as defined herein.

15 An "amino" group refers to an  $-NH_2$  group.

A "C-amido" group refers to a  $-C(=O)NR^xR^y$  group with  $R^x$  and  $R^y$  as defined herein.

20 A "C-thioamido" group refers to a  $-C(=S)NR^xR^y$  group, with  $R^x$  and  $R^y$  as defined herein.

A "N-amido" group refers to a  $R^xC(=O)NR^y-$  group, with  $R^x$  and  $R^y$  as defined herein.

25

An "ureido" group refers to a  $-NR^xC(=O)NR^yR^{y2}$  group with  $R^x$  and  $R^y$  as defined herein and  $R^{y2}$  defined the same as  $R^x$  and  $R^y$ .

30 A "guanidino" group refers to a  $-R^xNC(=N)NR^yR^{y2}$  group, with  $R^x$ ,  $R^y$  and  $R^{y2}$  as defined herein.

A "guanyl" group refers to a  $R^xR^yNC(=N)-$  group, with  $R^x$  and  $R^y$  as defined herein.

35 A "cyano" group refers to a  $-CN$  group.

A "silyl" group refers to a  $-Si(R'')_3$ , with  $R''$  as defined herein.

A "phosphonyl" group refers to a  $P(=O)(OR^x)_2$  with  $R^x$  as defined herein.

A "hydrazino" group refers to a  $-NR^xNR^yR^{y2}$  group with  $R^x$ ,  $R^y$  and  $R^{y2}$  as defined herein.

5

Any two adjacent R groups may combine to form an additional aryl, cycloalkyl, heteroaryl or heterocyclic ring fused to the ring initially bearing those R groups.

10

It is known in the art that nitrogen atoms in heteroaryl systems can be "participating in a heteroaryl ring double bond", and this refers to the form of double bonds in the two tautomeric structures which comprise five-member ring heteroaryl groups. This dictates whether nitrogens can be substituted as well understood by chemists in the art. The disclosure and claims of the present invention are based on the known general principles of chemical bonding. It is understood that the claims do not encompass structures known to be unstable or not able to exist based on the literature.

15

Physiologically acceptable salts and prodrugs of compounds disclosed herein are within the scope of this invention. The term "pharmaceutically acceptable salt" as used herein and in the claims is intended to include nontoxic base addition salts. Suitable salts include those derived from organic and inorganic acids such as, without limitation, hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, methanesulfonic acid, acetic acid, tartaric acid, lactic acid, sulfinic acid, citric acid, maleic acid, fumaric acid, sorbic acid, aconitic acid, salicylic acid, phthalic acid, and the like. The term "pharmaceutically acceptable salt" as used herein is also intended to include salts of acidic groups, such as a carboxylate, with such counterions as ammonium, alkali metal salts, particularly sodium or potassium, alkaline earth metal salts, particularly calcium or magnesium, and salts with suitable organic bases such as lower alkylamines (methylamine, ethylamine, cyclohexylamine, and the like) or with substituted lower alkylamines (e.g. hydroxyl-substituted alkylamines such as diethanolamine, triethanolamine or tris(hydroxymethyl)-aminomethane), or with bases such as piperidine or morpholine.

25

30

35

In the method of the present invention, the term "antiviral effective amount" means the total amount of each active component of the method that is sufficient to show a meaningful patient benefit, i.e., healing of acute conditions characterized by inhibition of the HIV infection. When applied to an individual active ingredient,

administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously. The terms "treat, treating, treatment" as used herein and in the  
5 claims means preventing or ameliorating diseases associated with HIV infection.

The present invention is also directed to combinations of the compounds with one or more agents useful in the treatment of AIDS. For example, the compounds of this invention may be effectively administered, whether at periods of pre-exposure  
10 and/or post-exposure, in combination with effective amounts of the AIDS antivirals, immunomodulators, antiinfectives, or vaccines, such as those in the following table.

#### ANTIVIRALS

15	<u>Drug Name</u>	<u>Manufacturer</u>	<u>Indication</u>
20	097	Hoechst/Bayer	HIV infection, AIDS, ARC (non-nucleoside reverse trans- criptase (RT) inhibitor)
25	Amprenivir 141 W94 GW 141	Glaxo Wellcome	HIV infection, AIDS, ARC (protease inhibitor)
30	Abacavir (1592U89) GW 1592	Glaxo Wellcome	HIV infection, AIDS, ARC (RT inhibitor)
	Acemannan	Carrington Labs (Irving, TX)	ARC
35	Acyclovir	Burroughs Wellcome	HIV infection, AIDS, ARC, in combination with AZT
40	AD-439	Tanox Biosystems	HIV infection, AIDS, ARC



	AD-519	Tanox Biosystems	HIV infection, AIDS, ARC
5	Adefovir dipivoxil AL-721	Gilead Sciences Ethigen (Los Angeles, CA)	HIV infection ARC, PGL HIV positive, AIDS
10	Alpha Interferon	Glaxo Wellcome	Kaposi's sarcoma, HIV in combination w/Retrovir
15	Ansamycin LM 427	Adria Laboratories (Dublin, OH) Erbamont (Stamford, CT)	ARC
20	Antibody which Neutralizes pH Labile alpha aberrant Interferon	Advanced Biotherapy Concepts (Rockville, MD)	AIDS, ARC
	AR177	Aronex Pharm	HIV infection, AIDS, ARC
25	Beta-fluoro-ddA	Nat'l Cancer Institute	AIDS-associated diseases
30	BMS-232623 (CGP-73547)	Bristol-Myers Squibb/ Novartis	HIV infection, AIDS, ARC (protease inhibitor)
35	BMS-234475 (CGP-61755)	Bristol-Myers Squibb/ Novartis	HIV infection, AIDS, ARC (protease inhibitor)
	CI-1012	Warner-Lambert	HIV-1 infection
40	Cidofovir	Gilead Science	CMV retinitis, herpes, papillomavirus
	Curdan sulfate	AJI Pharma USA	HIV infection
	Cytomegalovirus Immune globin	MedImmune	CMV retinitis

	Cytovene	Syntex	Sight threatening
5	Ganciclovir		CMV peripheral CMV retinitis
10	Delaviridine	Pharmacia-Upjohn	HIV infection, AIDS, ARC (RT inhibitor)
15	Dextran Sulfate	Ueno Fine Chem. Ind. Ltd. (Osaka, Japan)	AIDS, ARC, HIV positive asymptomatic
	ddC Dideoxycytidine	Hoffman-La Roche	HIV infection, AIDS, ARC
20	ddI Dideoxyinosine	Bristol-Myers Squibb	HIV infection, AIDS, ARC; combination with AZT/d4T
25	DMP-450	AVID (Camden, NJ)	HIV infection, AIDS, ARC (protease inhibitor)
30	Efavirenz (DMP 266) (-)-6-Chloro-4-(S)- cyclopropylethynyl- 4(S)-trifluoro- methyl-1,4-dihydro- 2H-3,1-benzoxazin- 2-one, STOCRINE	DuPont Merck	HIV infection, AIDS, ARC (non-nucleoside RT inhibitor)
35	EL10	Elan Corp, PLC (Gainesville, GA)	HIV infection
40	Famciclovir	Smith Kline	herpes zoster, herpes simplex

5	FTC	Emory University	HIV infection, AIDS, ARC (reverse transcriptase inhibitor)
10	GS 840	Gilead	HIV infection, AIDS, ARC (reverse transcriptase inhibitor)
15	HBV097	Hoechst Marion Roussel	HIV infection, AIDS, ARC (non-nucleoside reverse transcriptase inhibitor)
20	Hypericin	VIMRx Pharm.	HIV infection, AIDS, ARC
	Recombinant Human Interferon Beta	Triton Biosciences (Alameda, CA)	AIDS, Kaposi's sarcoma, ARC
25	Interferon alfa-n3	Interferon Sciences	ARC, AIDS
30	Indinavir	Merck	HIV infection, AIDS, ARC, asymptomatic HIV positive, also in combination with AZT/ddI/ddC
	ISIS 2922	ISIS Pharmaceuticals	CMV retinitis
35	KNI-272	Nat'l Cancer Institute	HIV-assoc. diseases
40	Lamivudine, 3TC	Glaxo Wellcome	HIV infection, AIDS, ARC (reverse transcriptase inhibitor); also with AZT
	Lobucavir	Bristol-Myers Squibb	CMV infection

5	Nelfinavir	Agouron Pharmaceuticals	HIV infection, AIDS, ARC (protease inhibitor)
	Nevirapine	Boeheringer Ingelheim	HIV infection, AIDS, ARC (RT inhibitor)
10	Novapren	Novaferon Labs, Inc. (Akron, OH)	HIV inhibitor
15	Peptide T Octapeptide Sequence	Peninsula Labs (Belmont, CA)	AIDS
20	Trisodium Phosphonoformate	Astra Pharm. Products, Inc.	CMV retinitis, HIV infection, other CMV infections
	PNU-140690	Pharmacia Upjohn	HIV infection, AIDS, ARC (protease inhibitor)
25	Probucol	Vyrex	HIV infection, AIDS
	RBC-CD4	Sheffield Med. Tech (Houston, TX)	HIV infection, AIDS, ARC
30	Ritonavir	Abbott	HIV infection, AIDS, ARC (protease inhibitor)
35	Saquinavir	Hoffmann- LaRoche	HIV infection, AIDS, ARC (protease inhibitor)
40	Stavudine; d4T Didehydrodeoxy- thymidine	Bristol-Myers Squibb	HIV infection, AIDS, ARC
	Valaciclovir	Glaxo Wellcome	Genital HSV & CMV infections

	Virazole Ribavirin	Viratek/ICN (Costa Mesa, CA)	asymptomatic HIV positive, LAS, ARC
5	VX-478	Vertex	HIV infection, AIDS, ARC
	Zalcitabine	Hoffmann-LaRoche	HIV infection, AIDS, ARC, with AZT
10	Zidovudine; AZT	Glaxo Wellcome	HIV infection, AIDS, ARC, Kaposi's sarcoma, in combination with other therapies
15			

IMMUNOMODULATORS

	<u>Drug Name</u>	<u>Manufacturer</u>	<u>Indication</u>
20	AS-101	Wyeth-Ayerst	AIDS
	Bropirimine	Pharmacia Upjohn	Advanced AIDS
25	Acemannan	Carrington Labs, Inc. (Irving, TX)	AIDS, ARC
	CL246,738	American Cyanamid Lederle Labs	AIDS, Kaposi's sarcoma
30	EL10	Elan Corp, PLC (Gainesville, GA)	HIV infection
	FP-21399	Fuki ImmunoPharm	Blocks HIV fusion with CD4+ cells

	Gamma Interferon	Genentech	ARC, in combination w/TNF (tumor necrosis factor)
5	Granulocyte Macrophage Colony Stimulating Factor	Genetics Institute Sandoz	AIDS
10	Granulocyte Macrophage Colony Stimulating Factor	Hoechst-Roussel Immunex	AIDS
15	Granulocyte Macrophage Colony Stimulating Factor	Schering-Plough	AIDS, combination w/AZT
	HIV Core Particle Immunostimulant	Rorer	Seropositive HIV
20	IL-2 Interleukin-2	Cetus	AIDS, in combination w/AZT
25	IL-2 Interleukin-2	Hoffman-LaRoche Immunex	AIDS, ARC, HIV, in combination w/AZT
	IL-2 Interleukin-2 (aldeslukin)	Chiron	AIDS, increase in CD4 cell counts
30	Immune Globulin Intravenous (human)	Cutter Biological (Berkeley, CA)	Pediatric AIDS, in combination w/AZT
35	IMREG-1	Imreg (New Orleans, LA)	AIDS, Kaposi's sarcoma, ARC, PGL
40	IMREG-2	Imreg (New Orleans, LA)	AIDS, Kaposi's sarcoma, ARC, PGL
	Imuthiol Diethyl Dithio Carbamate	Merieux Institute	AIDS, ARC
45	Alpha-2 Interferon	Schering Plough	Kaposi's sarcoma w/AZT, AIDS

	Methionine- Enkephalin	TNI Pharmaceutical (Chicago, IL)	AIDS, ARC
5	MTP-PE Muramyl-Tripeptide	Ciba-Geigy Corp.	Kaposi's sarcoma
10	Granulocyte Colony Stimulating Factor	Amgen	AIDS, in combination w/AZT
	Remune	Immune Response Corp.	Immunotherapeutic
15	rCD4 Recombinant Soluble Human CD4	Genentech	AIDS, ARC
20	rCD4-IgG hybrids		AIDS, ARC
	Recombinant Soluble Human CD4	Biogen	AIDS, ARC
25	Interferon Alfa 2a	Hoffman-La Roche	Kaposi's sarcoma AIDS, ARC, in combination w/AZT
30	SK&F106528 Soluble T4	Smith Kline	HIV infection
35	Thymopentin	Immunobiology Research Institute (Annandale, NJ)	HIV infection
	Tumor Necrosis Factor; TNF	Genentech	ARC, in combination w/gamma Interferon

ANTI-INFECTIVES

	<u>Drug Name</u>	<u>Manufacturer</u>	<u>Indication</u>
5	Clindamycin with Primaquine	Pharmacia Upjohn	PCP
10	Fluconazole	Pfizer	Cryptococcal meningitis, candidiasis
	Pastille Nystatin Pastille	Squibb Corp.	Prevention of oral candidiasis
15	Ornidyl Eflornithine	Merrell Dow	PCP
20	Pentamidine Isethionate (IM & IV)	LyphoMed (Rosemont, IL)	PCP treatment
	Trimethoprim		Antibacterial
	Trimethoprim/sulfa		Antibacterial
25	Piritrexim	Burroughs Wellcome	PCP treatment
30	Pentamidine Isethionate for Inhalation	Fisons Corporation	PCP prophylaxis
	Spiramycin	Rhone-Poulenc diarrhea	Cryptosporidial
35	Intraconazole-R51211	Janssen-Pharm.	Histoplasmosis; cryptococcal meningitis
	Trimetrexate	Warner-Lambert	PCP



	Daunorubicin	NeXstar, Sequus	Kaposi's sarcoma
5	Recombinant Human Erythropoietin	Ortho Pharm. Corp.	Severe anemia assoc. with AZT therapy
10	Recombinant Human Growth Hormone	Serono	AIDS-related wasting, cachexia
	Megestrol Acetate	Bristol-Myers Squibb	Treatment of anorexia assoc. W/AIDS
15	Testosterone	Alza, Smith Kline	AIDS-related wasting
20	Total Enteral Nutrition	Norwich Eaton Pharmaceuticals	Diarrhea and malabsorption related to AIDS

Additionally, the compounds of the invention herein may be used in combination with another class of agents for treating AIDS which are called HIV entry inhibitors. Examples of such HIV entry inhibitors are discussed in DRUGS OF THE FUTURE 1999, 24(12), pp. 1355-1362; CELL, Vol. 9, pp. 243-246, Oct. 29, 1999; and DRUG DISCOVERY TODAY, Vol. 5, No. 5, May 2000, pp. 183-194.

It will be understood that the scope of combinations of the compounds of this invention with AIDS antivirals, immunomodulators, anti-infectives, HIV entry inhibitors or vaccines is not limited to the list in the above Table, but includes in principle any combination with any pharmaceutical composition useful for the treatment of AIDS.

Preferred combinations are simultaneous or alternating treatments of with a compound of the present invention and an inhibitor of HIV protease and/or a non-nucleoside inhibitor of HIV reverse transcriptase. An optional fourth component in the combination is a nucleoside inhibitor of HIV reverse transcriptase, such as AZT, 3TC, ddC or ddI. A preferred inhibitor of HIV protease is indinavir, which is the sulfate salt of N-(2(R)-hydroxy-1-(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-

(1-(4-(3-pyridyl-methyl)-2(S)-N<sup>7</sup>-(t-butylcarboxamido)-piperazinyl))-pentaneamide ethanolate, and is synthesized according to U.S. 5,413,999. Indinavir is generally administered at a dosage of 800 mg three times a day. Other preferred protease inhibitors are nelfinavir and ritonavir. Another preferred inhibitor of HIV protease is saquinavir which is administered in a dosage of 600 or 1200 mg tid. Preferred non-nucleoside inhibitors of HIV reverse transcriptase include efavirenz. The preparation of ddC, ddI and AZT are also described in EPO 0,484,071. These combinations may have unexpected effects on limiting the spread and degree of infection of HIV. Preferred combinations include those with the following (1) indinavir with efavirenz, and, optionally, AZT and/or 3TC and/or ddI and/or ddC; (2) indinavir, and any of AZT and/or ddI and/or ddC and/or 3TC, in particular, indinavir and AZT and 3TC; (3) stavudine and 3TC and/or zidovudine; (4) zidovudine and lamivudine and 141W94 and 1592U89; (5) zidovudine and lamivudine.

In such combinations the compound of the present invention and other active agents may be administered separately or in conjunction. In addition, the administration of one element may be prior to, concurrent to, or subsequent to the administration of other agent(s).

The preparative procedures and anti-HIV-1 activity of the novel azaindole piperazine diamide analogs of Formula I are summarized below in Schemes 1-64.

### Abbreviations

The following abbreviations, most of which are conventional abbreviations well known to those skilled in the art, are used throughout the description of the invention and the examples. Some of the abbreviations used are as follows:

h	=	hour(s)
rt	=	room temperature
mol	=	mole(s)
mmol	=	millimole(s)
g	=	gram(s)
mg	=	milligram(s)
mL	=	milliliter(s)
TFA	=	Trifluoroacetic Acid

	DCE	=	1,2-Dichloroethane
	CH <sub>2</sub> Cl <sub>2</sub>	=	Dichloromethane
	TPAP	=	tetrapropylammonium perruthenate
	THF	=	Tetrahydrofuran
5	DEPBT	=	3-(Diethoxyphosphoryloxy)-1,2,3-benzotriazin-4(3H)-one
	DMAP	=	4-dimethylaminopyridine
	P-EDC	=	Polymer supported 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide
10	EDC	=	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide
	DMF	=	<i>N,N</i> -dimethylformamide
	Hunig's Base	=	<i>N,N</i> -Diisopropylethylamine
	mCPBA	=	<i>meta</i> -Chloroperbenzoic Acid
	azaindole	=	1 <i>H</i> -Pyrrolo-pyridine
15	4-azaindole	=	1 <i>H</i> -pyrrolo[3,2- <i>b</i> ]pyridine
	5-azaindole	=	1 <i>H</i> -Pyrrolo[3,2- <i>c</i> ]pyridine
	6-azaindole	=	1 <i>H</i> -pyrrolo[2,3- <i>c</i> ]pyridine
	7-azaindole	=	1 <i>H</i> -Pyrrolo[2,3- <i>b</i> ]pyridine
	PMB	=	4-Methoxybenzyl
20	DDQ	=	2, 3-Dichloro-5, 6-dicyano-1, 4-benzoquinone
	OTf	=	Trifluoromethanesulfonyl
	NMM	=	4-Methylmorpholine
	PIP-COPh	=	1-Benzoylpiperazine
	NaHMDS	=	Sodium hexamethyldisilazide
25	EDAC	=	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide
	TMS	=	Trimethylsilyl
	DCM	=	Dichloromethane
30	DCE	=	Dichloroethane


	MeOH	=	Methanol
	THF	=	Tetrahydrofuran
5	EtOAc	=	Ethyl Acetate
	LDA	=	Lithium diisopropylamide
	TMP-Li	=	2,2,6,6-tetramethylpiperidinyllithium
10	DME	=	Dimethoxyethane
	DIBALH	=	Diisobutylaluminum hydride
15	HOBT	=	1-hydroxybenzotriazole
	CBZ	=	Benzoyloxycarbonyl
	PCC	=	Pyridinium chlorochromate
20			

### Chemistry

The present invention comprises compounds of Formula I, their pharmaceutical formulations, and their use in patients suffering from or susceptible to HIV infection. The compounds of Formula I include pharmaceutically acceptable salts thereof.

General procedures to construct substituted azaindole piperazine diamides of Formula I and intermediates useful for their synthesis are described in the following Schemes.

Scheme 1



- 5 Step A in Scheme 1 depicts the synthesis of an aza indole intermediate, 2, via the well known Bartoli reaction in which vinyl magnesium bromide reacts with an aryl or heteroaryl nitro group, such as in 1, to form a five-membered nitrogen containing ring as shown. Some references for the above transformation include: Bartoli et al. a) *Tetrahedron Lett.* **1989**, 30, 2129. b) *J. Chem. Soc. Perkin Trans. I*
- 10 **1991**, 2757. c) *J. Chem. Soc. Perkin Trans. II* **1991**, 657. d) *Synthesis* (1999), 1594. In the preferred procedure, a solution of vinyl Magnesium bromide in THF (typically 1.0M but from 0.25 to 3.0M) is added dropwise to a solution of the nitro pyridine in THF at  $-78^\circ$  under an inert atmosphere of either nitrogen or Argon. After addition is completed, the reaction temperature is allowed to warm to  $-20^\circ$  and then is stirred for
- 15 approximately 12h before quenching with 20% aq ammonium chloride solution. The reaction is extracted with ethyl acetate and then worked up in a typical manner using a drying agent such as anhydrous magnesium sulfate or sodium sulfate. Products are generally purified using chromatography over Silica gel. Best results are generally

achieved using freshly prepared vinyl Magnesium bromide. In some cases, vinyl Magnesium chloride may be substituted for vinyl Magnesium bromide.

Substituted azaindoles may be prepared by methods described in the literature or may be available from commercial sources. Thus there are many methods for carrying out step A in the literature and the specific examples are too numerous to even list. Alternative syntheses of aza indoles and general methods for carrying out step A include, but are not limited to, those described in the following references (a-k below): a) Prokopov, A. A.; Yakhontov, L. N. *Khim.-Farm. Zh.* **1994**, *28*(7), 30-51; b) Lablache-Combier, A. **Heteroaromatics**. Photoinduced Electron Transfer **1988**, Pt. C, 134-312; c) Saify, Zafar Said. *Pak. J. Pharmacol.* **1986**, *2*(2), 43-6; d) Bisagni, E. *Jerusalem Symp. Quantum Chem. Biochem.* **1972**, *4*, 439-45; e) Yakhontov, L. N. *Usp. Khim.* **1968**, *37*(7), 1258-87; f) Willette, R. E. *Advan. Heterocycl. Chem.* **1968**, *9*, 27-105; g) Mahadevan, I.; Rasmussen, M. *Tetrahedron* **1993**, *49*(33), 7337-52; h) Mahadevan, I.; Rasmussen, M. *J. Heterocycl. Chem.* **1992**, *29*(2), 359-67; i) Spivey, A. C.; Fekner, T.; Spey, S. E.; Adams, H. *J. Org. Chem.* **1999**, *64*(26), 9430-9443; j) Spivey, A.C.; Fekner, T.; Adams, H. *Tetrahedron Lett.* **1998**, *39*(48), 8919-8922; k) *Advances in Heterocyclic Chemistry* (Academic press) **1991**, Vol. 52, pg 235-236 and references therein.

20

**Step B.** Intermediate 3 can be prepared by reaction of aza-indole, intermediate 2, with an excess of ClCOCOOME in the presence of AlCl<sub>3</sub> (aluminum chloride) (Sycheva et al, Ref. 26, Sycheva, T.V.; Rubtsov, N.M.; Sheinker, Yu.N.; Yakhontov, L.N. Some reactions of 5-cyano-6-chloro-7-azaindoles and lactam-lactim tautomerism in 5-cyano-6-hydroxy-7-azaindoles. *Khim. Geterotsikl. Soedin.*, **1987**, 100-106). Typically an inert solvent such as CH<sub>2</sub>Cl<sub>2</sub> is used but others such as THF, Et<sub>2</sub>O, DCE, dioxane, benzene, or toluene may find applicability either alone or in mixtures. Other oxalate esters such as ethyl or benzyl mono esters of oxalic acid could also suffice for either method shown above. More lipophilic esters ease isolation during aqueous extractions. Phenolic or substituted phenolic (such as pentafluorophenol) esters enable direct coupling of the HW(C=O)A group, such as a piperazine, in Step D without activation. Lewis acid catalysts, such as tin tetrachloride, titanium IV chloride, and aluminum chloride are employed in Step B with aluminum chloride being most preferred. Alternatively, the azaindole is treated with a Grignard reagent such as MeMgI (methyl magnesium iodide), methyl magnesium bromide or ethyl magnesium bromide and a zinc halide, such as ZnCl<sub>2</sub> (zinc chloride) or zinc bromide, followed by the addition of an oxalyl chloride mono ester, such as ClCOCOOME (methyl chlorooxoacetate) or another ester as above, to

35

- afford the aza-indole glyoxyl ester (Shadrina et al, Ref. 25). Oxalic acid esters such as methyl oxalate, ethyl oxalate or as above are used. Aprotic solvents such as  $\text{CH}_2\text{Cl}_2$ ,  $\text{Et}_2\text{O}$ , benzene, toluene, DCE, or the like may be used alone or in combination for this sequence. In addition to the oxalyl chloride mono esters, oxalyl chloride itself may
- 5 be reacted with the azaindole and then further reacted with an appropriate amine, such as a piperazine derivative (See Scheme 52, for example).

- Step C.** Hydrolysis of the methyl ester, (intermediate 3, Scheme 1) affords a potassium salt of intermediate 4, which is coupled with mono-benzoylated
- 10 piperazine derivatives as shown in Step D of Scheme 1. Some typical conditions employ methanolic or ethanolic sodium hydroxide followed by careful acidification with aqueous hydrochloric acid of varying molarity but 1M HCl is preferred. The acidification is not utilized in many cases as described above for the preferred conditions. Lithium hydroxide or potassium hydroxide could also be employed and
- 15 varying amounts of water could be added to the alcohols. Propanols or butanols could also be used as solvents. Elevated temperatures up to the boiling points of the solvents may be utilized if ambient temperatures do not suffice. Alternatively, the hydrolysis may be carried out in a non polar solvent such as  $\text{CH}_2\text{Cl}_2$  or THF in the presence of Triton B. Temperatures of  $-78^\circ\text{C}$  to the boiling point of the solvent may
- 20 be employed but  $-10^\circ\text{C}$  is preferred. Other conditions for ester hydrolysis are listed in reference 41 and both this reference and many of the conditions for ester hydrolysis are well known to chemists of average skill in the art.


**Alternative procedures for step B and C:**

25 **Imidazolium Chloroaluminate:**

- We found that ionic liquid 1-alkyl-3-alkylimidazolium chloroaluminate is generally useful in promoting the Friedel-Crafts type acylation of indoles and azaindoles. The ionic liquid is generated by mixing 1-alkyl-3-alkylimidazolium
- 30 chloride with aluminium chloride at room temperature with vigorous stirring. 1:2 or 1:3 molar ratio of 1-alkyl-3-alkylimidazolium chloride to aluminium chloride is preferred. One particular useful imidazolium chloroaluminate for the acylation of azaindole with methyl or ethyl chlorooxoacetate is the 1-ethyl-3-methylimidazolium chloroaluminate. The reaction is typically performed at ambient temperature and the
- 35 azaindoleglyoxyl ester can be isolated. More conveniently, we found that the glyoxyl ester can be hydrolyzed *in situ* at ambient temperature on prolonged reaction time

(typically overnight) to give the corresponding glyoxyl acid for amide formation (Scheme 1).

Scheme 1



A representative experimental procedure is as follows: 1-ethyl-3-methylimidazolium chloride (2 equiv.; purchased from TCI; weighted under a stream of nitrogen) was stirred in an oven-dried round bottom flask at r.t. under a nitrogen atmosphere, and added aluminium chloride (6 equiv.; anhydrous powder packaged under argon in ampules purchased from Aldrich preferred; weighted under a stream of nitrogen). The mixture was vigorously stirred to form a liquid, which was then added azaindole (1 equiv.) and stirred until a homogenous mixture resulted. The reaction mixture was added dropwise ethyl or methyl chlorooxoacetate (2 equiv.) and then stirred at r.t. for 16 h. After which time, the mixture was cooled in an ice-water bath and the reaction quenched by carefully adding excess water. The precipitates were filtered, washed with water and dried under high vacuum to give the azaindoleglyoxyl acid. For some examples, 3 equivalents of 1-ethyl-3-methylimidazolium chloride and chlorooxoacetate may be required.


20

Related references: (1) Welton, T. *Chem Rev.* **1999**, *99*, 2071; (2) Surette, J. K. D.; Green, L.; Singer, R. D. *Chem. Commun.* **1996**, 2753; (3) Saleh, R. Y. WO 0015594.

25 **Step D.** The acid intermediate, 4, from step C of Scheme 1 is coupled with an amine A(C=O)WH preferably in the presence of DEPBT (3-(diethoxyphosphoryloxy)-1,2,3-benzotriazin-4(3H)-one) and *N,N*-diisopropylethylamine, commonly known as Hunig's base, to provide azaindole piperazine diamides. DEPBT was prepared according to the procedure of Ref. 28, Li, H.; Jiang, X.; Ye, Y.-H.; Fan, C.; Romoff, T.; Goodman, M. *Organic Lett.*, **1999**, *1*, 91-93. Typically an inert solvent such as DMF or THF is used but other aprotic solvents could be used. The group W as referred to herein is

30





The amide bond construction reaction could be carried out using the preferred conditions described above, the EDC conditions described below, other coupling  
 5 conditions described in this application, or alternatively by applying the conditions or coupling reagents for amide bond construction described later in this application for construction of substituents R<sub>1</sub>-R<sub>4</sub>. Some specific nonlimiting examples are given in this application.

10 The mono-substituted piperazine derivatives can be prepared according to well established procedures such as those described by Desai et al, Ref. 27(a), Adamczyk et al, Ref. 27(b), Rossen et al, Ref. 27(c), and Wang et al, 27(d).


Additional procedures for synthesizing, modifying and attaching groups :  
 15 (C=O)<sub>m</sub>-WC(O)-A are contained in PCT WO 00/71535.

20

25

30

Scheme 2



- 5 Scheme 2 provides a more specific example of the transformations previously described in Scheme 1. Intermediates 6-10 are prepared by the methodologies as described for intermediates 1a-5a in Scheme 1. Scheme 2A is another embodiment of the transformations described in Schemes 1 and 2. Conversion of the phenol to the chloride (Step S, Scheme 2A) may be accomplished according to the procedures described in Reimann, E.; Wichmann, P.; Hoefner, G.; *Sci. Pharm.* **1996**, *64*(3), 637-646; and Katritzky, A.R.; Rachwal, S.; Smith, T.P.; Steel, P.J.; *J. Heterocycl. Chem.* **1995**, *32*(3), 979-984. Step T of Scheme 2A can be carried out as described for Step A of Scheme 1. The bromo intermediate can then be converted into alkoxy, chloro, or fluoro intermediates as shown in Step U of Scheme 2A. Scheme 2A describes the preferred method for preparing intermediate 6c or other closely related compounds containing a 4 methoxy group in the 6-azaindole system. When step U is the conversion of the bromide into alkoxy derivatives, the conversion may be carried out by reacting the bromide with an excess of sodium methoxide in methanol with cuprous salts, such as copper I bromide, copper I iodide, and copper I cyanide. The
- 10  
 15


temperature may be carried out at temperatures of between ambient and 175° but most likely will be around 115°C or 100°C. The reaction may be run in a pressure vessel or sealed tube to prevent escape of volatiles such as methanol. The preferred conditions utilize 3eq of sodium methoxide in methanol, CuBr as the reaction catalyst (0.2 to 3 equivalents with the preferred being 1 eq or less) , and a reaction temperature of 115° C. The reaction is carried out in a sealed tube or sealed reaction vessel. The conversion of the bromide into alkoxy derivatives may also be carried out according to procedures described in Palucki, M.; Wolfe, J.P.; Buchwald, S.L.; *J. Am. Chem. Soc.* **1997**, *119*(14), 3395-3396; Yamato, T.; Komine, M.; Nagano, Y.; *Org. Prep. Proc. Int.* **1997**, *29*(3), 300-303; Rychnovsky, S.D.; Hwang, K.; *J. Org. Chem.* **1994**, *59*(18), 5414-5418. Conversion of the bromide to the fluoro derivative (Step U, Scheme 2A) may be accomplished according to Antipin, I.S.; Vigalok, A.I.; Konovalov, A.I.; *Zh. Org. Khim.* **1991**, *27*(7), 1577-1577; and Uchibori, Y.; Umeno, M.; Seto, H.; Qian, Z.; Yoshioka, H.; *Synlett.* **1992**, *4*, 345-346. Conversion of the bromide to the chloro derivative (Step U, Scheme 2A) may be accomplished according to procedures described in Gilbert, E.J.; Van Vranken, D.L.; *J. Am. Chem. Soc.* **1996**, *118*(23), 5500-5501; Mongin, F.; Mongin, O.; Trecourt, F.; Godard, A.; Queguiner, G.; *Tetrahedron Lett.* **1996**, *37*(37), 6695-6698; and O'Connor, K.J.; Burrows, C.J.; *J. Org. Chem.* **1991**, *56*(3), 1344-1346. Steps V, W and X of Scheme 2A are carried out according to the procedures previously described for Steps B, C, and D of Scheme 1, respectively. The steps of Scheme 2A may be carried out in a different order as shown in Scheme 2B and Scheme 2C.

25


30

35


## Scheme 2A



## Scheme 2B



## Scheme 2C




5

10

15


Scheme 3



5 Scheme 3 shows the synthesis of 4-azaindole derivatives 1b-5b, 5-azaindole derivatives 1c-5c, and 7-azaindole derivatives 1d-5d. The methods used to synthesize 1b-5b, 1c-5c, and 1d-5d are the same methods described for the synthesis of 1a-5a as described in Scheme 1. It is understood, for the purposes of Scheme 3, that 1b is used to synthesize 2b-5b, 1c provides 2c-5c and 1d provides 2d-5d.

The compounds where there is a single carbonyl between the azaindole and group W can be prepared by the method of Kelarev, V. I.; Gasanov, S. Sh.; Karakhanov, R. A.; Polivin, Yu. N.; Kuatbekova, K. P.; Panina, M. E.; *Zh. Org. Khim* 1992, 28(12), 2561-2568. In this method azaindoles are reacted with trichloroacetyl chloride in pyridine and then subsequently with KOH in methanol to provide the 3-carbomethoxy azaindoles shown in Scheme 4 which can then be hydrolyzed to the acid and carried through the coupling sequence with  $\text{HW}(\text{C}=\text{O})\text{A}$  to provide the compounds of Formula I wherein a single carbonyl links the azaindole moiety and group W.

Scheme 4




15

An alternative method for carrying out the sequence outlined in steps B-D (shown in Scheme 5) involves treating an azaindole, such as 11, obtained by procedures described in the literature or from commercial sources, with  $\text{MeMgI}$  and  $\text{ZnCl}_2$ , followed by the addition of  $\text{ClCOCOC}(\text{Cl})$  (oxalyl chloride) in either THF or  $\text{Et}_2\text{O}$  to afford a mixture of a glyoxyl chloride azaindole, 12a, and an acyl chloride azaindole, 12b. The resulting mixture of glyoxyl chloride azaindole and acyl chloride azaindole is then coupled with mono-benzoylated piperazine derivatives under basic conditions to afford the products of step D as a mixture of compounds, 13a and 13b, where either one or two carbonyl groups link the azaindole and group W. Separation via chromatographic methods which are well known in the art provides the pure 13a and 13b. This sequence is summarized in Scheme 5, below.

25




Scheme 5



5

Scheme 6



- Scheme 6 depicts a general method for modifying the substituent A.
- 10 Coupling of H-W-C(O)OtBu using the conditions described previously for W in Scheme 1, Step D provides Boc protected intermediate, 15. Intermediate 15 is then deprotected by treatment with an acid such as TFA, hydrochloric acid or formic acid using standard solvents or additives such as CH<sub>2</sub>Cl<sub>2</sub>, dioxane, or anisole and temperatures between -78 °C and 100 °C. Other acids such as aq hydrochloric or

perchloric may also be used for deprotection. Alternatively other nitrogen protecting groups on W such as Cbz or TROC, may be utilized and could be removed via hydrogenation or treatment with zinc respectively. A stable silyl protecting group such as phenyl dimethylsilyl could also be employed as a nitrogen protecting group  
5 on W and can be removed with fluoride sources such as tetrabutylammonium fluoride. Finally, the free amine is coupled to acid A-C(O)OH using standard amine-acid coupling conditions such as those used to attach group W or as shown below for amide formation on positions R<sub>1</sub>-R<sub>4</sub> to provide compound 16.

10 Some specific examples of general methods for preparing functionalized azaindoles or for interconverting functionality on aza indoles which will be useful for preparing the compounds of this invention are shown in the following sections for illustrative purposes. It should be understood that this invention covers substituted 4,  
5, 6, and 7 azaindoles and that the methodology shown below may be applicable to  
15 all of the above series while other shown below will be specific to one or more. A typical practitioner of the art can make this distinction when not specifically delineated. Many methods are intended to be applicable to all the series, particularly functional group installations or interconversions. For example, a general strategy for providing further functionality of this invention is to position or install a halide such  
20 as bromo, chloro, or iodo, aldehyde, cyano, or a carboxy group on the azaindole and then to convert that functionality to the desired compounds. In particular, conversion to substituted heteroaryl, aryl, and amide groups on the ring are of particular interest.

General routes for functionalizing azaindole rings are shown in Schemes 7, 8  
25 and 9. As depicted in Scheme 7, the azaindole, 17, can be oxidized to the corresponding *N*-oxide derivative, 18, by using mCPBA (meta-Chloroperbenzoic Acid) in acetone or DMF (eq. 1, Harada et al, Ref. 29 and Antonini et al, Ref. 34). The *N*-oxide, 18, can be converted to a variety of substituted azaindole derivatives by using well documented reagents such as phosphorus oxychloride (POCl<sub>3</sub>) (eq. 2,  
30 Schneller et al, Ref. 30), tetramethylammonium fluoride (Me<sub>4</sub>NF) (eq. 3), Grignard reagents RMgX (R = alkyl or aryl, X = Cl, Br or I) (eq. 4, Shiotani et al, Ref. 31), trimethylsilyl cyanide (TMSCN) (eq. 5, Minakata et al, Ref. 32) or Ac<sub>2</sub>O (eq. 6, Klemm et al, Ref. 33). Under such conditions, a chlorine (in 19), fluorine (in 20), nitrile (in 22), alkyl (in 21), aromatic (in 21) or hydroxyl group (in 24) can be  
35 introduced to the pyridine ring. Nitration of azaindole *N*-oxides results in introduction of a nitro group to azaindole ring, as shown in Scheme 8 (eq. 7, Antonini et al, Ref. 34). The nitro group can subsequently be displaced by a variety of nucleophilic agents, such as OR, NR<sup>1</sup>R<sup>2</sup> or SR, in a well established chemical fashion

(eq. 8, Regnouf De Vains et al, Ref. 35(a), Miura et al, Ref. 35(b), Profft et al, Ref. 35(c)). The resulting N-oxides, 26, are readily reduced to the corresponding azaindole, 27, using phosphorus trichloride (PCl<sub>3</sub>) (eq. 9, Antonini et al, Ref. 34 and Nesi et al, Ref. 36). Similarly, nitro-substituted *N*-oxide, 25, can be reduced to the azaindole, 28, using phosphorus trichloride (eq. 10). The nitro group of compound 28 can be reduced to either a hydroxylamine (NHOH), as in 29, (eq. 11, Walser et al, Ref. 37(a) and Barker et al, Ref. 37(b)) or an amino (NH<sub>2</sub>) group, as in 30, (eq. 12, Nesi et al, Ref. 36 and Ayyangar et al, Ref. 38) by carefully selecting different reducing conditions.

10

15







20

25







30

35

Scheme 7




## Scheme 8



The alkylation of the nitrogen atom at position 1 of the azaindole derivatives  
 5 can be achieved using  $\text{NaH}$  as the base,  $\text{DMF}$  as the solvent and an alkyl halide or sulfonate as alkylating agent, according to a procedure described in the literature (Mahadevan et al, Ref. 39) (Scheme 9).


## Scheme 9



In the general routes for substituting the azaindole ring described above, each process can be applied repeatedly and combinations of these processes is permissible in order to provide azaindoles incorporating multiple substituents. The application of such processes provides additional compounds of Formula I.

5

Scheme 10



10 The synthesis of 4-aminoazaindoles which are useful precursors for 4, 5, and/or 7-substituted azaindoles is shown in Scheme 10 above.

The synthesis of 3, 5-dinitro-4-methylpyridine, 32, is described in the following two references by Achremowicz et.al.: Achremowicz, Lucjan. *Pr. Nauk. Inst. Chem. Org. Fiz. Politech. Wroclaw*. 1982, 23, 3-128; Achremowicz, Lucjan. *Synthesis* 1975, 10, 653-4. In the first step of Scheme 10, the reaction with dimethylformamide dimethyl acetal in an inert solvent or neat under conditions for forming Batcho-Leimgruber precursors provides the cyclization precursor, 33, as shown. Although the step is anticipated to work as shown, the pyridine may be oxidized to the N-oxide prior to the reaction using a peracid such as MCPBA or a more potent oxidant like meta-trifluoromethyl or meta nitro peroxy benzoic acids. In the second step of Scheme 10, reduction of the nitro group using for example hydrogenation over Pd/C catalyst in a solvent such as MeOH, EtOH, or EtOAc provides the cyclized product, 34. Alternatively the reduction may be carried out using tin dichloride and HCl, hydrogenation over Raney nickel or other catalysts, or by using other methods for nitro reduction such as described elsewhere in this application.

20 The amino indole, 34, can now be converted to compounds of Formula I via, for example, diazotization of the amino group, and then conversion of the diazonium salt to the fluoride, chloride or alkoxy group. See the discussion of such conversions in the descriptions for Schemes 17 and 18. The conversion of the amino moiety into desired functionality could then be followed by installation of the oxoacetopiperazine moiety by the standard methodology described above. 5 or 7-substitution of the azaindole can arise from N-oxide formation at position 6 and subsequent conversion to the chloro via conditions such as POCl<sub>3</sub> in chloroform, acetic anhydride followed

by  $\text{POCl}_3$  in DMF, or alternatively  $\text{TsCl}$  in DMF. Literature references for these and other conditions are provided in some of the later Schemes in this application. The synthesis of 4-bromo-7-hydroxy or protected hydroxy-4-azaindole is described below as this is a useful precursor for 4 and/or 7 substituted 6-aza indoles.


5

The synthesis of 5-bromo-2-hydroxy-4-methyl-3-nitro pyridine, 35, may be carried out as described in the following reference: Betageri, R.; Beaulieu, P.L.; Llinas-Brunet, M; Ferland, J.M.; Cardozo, M.; Moss, N.; Patel, U.; Proudfoot, J.R. PCT Int. Appl. WO 9931066, 1999. Intermediate 36 is prepared from 35 according to the method as described for Step 1 of Scheme 10. PG is an optional hydroxy protecting group such as triallylsilyl or the like. Intermediate 37 is then prepared from 36 by the selective reduction of the nitro group in the presence of bromide and subsequent cyclization as described in the second step of Scheme 10.  $\text{Fe}(\text{OH})_2$  in DMF with catalytic tetrabutylammonium bromide can also be utilized for the reduction of the nitro group. The bromide may then be converted to fluoride via displacement with fluoride anions or to other substituents. The compounds are then converted to compounds of Formula I as above.

15

### Scheme 11

20




25

An alternate method for preparing substituted 6-azaindoles is shown below in Schemes 12 and 13. It should be recognized that slight modifications of the route depicted below are possible. For example, acylation reactions of the 3 position of what will become the azaindole five membered ring, prior to aromatization of the azaindole, may be carried out in order to obtain higher yields. In addition to a paramethoxybenzyl group (PMB), a benzyl group can be carried through the sequence and removed during azaindole formation by using  $\text{TsOH}$ , p-Chloranil, in benzene as the oxidant if DDQ is not optimal. The benzyl intermediate, 38, has been described by Ziegler et al. in *J. Am. Chem. Soc.* 1973, 95(22), 7458. The transformation of 38 to 40 is analogous to the transformation described in *Heterocycles* 1984, 22, 2313.


30

## Scheme 12



- 5 Scheme 13 describes various transformations of intermediate 40 which ultimately provide compounds of Formula I. The conversions of the phenol moiety to other functionality at position 4 ( $R_2$  position in Scheme 13) may be carried out by the following methods: 1) conversion of a phenol to methoxy group with silver oxide and MeI or diazomethane; 2) conversion of a phenolic hydroxy group to chloro using cat  $ZnCl_2$  and  $N,N$  dimethylaniline in  $CH_2Cl_2$  or  $PCl_5$  and  $POCl_3$  together; 3)
- 10 conversion of a phenolic hydroxy group to fluoro using diethylamine- $SF_5$  as in *Org. Prep. Proc. Int.* 1992, 24(1), 55-57. The method described in EP 427603, 1991, using the chloroformate and HF will also be useful. Other transformations are possible. For example the phenol can be converted to a triflate by standard methods
- 15 and used in coupling chemistries described later in this application.

## Scheme 13




- 1) Ketone alkylation to install  $R_3$
- 2) DDQ oxidation to form azaindole
- 3) Transformation of phenol ( $R_2 = OH$ ) into methyl ether, or Fluoro, chloro, etc
- 4) Use of C-7 directing group to functionalize at  $R_4$  or formation of N-oxide and  $POCl_3$  to make  $R_4 = chloro$
- 5) Conversion to compounds of Formula I as above

- 20 **Step E** Scheme 14 depicts the nitration of an azaindole, 41, ( $R_2 = H$ ). Numerous conditions for nitration of the azaindole may be effective and have been




- described in the literature.  $N_2O_5$  in nitromethane followed by aqueous sodium bisulfite according to the method of Bakke, J. M.; Raney, E.; *Synthesis* 1997, 3, 281-283 could be utilized. Nitric acid in acetic may also be employed as described in Kimura, H.; Yotsuya, S.; Yuki, S.; Sugi, H.; Shigehara, I.; Haga, T.; *Chem. Pharm. Bull.* 1995, 43(10), 1696-1700. Sulfuric acid followed by nitric acid may be employed as in Ruefenacht, K.; Kristinsson, H.; Mattern, G.; *Helv Chim Acta* 1976, 59, 1593. Coombes, R. G.; Russell, L. W.; *J. Chem. Soc., Perkin Trans. 1* 1974, 1751 describes the use of a Titanium based reagent system for nitration. Other conditions for the nitration of the azaindoles can be found in the following references:
- 10 Lever, O.W.J.; Werblood, H. M.; Russell, R. K.; *Synth. Comm.* 1993, 23(9), 1315-1320; Wozniak, M.; Van Der Plas, H. C.; *J. Heterocycl Chem.* 1978, 15, 731.

Scheme 14



Scheme 15



## Step F

- As shown above in Scheme 15, Step F, substituted azaindoles containing a chloride, bromide, iodide, triflate, or phosphonate undergo coupling reactions with a boronate (Suzuki type reactions) or a stannane to provide substituted azaindoles. Stannanes and boronates are prepared via standard literature procedures or as described in the experimental section of this application. The substituted indoles may undergo metal mediated coupling to provide compounds of Formula I wherein
- 25

R<sub>4</sub> is aryl, heteroaryl, or heteroalicyclic for example. The bromoazaindole intermediates, (or azaindole triflates or iodides) may undergo Stille-type coupling with heteroarylstannanes as shown in Scheme 15. Conditions for this reaction are well known in the art and the following are three example references a) Farina, V.; Roth, G.P. Recent advances in the Stille reaction; *Adv. Met.-Org. Chem.* **1996**, *5*, 1-53. b) Farina, V.; Krishnamurthy, V.; Scott, W.J. The Stille reaction ; *Org. React.* (N. Y.) **1997**, *50*, 1-652. and c) Stille, J. K. *Angew. Chem. Int. Ed. Engl.* **1986**, *25*, 508-524. Other references for general coupling conditions are also in the reference by Richard C. Larock *Comprehensive Organic Transformations* 2nd Ed. 1999, John Wiley and Sons New York. All of these references provide numerous conditions at the disposal of those skilled in the art in addition to the specific examples provided in Scheme 15 and in the specific embodiments. It can be well recognized that an indole stannane could also couple to a heterocyclic or aryl halide or triflate to construct compounds of Formula I. Suzuki coupling (Norio Miyaura and Akio Suzuki *Chem Rev.* **1995**, *95*, 2457.) between a triflate, bromo, or chloro azaindole intermediate and a suitable boronate could also be employed and some specific examples are contained in this application. Palladium catalyzed couplings of stannanes and boronates between chloro azaindole intermediates are also feasible and have been utilized extensively for this invention. Preferred procedures for coupling of a chloro azaindole and a stannane employ dioxane, stoichiometric or an excess of the tin reagent (up to 5 equivalents), 0.1 to 1 eq of Palladium (0) tetrakis triphenyl phosphine in dioxane heated for 5 to 15 h at 110 to 120°. Other solvents such as DMF, THF, toluene, or benzene could be employed. Preferred procedures for Suzuki coupling of a chloro azaindole and a boronate employ 1:1 DMF water as solvent, 2 equivalents of potassium carbonate as base stoichiometric or an excess of the boron reagent (up to 5 equivalents), 0.1 to 1 eq of Palladium (0) tetrakis triphenyl phosphine heated for 5 to 15 h at 110 to 120°. If standard conditions fail new specialized catalysts and conditions can be employed. Some references (and the references therein) describing catalysts which are useful for coupling with aryl and heteroaryl chlorides are:

Littke, A. F.; Dai, C.; Fu, G. C. *J. Am. Chem. Soc.* **2000**, *122*(17), 4020-4028; Varma, R. S.; Naicker, K. P. *Tetrahedron Lett.* **1999**, *40*(3), 439-442; Wallow, T. I.; Novak, B. M. *J. Org. Chem.* **1994**, *59*(17), 5034-7; Buchwald, S.; Old, D. W.; Wolfe, J. P.; Palucki, M.; Kamikawa, K.; Chieffi, A.; Sadighi, J. P.; Singer, R. A.; Ahman, J PCT Int. Appl. WO 0002887 2000; Wolfe, J. P.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **1999**, *38*(23), 3415; Wolfe, J. P.; Singer, R. A.; Yang, B. H.; Buchwald, S. L. *J. Am. Chem. Soc.* **1999**, *121*(41), 9550-9561; Wolfe, J. P.;


Buchwald, S. L. *Angew. Chem., Int. Ed.* 1999, 38(16), 2413-2416; Bracher, F.; Hildebrand, D.; *Liebigs Ann. Chem.* 1992, 12, 1315-1319; and Bracher, F.; Hildebrand, D.; *Liebigs Ann. Chem.* 1993, 8, 837-839.

- 5           Alternatively, the boronate or stannane may be formed on the azaindole via methods known in the art and the coupling performed in the reverse manner with aryl or heteroaryl based halogens or triflates.

10           Known boronate or stannane agents could be either purchased from commercial resources or prepared following disclosed documents. Additional examples for the preparation of tin reagents or boronate reagents are contained in the experimental section.

15           Novel stannane agents could be prepared from one of the following routes.

**Scheme Tin-01**




**Base = LDA, TMP-Li, n-BuLi, S-BuLi, t-BuLi**

**Solvent = THF, ether, DME**

**R = Me, Bu**

20

**Scheme Tin-02**




**Base = n-BuLi, S-BuLi, t-BuLi**

**Solvent = THF, ether, DME**

**R = Me, Bu**

## Scheme Tin-03




Solvent = THF, ether, DME

R = Me, Bu

5

## Scheme Tin-04




Solvent = Dioxane, Toluene

R = Me, Bu

## Scheme Tin-05

10

E = Electrophile = R'-halide, R'COCl, R'OCOCl, R'R''NCOCi, RSO<sub>2</sub>Cl, R'NCO, R'NSO, R'NCNR''Solvent = CH<sub>2</sub>Cl<sub>2</sub>, THF, Ether, DMF

R = Me, Bu

Base = NaH, BuLi, LDA, K<sub>2</sub>CO<sub>3</sub>, Et<sub>3</sub>N, DBU, DMAP, NaHMDS


Boronate reagents are prepared as described in reference 71. Reaction of lithium or Grignard reagents with trialkyl borates generates boronates. Alternatively, Palladium catalyzed couplings of alkoxy diboron or alkyl diboron reagents with aryl or heteroaryl halides can provide boron reagents for use in Suzuki type couplings. Some example conditions for coupling a halide with (MeO)BB(OMe)<sub>2</sub> utilize PdCl<sub>2</sub> (dppf), KOAc, DMSO, at 80°C until reaction is complete when followed by TLC or HPLC analysis.

20

Related examples are provided in the following experimental section.

Methods for direct addition of aryl or heteroaryl organometallic reagents to alpha chloro nitrogen containing heterocycles or the N-oxides of nitrogen containing heterocycles are known and applicable to the azaindoles. Some examples are  
 5 Shiotani et. Al. *J. Heterocyclic Chem.* 1997, 34(3), 901-907; Fourmigue et.al. *J. Org. Chem.* 1991, 56(16), 4858-4864.

Scheme 16



10

The preparation of a key aldehyde intermediate, 43, using a procedure adapted from the method of Gilmore et. Al. *Synlett* 1992, 79-80. Is shown in Scheme 16 above. The aldehyde substituent is shown only at the R<sub>4</sub> position for the sake of clarity, and should not be considered as a limitation of the methodology. The  
 15 bromide or iodide intermediate is converted into an aldehyde intermediate, 43, by metal-halogen exchange and subsequent reaction with dimethylformamide in an appropriate aprotic solvent. Typical bases used include, but are not limited to, alkyl lithium bases such as n-butyl lithium, sec butyl lithium or tert butyl lithium or a metal  
 20 such as lithium metal. A preferred aprotic solvent is THF. Typically the transmetalation is initiated at -78 °C. The reaction may be allowed to warm to allow the transmetalation to go to completion depending on the reactivity of the bromide intermediate. The reaction is then recooled to -78 °C and allowed to react with dimethylformamide. (allowing the reaction to warm may be required to enable  
 25 complete reaction) to provide an aldehyde which is elaborated to compounds of Formula I. Other methods for introduction of an aldehyde group to form intermediates of formula 43 include transition metal catalyzed carbonylation reactions of suitable bromo, trifluoromethane sulfonyl, or stannyl azaindoles.


Alternative the aldehydes can be introduced by reacting indolyl anions or indolyl Grignard reagents with formaldehyde and then oxidizing with MnO<sub>2</sub> or TPAP/NMO or other suitable oxidants to provide intermediate 43.

5           The methodology described in T. Fukuda et.al. *Tetrahedron* 1999, 55, 9151 and M. Iwao et. Al. *Heterocycles* 1992, 34(5), 1031 provide methods for preparing indoles with substituents at the 7-position. The Fukuda references provide methods for functionalizing the C-7 position of indoles by either protecting the indole nitrogen with 2,2-diethyl propanoyl group and then deprotonating the 7-position with sec/Buli  
10 in TMEDA to give an anion. This anion may be quenched with DMF, formaldehyde, or carbon dioxide to give the aldehyde, benzyl alcohol, or carboxylic acid respectively and the protecting group removed with aqueous t butoxide. Similar transformations can be achieved by converting indoles to indoline, lithiation at C-7 and then reoxidation to the indole such as described in the Iwao reference above. The  
15 oxidation level of any of these products may be adjusted by methods well known in the art as the interconversion of alcohol, aldehyde, and acid groups has been well studied. It is also well understood that a cyano group can be readily converted to an aldehyde. A reducing agent such as DIBALH in hexane such as used in Weyerstahl, P.; Schlicht, V.; *Liebigs Ann/Recl.* 1997, 1, 175-177 or alternatively catecholalane in  
20 THF such as used in Cha, J. S.; Chang, S. W.; Kwon, O. O.; Kim, J. M.; *Synlett.* 1996, 2, 165-166 will readily achieve this conversion to provide intermediates such as 44 (Scheme 16). Methods for synthesizing the nitriles are shown later in this application. It is also well understood that a protected alcohol, aldehyde, or acid group could be present in the starting azaindole and carried through the synthetic  
25 steps to a compound of Formula I in a protected form until they can be converted into the desired substituent at R<sub>1</sub> through R<sub>4</sub>. For example, a benzyl alcohol can be protected as a benzyl ether or silyl ether or other alcohol protecting group; an aldehyde may be carried as an acetal, and an acid may be protected as an ester or ortho ester until deprotection is desired and carried out by literature methods.

30

35

Scheme 17



- 5           **Step G**       Step 1 of Scheme 17 shows the reduction of a nitro group on 45 to the amino group of 46. Although shown on position 4 of the azaindole, the chemistry is applicable to other nitro isomers. The procedure described in Ciurla, H.; Puzsko, A.; *Khim Geterotsikl Soedin* 1996, 10, 1366-1371 uses hydrazine Raney-Nickel for the reduction of the nitro group to the amine. Robinson, R. P.; Donahue, O. K. M.; Son, P. S.; Wagdy, S. D.; *J. Heterocycl. Chem.* 1996, 33(2), 287-293 describes the use of hydrogenation and Raney Nickel for the reduction of the nitro group to the amine. Similar conditions are described by Nicolai, E.; Claude, S.; Teulon, J. M.; *J. Heterocycl. Chem.* 1994, 31(1), 73-75 for the same transformation. The following two references describe some trimethylsilyl sulfur or chloride based reagents which may be used for the reduction of a nitro group to an amine. Hwu, J.R.; Wong, F.F.; Shiao, M.J.; *J. Org. Chem.* 1992, 57(19), 5254-5255; Shiao, M.J.; Lai, L.L.; Ku, W.S.; Lin, P.Y.; Hwu, J.R.; *J. Org. Chem.* 1993, 58(17), 4742-4744.

20           Step 2 of Scheme 17 describes general methods for conversion of amino groups on azaindoles into other functionality. Scheme 18 also depicts transformations of an amino azaindole into various intermediates and compounds of Formula I.

25           The amino group at any position of the azaindole, such as 46 (Scheme 17), may be converted to a hydroxy group using sodium nitrite, sulfuric acid, and water via the method of Klemm, L. H.; Zell, R.; *J. Heterocycl. Chem.* 1968, 5, 773. Bradsher, C. K.; Brown, F. C.; Porter, H. K.; *J. Am. Chem. Soc.* 1954, 76, 2357

describes how the hydroxy group may be alkylated under standard or Mitsunobu conditions to form ethers. The amino group may be converted directly into a methoxy group by diazotization (sodium nitrite and acid) and trapping with methanol.


- 5           The amino group of an azaindole, such as 46, can be converted to fluoro via the method of Sanchez using  $\text{HPF}_6$ ,  $\text{NaNO}_2$ , and water by the method described in Sanchez, J. P.; Gogliotti, R. D.; *J. Heterocycl. Chem.* 1993, 30(4), 855-859. Other methods useful for the conversion of the amino group to fluoro are described in Rocca, P.; Marsais, F.; Godard, A.; Queguiner, G.; *Tetrahedron Lett.* 1993, 34(18),  
10   2937-2940 and Sanchez, J. P.; Rogowski, J.W.; *J. Heterocycl. Chem.* 1987, 24, 215.

- The amino group of the azaindole, 46, can also be converted to a chloride via diazotization and chloride displacement as described in Ciurla, H.; Puszko, A.; *Khim Geterotsikl Soedin* 1996, 10, 1366-1371 or the methods in Raveglia, L.F.; Giardina,  
15   G.A.; Grugni, M.; Rigolio, R.; Farina, C.; *J. Heterocycl. Chem.* 1997, 34(2), 557-559 or the methods in Matsumoto, J. I.; Miyamoto, T.; Minamida, A.; Mishimura, Y.; Egawa, H.; Mishimura, H.; *J. Med. Chem.* 1984, 27(3), 292; or as in Lee, T.C.; Salemnick, G.; *J. Org. Chem.* 1975, 24, 3608.

- 20           The amino group of the azaindole, 46, can also be converted to a bromide via diazotization and displacement by bromide as described in Raveglia, L.F.; Giardina, G.A.; Grugni, M.; Rigolio, R.; Farina, C.; *J. Heterocycl. Chem.* 1997, 34(2), 557-559; Talik, T.; Talik, Z.; Ban-Oganowska, H.; *Synthesis* 1974, 293; and Abramovitch, R.A.; Saha, M.; *Can. J. Chem.* 1966, 44, 1765.



Scheme 18



Steps 2 and 3 may be reversed as appropriate


5           The preparation of 4-amino 4-azaindole and 7-methyl-4-azaindole is described by Mahadevan, I.; Rasmussen, M. *J. Heterocycl. Chem.* **1992**, 29(2), 359-67. The amino group of the 4-amino 4-azaindole can be converted to halogens, hydroxy, protected hydroxy, triflate, as described above in Schemes 17-18 for the 4-amino compounds or by other methods known in the art. Protection of the indole nitrogen of the 7-methyl-4-azaindole via acetylation or other strategy followed by oxidation of the 7-methyl group with potassium permanganate or chromic acid provides the 7-acid /4-N-oxide. Reduction of the N-oxide, as described below, provides an intermediate from which to install various substituents at position R<sub>4</sub>. Alternatively the parent 4-azaindole which was prepared as described in Mahadevan, I.; Rasmussen, M. *J. Heterocycl. Chem.* **1992**, 29(2), 359-67 could be derivatized at nitrogen to provide the 1-(2,2-diethylbutanoyl)azaindole which could then be lithiated using TMEDA /sec BuLi as described in T. Fukuda et. Al. *Tetrahedron* **1999**, 55, 9151-9162; followed by conversion of the lithio species to the 7-carboxylic acid or 7-halogen as described. Hydrolysis of the N-amide using aqueous tert-butoxide in THF regenerates the free NH indole which can now be converted to compounds of Formula I. The chemistry used to functionalize position 7 can also be applied to the 5 and 6 indole series.

25           Scheme 19 shows the preparation of a 7-chloro-4-azaindole, 50, which can be converted to compounds of Formula I by the chemistry previously described, especially the palladium catalyzed tin and boron based coupling methodology described above. The chloro nitro indole, 49, is commercially available or can be

prepared from 48 according to the method of Delarge, J.; Lapiere, C. L. *Pharm. Acta Helv.* **1975**, 50(6), 188-91.

Scheme 19


5



Scheme 20, below, shows another synthetic route to substituted 4-aza indoles. The 3-aminopyrrole, 51, was reacted to provide the pyrrolopyridinone, 52, which was then reduced to give the hydroxy azaindole, 53. The pyrrolo[2,3-b]pyridines described were prepared according to the method of Britten, A.Z.; Griffiths, G.W.G. *Chem. Ind. (London)* **1973**, 6, 278. The hydroxy azaindole, 53, can then be converted to the triflate then further reacted to provide compounds of Formula I.

15

Scheme 20




The following references describe the synthesis of 7-halo or 7 carboxylic acid, or 7-amido derivatives of 5-azaindoline which can be used to construct compounds of Formula I. Bychikhina, N. N.; Azimov, V. A.; Yakhontov, L.N. *Khim. Geterotsikl. Soedin.* **1983**, 1, 58-62; Bychikhina, N. N.; Azimov, V. A.; Yakhontov, L. N. *Khim. Geterotsikl. Soedin.* **1982**, 3, 356-60; Azimov, V. A.; Bychikhina, N. N.; Yakhontov, L. N. *Khim. Geterotsikl. Soedin.* **1981**, 12, 1648-53; Spivey, A.C.; Fekner, T.; Spey,

20

S.E.; Adams, H. *J. Org. Chem.* 1999, 64(26), 9430-9443; Spivey, A.C.; Fekner, T.; Adams, H. *Tetrahedron Lett.* 1998, 39(48), 8919-8922. The methods described in Spivey et al. (preceding two references) for the preparation of 1-methyl-7-bromo-4-azaindoline can be used to prepare the 1-benzyl-7-bromo-4-azaindoline, 54, shown below in Scheme 21. This can be utilized in Stille or Suzuki couplings to provide 55, which is deprotected and dehydrogenated to provide 56. Other useful azaindole intermediates, such as the cyano derivatives, 57 and 58, and the aldehyde derivatives, 59 and 60, can then be further elaborated to compounds of Formula I.

10


Scheme 21



Alternatively the 7-functionalized 5-azaindole derivatives may be obtained by functionalization using the methodologies of T. Fukuda et.al. *Tetrahedron* 1999, 55, 9151 and M. Iwao et. Al. *Heterocycles* 1992, 34(5), 1031 described above for the 4 or 6 azaindoles. The 4 or 6 positions of the 5 aza indoles can be functionalized by using the azaindole N-oxide.

The conversion of indoles to indolines is well known in the art and can be carried out as shown or by the methods described in Somei, M.; Saida, Y.; Funamoto, T.; Ohta, T. *Chem. Pharm. Bull.* 1987, 35(8), 3146-54; M. Iwao et. Al. *Heterocycles* 1992, 34(5), 1031; and Akagi, M.; Ozaki, K. *Heterocycles* 1987, 26(1), 61-4.

Scheme 22



5           The preparation of azaindole oxoacetyl or oxo piperidines with carboxylic acids can be carried out from nitrile, aldehyde, or anion precursors via hydrolysis, oxidation, or trapping with  $CO_2$  respectively. As shown in the Scheme 22, Step 1, or the scheme below step a12 one method for forming the nitrile intermediate, 62, is by cyanide displacement of a halide in the aza-indole ring. The cyanide reagent used can be sodium cyanide, or more preferably copper or zinc cyanide. The reactions may be carried out in numerous solvents which are well known in the art. For example DMF is used in the case of copper cyanide. Additional procedures useful for carrying out step 1 of Scheme 24 are Yamaguchi, S.; Yoshida, M.; Miyajima, I.; Araki, T.; Hirai, Y.; *J. Heterocycl. Chem.* 1995, 32(5), 1517-1519 which describes methods for copper cyanide; Yutilov, Y.M.; Svertilova, I.A.; *Khim Geterotsikl Soedin* 1994, 8, 1071-1075 which utilizes potassium cyanide; and Prager, R.H.; Tsopelas, C.; Heisler, T.; *Aust. J. Chem.* 1991, 44 (2), 277-285 which utilizes copper cyanide in the presence of  $MeOS(O)_2F$ . The chloride or more preferably a bromide on the azaindole may be displaced by sodium cyanide in dioxane via the method described in *Synlett*. 1998, 3, 243-244. Alternatively, Nickel dibromide, Zinc, and triphenyl phosphine in can be used to activate aromatic and heteroaryl chlorides to displacement via potassium cyanide in THF or other suitable solvent by the methods described in Eur. Pat. Appl., 831083, 1998.

25           The conversion of the cyano intermediate, 62, to the carboxylic acid intermediate, 63, is depicted in step 2, Scheme 22 or in step a12, Scheme 23. Many methods for the conversion of nitriles to acids are well known in the art and may be employed. Suitable conditions for step 2 of Scheme 22 or the conversion of

intermediate 65 to intermediate 66 below employ potassium hydroxide, water, and an aqueous alcohol such as ethanol. Typically the reaction must be heated at refluxing temperatures for one to 100 h. Other procedures for hydrolysis include those described in:

5

Shiotani, S.; Taniguchi, K.; *J. Heterocycl. Chem.* **1997**, *34*(2), 493-499; Boogaard, A. T.; Pandit, U. K.; Koomen, G.-J.; *Tetrahedron* **1994**, *50*(8), 2551-2560; Rivalle, C.; Bisagni, E.; *Heterocycles* **1994**, *38*(2), 391-397; Macor, J.E.; Post, R.; Ryan, K.; *J. Heterocycl. Chem.* **1992**, *29*(6), 1465-1467.


10

The acid intermediate, 66 (Scheme 23), may then be esterified using conditions well known in the art. For example, reaction of the acid with diazomethane in an inert solvent such as ether, dioxane, or THF would give the methyl ester. Intermediate 67 may then be converted to intermediate 68 according to the procedure described in Scheme 2. Intermediate 68 may then be hydrolyzed to provide intermediate 69.

15


20

## Scheme 23




As shown in Scheme 24, step a13 another preparation of the  
 5 indoleoxoacetyl piperazine 7-carboxylic acids, 69, is carried out by oxidation of the  
 corresponding 7-carboxaldehyde, 70. Numerous oxidants are suitable for the  
 conversion of aldehyde to acid and many of these are described in standard organic  
 chemistry texts such as: Larock, Richard C., Comprehensive organic transformations  
 : a guide to functional group preparations 2<sup>nd</sup> ed. New York : Wiley-VCH, 1999. One  
 10 preferred method is the use of silver nitrate or silver oxide in a solvent such as  
 aqueous or anhydrous methanol at a temperature of ~25 °C or as high as reflux. The  
 reaction is typically carried out for one to 48 h and is typically monitored by TLC or  
 LC/MS until complete conversion of product to starting material has occurred.  
 Alternatively,  $\text{KMnO}_4$  or  $\text{CrO}_3/\text{H}_2\text{SO}_4$  could be utilized.

Scheme 24




Scheme 25 gives a specific example of the oxidation of an aldehyde intermediate, 70a, to provide the carboxylic acid intermediate, 69a.

Scheme 25




Alternatively, intermediate 69 can be prepared by the nitrile method of synthesis carried out in an alternative order as shown in Scheme 26. The nitrile hydrolysis step can be delayed and the nitrile carried through the synthesis to provide a nitrile which can be hydrolyzed to provide the free acid, 69, as above.

## Scheme 26



## Scheme 27



5

**Step H** The direct conversion of nitriles, such as 72, to amides, such as 73, shown in Scheme 27, Step H, can be carried out using the conditions as described in Shiotani, S.; Taniguchi, K.; *J. Heterocycl. Chem.* **1996**, *33*(4), 1051-1056 (describes the use of aqueous sulfuric acid); Memoli, K.A.; *Tetrahedron Lett.* **1996**, *37*(21), 3617-3618; Adolfsson, H.; Waernmark, K.; Moberg, C.; *J. Org. Chem.* **1994**, *59*(8), 2004-2009; and El Hadri, A.; Leclerc, G.; *J. Heterocycl. Chem.* **1993**, *30*(3), 631-635.

**Step I** For  $\text{NH}_2$

15


Shiotani, S.; Taniguchi, K.; *J. Heterocycl. Chem.* **1997**, *34*(2), 493-499; Boogaard, A. T.; Pandit, U. K.; Koomen, G.-J.; *Tetrahedron* **1994**, *50*(8), 2551-2560; Rivalle, C.; Bisagni, E.; *Heterocycles* **1994**, *38*(2), 391-397; Macor, J.E.; Post, R.; Ryan, K.; *J. Heterocycl. Chem.* **1992**, *29*(6), 1465-1467.

20




Step J

Scheme 28




## Scheme 28A

Either:





Prepared as in US 5,811,432

Or:




Prepared as in US 5,811,432

Tetrahedron Letters 1986, 27,837.




Scheme 29



Steps a16, a17, and a18 encompasses reactions and conditions for 1°, 2° and 3° amide bond formation as shown in Schemes 28 and 29 which provide compounds such as those of Formula 73.

The reaction conditions for the formation of amide bonds encompass any reagents that generate a reactive intermediate for activation of the carboxylic acid to amide formation, for example (but not limited to), acyl halide, from carbodiimide, acyl iminium salt, symmetrical anhydrides, mixed anhydrides (including phosphonic/phosphinic mixed anhydrides), active esters (including silyl ester, methyl ester and thioester), acyl carbonate, acyl azide, acyl sulfonate and acyloxy *N*-phosphonium salt. The reaction of the indole carboxylic acids with amines to form amides may be mediated by standard amide bond forming conditions described in the art. Some examples for amide bond formation are listed in references 41-53 but this list is not limiting. Some carboxylic acid to amine coupling reagents which are applicable are EDC, Diisopropylcarbodiimide or other carbodiimides, PyBop (benzotriazolyloxytris(dimethylamino) phosphonium hexafluorophosphate), 2-(1H-benzotriazole-1-yl)-1, 1, 3, 3-tetramethyl uronium hexafluorophosphate (HBTU). A particularly useful method for azaindole 7-carboxylic acid to amide reactions is the use of carbonyl imidazole as the coupling reagent as described in reference 53. The temperature of this reaction may be lower than in the cited reference, from 80 °C (or possibly lower) to 150 °C or higher. A more specific application is depicted in Scheme 30.

Scheme 30



5        The following four general methods provide a more detailed description for the preparation of indolecarboamides and these methods were employed for the synthesis of compounds of Formula I.

Method 1:

10

To a mixture of an acid intermediate, such as 69, (1 equiv., 0.48 mmol), an appropriate amine (4 equiv.) and DMAP (58 mg, 0.47 mmol) dissolved  $\text{CH}_2\text{Cl}_2$  (1 mL) was added EDC (90 mg, 0.47 mmol). The resulting mixture was shaken at rt for 12h, and then evaporated *in vacuo*. The residue was dissolved in MeOH, and  
15 subjected to preparative reverse phase HPLC purification.

Method 2:

To a mixture of an appropriate amine (4 equiv.) and HOBt (16 mg, 0.12 mmol) in THF (0.5 mL) was added an acid intermediate, such as 69, (25 mg, 0.06 mmol) and NMM (50  $\mu\text{L}$ , 0.45 mmol), followed by EDC (23 mg, 0.12 mmol). The reaction mixture was shaken at rt for 12 h. The volatiles were evaporated *in vacuo*; and the residue dissolved in MeOH and subjected to preparative reverse phase HPLC  
20 purification.

25

Method 3:

To a mixture of an acid intermediate, such as 69, (0.047 mmol), amine (4 equiv.) and DEPBT (prepared according to Li, H.; Jiang, X. Ye, Y.; Fan, C.; Todd, R.; Goodman, M. *Organic Letters* 1999, 1, 91; 21 mg, 0.071 mmol) in DMF (0.5 mL) was added TEA (0.03 mL, 0.22 mmol). The resulting mixture was shaken at rt for 12  
30

h; and then diluted with MeOH (2 mL) and purified by preparative reverse phase HPLC.

Method 4:

5


A mixture of an acid intermediate, such as 69, (0.047mmol) and 8.5 mg (0.052mmol) of 1,1-carbonyldiimidazole in anhydrous THF (2 mL) was heated to reflux under nitrogen. After 2.5h, 0.052 mmol of amine was added and heating continued. After an additional period of 3~20 h at reflux, the reaction mixture was cooled and concentrated in vacuo. The residue was purified by chromatography on silica gel to provide a compound of Formula I

In addition, the carboxylic acid may be converted to an acid chloride using reagents such as thionyl chloride (neat or in an inert solvent) or oxalyl chloride in a solvent such as benzene, toluene, THF, or CH<sub>2</sub>Cl<sub>2</sub>. The amides may alternatively, be formed by reaction of the acid chloride with an excess of ammonia, primary, or secondary amine in an inert solvent such as benzene, toluene, THF, or CH<sub>2</sub>Cl<sub>2</sub> or with stoichiometric amounts of amines in the presence of a tertiary amine such as triethylamine or a base such as pyridine or 2,6-lutidine. Alternatively, the acid chloride may be reacted with an amine under basic conditions (Usually sodium or potassium hydroxide) in solvent mixtures containing water and possibly a miscible co solvent such as dioxane or THF. Scheme 25B depicts a typical preparation of an acid chloride and derivatization to an amide of Formula I. Additionally, the carboxylic acid may be converted to an ester preferably a methyl or ethyl ester and then reacted with an amine. The ester may be formed by reaction with diazomethane or alternatively trimethylsilyl diazomethane using standard conditions which are well known in the art. References and procedures for using these or other ester forming reactions can be found in reference 52 or 54.

Additional references for the formation of amides from acids are: Norman, M.H.; Navas, F. III; Thompson, J.B.; Rigdon, G.C.; *J. Med. Chem.* 1996, 39(24), 4692-4703; Hong, F.; Pang, Y.-P.; Cusack, B.; Richelson, E.; *J. Chem. Soc., Perkin Trans 1* 1997, 14, 2083-2088; Langry, K.C.; *Org. Prep. Proc. Int.* 1994, 26(4), 429-438; Romero, D.L.; Morge, R.A.; Biles, C.; Berrios-Pena, N.; May, P.D.; Palmer, J.R.; Johnson, P.D.; Smith, H.W.; Busso, M.; Tan, C.-K.; Voorman, R.L.; Reusser, F.; Althaus, I.W.; Downey, K.M.; et al.; *J. Med. Chem.* 1994, 37(7), 999-1014;

Bhattacharjee, A.; Mukhopadhyay, R.; Bhattacharjya, A.; *Indian J. Chem., Sect B* 1994, 33(7), 679-682.

Scheme 31



5  
10  
Scheme 31 shows synthetic transformations on a chloro nitro azaindole. Step F-1 of Scheme 31 can be carried out according to the following procedures: Yamaguchi, S.; Yoshida, M.; Miyajima, I.; Araki, T.; Hirai, Y.; *J. Heterocycl. Chem.* 1995, 32(5), 1517-1519;

Yutilov, Y.M.; Svertilova, I. A.; *Khim Geterotsikl Soedin* 1994, 8, 1071-1075; and Prager, R.H.; Tsopelas, C.; Heisler, T.; *Aust. J. Chem.* 1991, 44(2), 277-285. Step F-2 of Scheme 31 may be accomplished according to the procedures set forth in:  
15 Ciurla, H.; Puszek, A.; *Khim Geterotsikl Soedin* 1996, 10, 1366-1371; Robinson, R.P.; Donahue, K.M.; Son, P.S.; Wagdy, S.D.; *J. Heterocycl. Chem.* 1996, 33(2), 287-293; Nicolai, E.; Claude, S.; Teulon, J. M.; *J. Heterocycl. Chem.* 1994, 31(1), 73-75; Hwu, J.R.; Wong, F.F.; Shiao, M.-J.; *J. Org. Chem.* 1992, 57(19), 5254-5255; Shiao, M.-J.; Lai, L.-L.; Ku, W.-S.; Lin, P.-Y.; Hwu, J.R.; *J. Org. Chem.* 1993, 58(17),  
20 4742-4744.

The introduction of an alkoxy or aryloxy substituent onto the azaindole (Step G, Scheme 31, R<sub>2</sub> is alkoxy or aryloxy) may be accomplished by the f procedures described in Klemm, L.H.; Zell, R.; *J. Heterocycl. Chem.* 1968, 5, 773; Bradsher, C. K.; Brown, F. C.; Porter, H. K.; *J. Am. Chem. Soc.* 1954, 76, 2357; and Hodgson, H. H.; Foster, C. K.; *J. Chem. Soc.* 1942, 581.

The introduction of a fluorine substituent onto the azaindole (Step G, Scheme 31) may be accomplished according to the procedures as described in Sanchez, J. P.; Gogliotti, R. D.; *J. Heterocycl. Chem.* 1993, 30(4), 855-859; Rocca, P.; Marsais, F.; Godard, A.; Queguiner, G.; *Tetrahedron Lett.* 1993, 34(18), 2937-2940; and Sanchez, J.P.; Rogowski, J.W.; *J. Heterocycl. Chem.* 1987, 24, 215.

The introduction of a chlorine substituent onto the azaindole (Step G, Scheme 31) may be accomplished according to the procedures as described in Ciurla, H.; Puszko, A.; *Khim Geterotsikl Soedin* 1996, 10, 1366-1371; Raveglia, L.F.; Giardina, G.A.M.; Grugni, M.; Rigolio, R.; Farina, C. ; *J. Heterocycl. Chem.* 1997, 34(2), 557-559; Matsumoto, J.I.; Miyamoto, T.; Minamida, A.; Mishimura, Y.; Egawa, H.; Mishimura, H.; *J. Med. Chem.* 1984, 27(3), 292; Lee, T.-C.; Salemnick, G.; *J. Org. Chem.* 1975, 24, 3608.


The introduction of a bromine substituent onto the azaindole (Step G, Scheme 31) may be accomplished according to the procedures as described in Raveglia, L.F.; Giardina, G.A.M.; Grugni, M.; Rigolio, R.; Farina, C. ; *J. Heterocycl. Chem.* 1997, 34(2), 557-559; Talik, T.; Talik, Z.; Ban-Oganowska, H.; *Synthesis* 1974, 293; Abramovitch, R. A.; Saha, M.; *Can. J. Chem.* 1966, 44, 1765.

It is well known in the art that heterocycles may be prepared from an aldehyde, carboxylic acid, carboxylic acid ester, carboxylic acid amide, carboxylic acid halide, or cyano moiety or attached to another carbon substituted by a bromide or other leaving group such as a triflate, mesylate, chloride, iodide, or phosponate. The methods for preparing such intermediates from intermediates typified by the carboxylic acid intermediate, 69, bromo intermediate, 76, or aldehyde intermediate, 70 described above are known by a typical chemist practitioner. The methods or types of heterocycles which may be constructed are described in the chemical literature. Some representative references for finding such heterocycles and their construction are included in reference 55 through 67 but should in no way be construed as limiting. However, examination of these references shows that many versatile methods are available for synthesizing diversely substituted heterocycles

and it is apparent to one skilled in the art that these can be applied to prepare compounds of Formula I. Chemists well versed in the art can now easily, quickly, and routinely find numerous reactions for preparing heterocycles, amides, oximes or other substituents from the above mentioned starting materials by searching for reactions or preparations using a conventional electronic database such as Scifinder (American Chemical Society), Crossfire (Beilstein), Theilheimer, or Reacccs (MDS). The reaction conditions identified by such a search can then be employed using the substrates described in this application to produce all of the compounds envisioned and covered by this invention. In the case of amides, commercially available amines can be used in the synthesis. Alternatively, the above mentioned search programs can be used to locate literature preparations of known amines or procedures to synthesize new amines. These procedures are then carried out by one with typical skill in the art to provide the compounds of Formula I for use as antiviral agents.


As shown below in Scheme 32, suitable substituted azaindoles, such as the bromoazaindole intermediate, 76, may undergo metal mediated couplings with aryl groups, heterocycles, or vinyl stannanes to provide compounds of Formula I wherein  $R_5$  is aryl, heteroaryl, or heteroalicyclic for example. The bromoazaindole intermediates, 76 (or azaindole triflates or iodides) may undergo Stille-type coupling with heteroarylstannanes as shown in Scheme 32, step a13. Conditions for this reaction are well known in the art and references 68-70 as well as reference 52 provide numerous conditions in addition to the specific examples provided in Scheme 14 and in the specific embodiments. It can be well recognized that an indole stannane could also couple to a heterocyclic or aryl halide or triflate to construct compounds of Formula I. Suzuki coupling (reference 71) between the bromo intermediate, 76, and a suitable boronate could also be employed and some specific examples are contained in this application.

Scheme 32






Scheme 33




As shown in Scheme 34, step a14, aldehyde intermediates, 70, may be used to generate numerous compounds of Formula I. The aldehyde group may be a precursor for any of the substituents R<sub>1</sub> through R<sub>6</sub>, but the transformation for R<sub>5</sub> is depicted above for simplicity. The aldehyde intermediate 70, may be reacted to become incorporated into a ring as

10

Scheme 34




described in the claims or be converted into an acyclic group. The aldehyde, 70, may be reacted with a Tosmic based reagent to generate oxazoles (references 42 and 43 for example). The aldehyde, 70, may be reacted with a Tosmic reagent and then an amine to give imidazoles as in reference 72 or the aldehyde intermediate, 70, may be reacted with hydroxylamine to give an oxime which is a compound of Formula I as described below. Oxidation of the oxime with NBS, t-butyl hypochlorite, or the other known reagents would provide the N-oxide which react with alkynes or 3 alkoxy vinyl esters to give isoxazoles of varying substitution. Reaction of the aldehyde intermediate 70, with the known reagent, 77 (reference 70) shown below under basic conditions would provide 4-aminotrityl oxazoles.



Removal of the trityl group would provide 4-amino oxazoles which could be substituted by acylation, reductive alkylation or alkylation reactions or heterocycle forming reactions. The trityl could be replaced with an alternate protecting group such as a monomethoxy trityl, CBZ, benzyl, or appropriate silyl group if desired. Reference 73 demonstrates the preparation of oxazoles containing a trifluoromethyl moiety and the conditions described therein demonstrates the synthesis of oxazoles with fluorinated methyl groups appended to them.


The aldehyde could also be reacted with a metal or Grignard (alkyl, aryl, or heteroaryl) to generate secondary alcohols. These would be efficacious or could be oxidized to the ketone with TPAP or  $\text{MnO}_2$  or PCC for example to provide ketones of Formula I which could be utilized for treatment or reacted with metal reagents to give tertiary alcohols or alternatively converted to oximes by reaction with hydroxylamine hydrochlorides in ethanolic solvents. Alternatively the aldehyde could be converted to benzyl amines via reductive amination. An example of oxazole formation via a Tosmic reagent is shown below in Scheme 35. The same reaction would work with aldehydes at other positions and also in the 5 and 6 aza indole series.

Scheme 35




Scheme 36 shows in step a15, a cyano intermediate, such as 62, which could be directly converted to compounds of Formula I via heterocycle formation or reaction with organometallic reagents.

### Scheme 36




Scheme 37 shows a method for acylation of a cyanoindole intermediate of formula 65 with oxalyl chloride which would give acid chloride, 79, which could then be coupled with the appropriate amine in the presence of base to provide 80.

### Scheme 37




The nitrile intermediate, 80, could be converted to the tetrazole of formula 81, which could then be alkylated with trimethylsilyldiazomethane to give the compound of formula 82 (Scheme 38).

Scheme 38



- 5 Tetrazole alkylation with alkyl halides would be carried out prior to azaindole acylation as shown in Scheme 39. Intermediate 65 could be converted to tetrazole, 83, which could be alkylated to provide 84. Intermediate 84 could then be acylated and hydrolyzed to provide 85 which could be subjected to amide formation conditions to provide 86. The group appended to the tetrazole may be quite diverse
- 10 and still exhibit impressive potency.

Scheme 39




15

Scheme 40 shows that an oxadiazole such as , 88, may be prepared by the addition of hydroxylamine to the nitrile, 80, followed by ring closure of intermediate 87 with phosgene. Alkylation of oxadiazole, 88, with trimethylsilyldiazomethane would give the compound of formula 89.

20

## Scheme 40




A 7-cyanoindole, such as 80, could be efficiently converted to the imidate ester under conventional Pinner conditions using 1,4-dioxane as the solvent. The imidate ester can be reacted with nitrogen, oxygen and sulfur nucleophiles to provide C7-substituted indoles, for example: imidazolines, benzimidazoles, azabenzimidazoles, oxazolines, oxadiazoles, thiazolines, triazoles, pyrimidines and amidines etc. For example the imidate may be reacted with acetyl hydrazide with heating in a nonparticipating solvent such as dioxane, THF, or benzene for example. (aqueous base or aqueous base in an alcoholic solvent may need to be added to effect final dehydrative cyclization in some cases) to form a methyl triazine. Other hydrazines can be used. Triazines can also be installed via coupling of stannyl triazines with 4,5,6, or 7-bromo or chloro azaindoles. The examples give an example of the formation of many of these heterocycles.

## References:

- (1) Das, B. P.; Boykin, D. W. *J. Med. Chem.* 1977, 20, 531.
- (2) Czarny, A.; Wilson, W. D.; Boykin, D. W. *J. Heterocyclic Chem.* 1996, 33, 1393.
- (3) Francesconi, I.; Wilson, W. D.; Tanious, F. A.; Hall, J. E.; Bender, B. C.; Tidwell, R. R.; McCurdy, D.; Boykin, D. W. *J. Med. Chem.* 1999, 42, 2260.


Scheme 41 shows addition of either hydroxylamine or hydroxylamine acetic acid to aldehyde intermediate 90 may give oximes of Formula 91.

**Scheme 41**




An acid may be a precursor for substituents R<sub>1</sub> through R<sub>5</sub> when it occupies the corresponding position such as R<sub>5</sub> as shown in Scheme 42.

## Scheme 41a




## Scheme 41a (continued)

5



## Scheme 42



5           An acid intermediate, such as 69, may be used as a versatile precursor to generate numerous substituted compounds. The acid could be converted to hydrazonyl bromide and then a pyrazole via reference 74. One method for general heterocycle synthesis would be to convert the acid to an alpha bromo ketone (ref 75) by conversion to the acid chloride using standard methods, reaction with  
10   diazomethane, and finally reaction with HBr. The alpha bromo ketone could be used to prepare many different compounds of Formula I as it can be converted to many heterocycles or other compounds of Formula I. Alpha amino ketones can be prepared by displacement of the bromide with amines. Alternatively, the alpha bromo ketone could be used to prepare heterocycles not available directly from the  
15   aldehyde or acid. For example, using the conditions of Hulton in reference 76 to react with the alpha bromo ketone would provide oxazoles. Reaction of the alpha bromoketone with urea via the methods of reference 77 would provide 2-amino oxazoles. The alpha bromoketone could also be used to generate furans using beta keto esters(ref 78-80) or other methods, pyrroles (from beta dicarbonyls as in ref 81 or by Hantzsch methods (ref 82) thiazoles, isoxazoles and imidazoles (ref 83) example  
20   using literature procedures. Coupling of the aforementioned acid chloride with N-methyl-O-methyl hydroxyl amine would provide a "Weinreb Amide" which could be used to react with alkyl lithiums or Grignard reagents to generate ketones. Reaction of the Weinreb anion with a dianion of a hydroxyl amine would generate isoxazoles  
25   (ref 84). Reaction with an acetylenic lithium or other carbanion would generate alkynyl indole ketones. Reaction of this alkynyl intermediate with diazomethane or other diazo compounds would give pyrazoles (ref 85). Reaction with azide or hydroxyl amine would give heterocycles after elimination of water. Nitrile oxides would react with the alkynyl ketone to give isoxazoles (ref 86). Reaction of the initial  
30   acid to provide an acid chloride using for example oxalyl chloride or thionyl chloride or triphenyl phosphine/ carbon tetrachloride provides a useful intermediate as noted above. Reaction of the acid chloride with an alpha ester substituted isocyanide and base would give 2-substituted oxazoles (ref 87). These could be converted to amines,




alcohols, or halides using standard reductions or Hoffman/Curtius type rearrangements.

5        Scheme 43 describes alternate chemistry for installing the oxoacetyl piperazine moiety onto the 3 position of the azaindoles. Step A''' in Scheme 43 depicts reaction with formaldehyde and dimethylamine using the conditions in Frydman, B.; Despuy, M.E.; Rapoport, H.; *J. Am. Chem. Soc.* **1965**, *87*, 3530 will provide the dimethylamino compound shown.

10        Step B''' shows displacement with potassium cyanide would provide the cyano derivative according to the method described in Miyashita, K.; Kondoh, K.; Tsuchiya, K.; Miyabe, H.; Imanishi, T.; *Chem. Pharm. Bull.* **1997**, *45*(5), 932-935 or in Kawase, M.; Sinhababu, A.K.; Borchardt, R.T.; *Chem. Pharm. Bull.* **1990**, *38*(11), 2939-2946. The same transformation could also be carried out using TMSCN and a  
15        tetrabutylammonium fluoride source as in Iwao, M.; Motoi, O.; *Tetrahedron Lett.* **1995**, *36*(33), 5929-5932. Sodium cyanide could also be utilized.

Scheme 43



5 Step C''' of Scheme 43 depicts hydrolysis of the nitrile with sodium hydroxide and methanol would provide the acid via the methods described in Iwao, M.; Motoi, O.; *Tetrahedron Lett.* 1995, 36(33), 5929-5932 for example. Other basic hydrolysis conditions using either NaOH or KOH as described in Thesing, J.; et al.; *Chem. Ber.* 1955, 88, 1295 and Geissman, T.A.; Armen, A.; *J. Am. Chem. Soc.* 1952, 10 74, 3916. The use of a nitrilase enzyme to achieve the same transformation is described by Klempier N, de Raadt A, Griengl H, Heinisch G, *J. Heterocycl. Chem.*, 1992 29, 93, and may be applicable.


15 Step D''' of Scheme 43 depicts an  $\alpha$  hydroxylation which may be accomplished by methods as described in Hanessian, S.; Wang, W.; Gai, Y.; *Tetrahedron Lett.* 1996, 37(42), 7477-7480; Robinson, R. A.; Clark, J. S.; Holmes, A. B.; *J. Am. Chem. Soc.* 1993, 115(22), 10400-10401 ( $KN(TMS)_2$  and then camphorsulfonyloxaziridine or another oxaziridine; and Davis, F.A.; Reddy, R.T.; Reddy, R.E.; *J. Org. Chem.* 1992, 57(24), 6387-6389.

Step E''' of Scheme 43 shows methods for the oxidation of the alpha hydroxy ester to the ketone which may be accomplished according to the methods described in Mohand, S.A.; Levina, A.; Muzart, J.; *Synth. Comm.* 1995, 25 (14), 2051-2059. A preferred method for step E''' is that of Ma, Z.; Bobbitt, J.M.; *J. Org. Chem.* 1991, 56(21), 6110-6114 which utilizes 4-(NH-Ac)-TEMPO in a solvent such as CH<sub>2</sub>Cl<sub>2</sub> in the presence of para toluenesulfonic acid. The method described in Corson, B.B.; Dodge, R.A.; Harris, S.A.; Hazen, R.K.; *Org. Synth.* 1941, 1, 241 for the oxidation of the alpha hydroxy ester to the ketone uses KmnO<sub>4</sub> as oxidant. Other methods for the oxidation of the alpha hydroxy ester to the ketone include those described in Hunaeus, ; Zincke,; *Ber. Dtsch Chem. Ges.* 1877, 10, 1489; Acree,; *Am. Chem.* 1913, 50, 391; and Claisen,; *Ber. Dtsch. Chem. Ges.* 1877, 10, 846.

Step F''' of Scheme 43 depicts the coupling reactions which may be carried out as described previously in the application and by a preferred method which is described in Li, H.; Jiang, X.; Ye, Y.-H.; Fan, C.; Romoff, T.; Goodman, M. *Organic Lett.*, 1999, 1, 91-93 and employs 3-(Diethoxyphosphoryloxy)-1,2,3-benzotriazin-4(3H)-one (DEPBT); a new coupling reagent with remarkable resistance to racemization.

20


Scheme 44



Scheme 44 depicts the preparation of Formula I compounds by coupling HWC(O)A to the acid as described in Step F''' of Scheme 43, followed by hydroxylation as in Step D''' of Scheme 43 and oxidation as described in Step E''' of Scheme 43.

25


Scheme 45



- Scheme 45 depicts a method for the preparation which could be used to obtain
- 5 amido compounds of Formula I. Step G' represents ester hydrolysis followed by amide formation (Step H' as described in Step F''' of Scheme 43). Step I' of Scheme 45 depicts the preparation of the N-oxide which could be accomplished according to the procedures in Suzuki, H.; Iwata, C.; Sakurai, K.; Tokumoto, K.; Takahashi, H.; Hanada, M.; Yokoyama, Y.; Murakami, Y.; *Tetrahedron* **1997**, *53*(5), 1593-1606;
  - 10 Suzuki, H.; Yokoyama, Y.; Miyagi, C.; Murakami, Y.; *Chem. Pharm. Bull.* **1991**, *39*(8), 2170-2172; and Ohmato, T.; Koike, K.; Sakamoto, Y.; *Chem. Pharm. Bull.* **1981**, *29*, 390. Cyanation of the N-oxide is shown in Step J' of Scheme 45 which

- may be accomplished according to Suzuki, H.; Iwata, C.; Sakurai, K.; Tokumoto, K.; Takahashi, H.; Hanada, M.; Yokoyama, Y.; Murakami, Y.; *Tetrahedron* **1997**, *53*(5), 1593-1606 and Suzuki, H.; Yokoyama, Y.; Miyagi, C.; Murakami, Y.; *Chem. Pharm. Bull.* **1991**, *39*(8), 2170-2172. Hydrolysis of the nitrile to the acid is depicted in Step
- 5 K' of Scheme 45 according to procedures such as Shiotani, S.; Tanigucchi, K.; *J. Heterocycl. Chem.* **1996**, *33*(4), 1051-1056; Memoli, K.A.; *Tetrahedron Lett.* **1996**, *37*(21), 3617-3618; Adolfsson, H.; Waernmark, K.; Moberg, C.; *J. Org. Chem.* **1994**, *59*(8), 2004-2009; and El Hadri, A.; Leclerc, G.; *J. Heterocycl. Chem.* **1993**, *30*(3), 631-635. Step L' of Scheme 45 depicts a method which could be utilized for the
- 10 preparation of amido compounds of Formula I from the cyano derivative which may be accomplished according to procedures described in Shiotani, S.; Taniguchi, K.; *J. Heterocycl. Chem.* **1997**, *34*(2), 493-499; Boogaard, A.T.; Pandit, U.K.; Koomen, G.-J.; *Tetrahedron* **1994**, *50*(8), 2551-2560; Rivalle, C.; Bisagni, E.; *Heterocycles* **1994**, *38*(2), 391-397; and Macor, J.E.; Post, R.; Ryan, K.; *J. Heterocycl. Chem.* **1992**,
- 15 *29*(6), 1465-1467. Step M' of Scheme 45 shows a method which could be used for the preparation of amido compounds of Formula I from the acid derivative which may be accomplished according to procedures described in Norman, M.H.; Navas, F. III; Thompson, J.B.; Rigdon, G.C.; *J. Med. Chem.* **1996**, *39*(24), 4692-4703; Hong, F.; Pang, Y.-P.; Cusack, B.; Richelson, E.; *J. Chem. Soc., Perkin Trans 1* **1997**, *14*,
- 20 2083-2088; Langry, K. C.; *Org. Prep. Proced. Int.* **1994**, *26*(4), 429-438; Romero, D.L.; Morge, R.A.; Biles, C.; Berrios-Pena, N.; May, P.D.; Palmer, J.R.; Johnson, P.D.; Smith, H.W.; Busso, M.; Tan, C.-K.; Voorman, R.L.; Reusser, F.; Althaus, I.W.; Downey, K.M.; et al.; *J. Med. Chem.* **1994**, *37*(7), 999-1014 and Bhattacharjee, A.; Mukhopadhyay, R.; Bhattacharjya, A.; *Indian J. Chem., Sect B* **1994**, *33*(7), 679-
- 25 682.


Scheme 46



Scheme 46 shows a method which could be used for the synthesis of an azaindole acetic acid derivative. Protection of the amine group could be effected by treatment with di-tert-butyl dicarbonate to introduce the t-Butoxycarbonyl (BOC) group. Introduction of the oxalate moiety may then be accomplished as shown in Step A of Scheme 46 according to the procedures described in Hewawasam, P.; Meanwell, N. A.; *Tetrahedron Lett.* 1994, 35(40), 7303-7306 (using t-Buli, or s-buli, THF); or Stanetty, P.; Koller, H.; Mihovilovic, M.; *J. Org. Chem.* 1992, 57(25), 6833-6837 (using t-Buli). The intermediate thus formed could then be cyclized to form the azaindole as shown in Step B of Scheme 46 according to the procedures described in Fuerstner, A.; Ernst, A.; Krause, H.; Ptock, A.; *Tetrahedron* 1996, 52(21), 7329-7344 (using  $\text{TiCl}_3$ , Zn, DME); or Fuerstner, A.; Hupperts, A.; *J. Am. Chem. Soc.* 1995, 117(16), 4468-4475 (using Zn, excess Tms-Cl,  $\text{TiCl}_3$  (cat.), MeCN).

15


Scheme 47



Scheme 47 describes an alternate synthesis which could be used to prepare azaindole acetic acid derivatives. Step C of Scheme 47 could be accomplished by using the procedures described in Harden, F.A.; Quinn, R.J.; Scammells, P.J.; *J. Med. Chem.* 1991, 34(9), 2892-2898 [use of 1.  $\text{NaNO}_2$ , conc.  $\text{HCl}$  2.  $\text{SnCl}_2$ , conc.  $\text{HCl}$  (cat.)]. Typically, 10 equivalents of  $\text{NaNO}_2$  and 1.0 equivalents of substrate reacted at 0 °C for 0.25 to 1h and to this reaction mixture was added 3.5 equivalents of  $\text{SnCl}_2$ . Alternatively, the procedure described in De Roos, K.B.; Salemink, C.A.; *Recl. Trav. Chim. Pays-Bas* 1971, 90, 1181 (use of  $\text{NaNO}_2$ , AcOH,  $\text{H}_2\text{O}$ ) could be used. The intermediate thus formed could be further reacted and cyclized to provide azaindole

- acetic acid derivatives as shown in Step D of Scheme 47 and according to the procedures described in Atkinson, C. M.; Mattocks, A. R.; *J. Chem. Soc.* **1957**, 3722; Ain Khan, M.; Ferreira Da Rocha, J.; *Heterocycles* **1978**, 9, 1617; Fusco, R.; Sannicola, F.; *Tetrahedron* **1980**, 36, 161 [use of HCl (conc)]; Abramovitch, R. A.; Spenser, I. D.; *Adv. Heterocycl. Chem.* **1964**, 3, 79 (use of ZnCl<sub>2</sub>, p-Cymene); and Clemo, G. R.; Holt, R. J. W.; *J. Chem. Soc.* **1953**, 1313; (use of ZnCl<sub>2</sub>, EtOH, Sealed tube).

Scheme 48




- Scheme 48 depicts another possible route to azaindole acetic acid derivatives. Step E of Scheme 48 could be carried out as shown or according to procedures such as those described in Yurovskaya, M.A.; Khamlova, I.G.; Nesterov, V.N.; Shishkin, O.V.; Struchkov, T.; *Khim Geterotsikl Soedin* **1995**, 11, 1543-1550; Grzegozek, M.; Wozniak, M.; Baranski, A.; Van Der Plas, H.C.; *J. Heterocycl. Chem.* **1991**, 28(4), 1075-1077 (use of NaOH, DMSO); Lawrence, N.J.; Liddle, J.; Jackson, D.A.; *Tetrahedron Lett.* **1995**, 36(46), 8477-8480 (use of NaH, DMSO); Haglund, O.; Nilsson, M.; *Synthesis* **1994**, 3, 242-244; (use of 2.5 equiv. CuCl, 3.5 equiv. TBu-OK, DME, Py); Makosza, M.; Sienkiewicz, K.; Wojciechowski, K.; *Synthesis* **1990**, 9, 850-852; (use of KO-tBu, DMF); Makosza, M.; Nizamov, S.; *Org. Prep. Proceed. Int.* **1997**, 29(6), 707-710; (use of tBu-OK, THF). Step F of Scheme 48 shows the cyclization reaction which could provide the azaindole acetic acid derivatives. This reaction could be accomplished according to procedures such as those described in Frydman, B.; Baldain, G.; Repetto, J. C.; *J. Org. Chem.* **1973**, 38, 1824 (use of H<sub>2</sub>, Pd-C, EtOH); Bistryakova, I. D.; Smirnova, N. M.; Safonova, T. S.; *Khim Geterotsikl Soedin* **1993**, 6, 800-803 (use of H<sub>2</sub>, Pd-C (cat.), MeOH); Taga, M.; Ohtsuka, H.; Inoue, I.; Kawaguchi, T.; Nomura, S.; Yamada, K.; Date, T.;

- Hiramatsu, H.; Sato, Y.; *Heterocycles* **1996**, *42*(1), 251-263 (use of  $\text{SnCl}_2$ ,  $\text{HCl}$ ,  $\text{Et}_2\text{O}$ ); Arcari, M.; Aveta, R.; Brandt, A.; Cecchetelli, L.; Corsi, G.B.; Dirella, M.; *Gazz. Chim. Ital.* **1991**, *121*(11), 499-504 [use of  $\text{Na}_2\text{S}_2\text{O}_6$ ,  $\text{THF}/\text{EtOH}/\text{H}_2\text{O}$  (2:2:1)];  
 5 Moody, C. J.; Rahimtoola, K. F.; *J. Chem. Soc., Perkin Trans 1* **1990**, 673 (use of  $\text{TiCl}_3$ ,  $\text{NH}_4\text{OAc}$ , acetone,  $\text{H}_2\text{O}$ ).

Scheme 49 provides another route to azaindole intermediates which could then be further elaborated to provide compounds of Formula I, such as the amido derivatives shown. Steps G'' and H'' of Scheme 49 may be carried out according to  
 10 the procedures described in Takahashi, K.; Shibasaki, K.; Ogura, K.; Iida, H.; *Chem. Lett.* **1983**, 859; and Itoh, N.; *Chem. Pharm. Bull.* **1962**, *10*, 55. Elaboration of the intermediate to the amido compound of Formula I could be accomplished as previously described for Steps I' - M' of Scheme 45.

15

Scheme 49




Scheme 50 shows the preparation of azaindole oxalic acid derivatives. The starting materials in Scheme 50 may be prepared according to *Tetrahedron Lett.* **1995**, *36*, 2389-2392. Steps A', B', C', and D' of Scheme 50 may be carried out  
 20 according to procedures described in Jones, R.A.; Pastor, J.; Siro, J.; Voro, T.N.; *Tetrahedron* **1997**, *53*(2), 479-486; and Singh, S.K.; Dekhane, M.; Le Hyaric, M.; Potier, P.; Dodd, R.H.; *Heterocycles* **1997**, *44*(1), 379-391. Step E' of Scheme 50 could be carried out according to the procedures described in Suzuki, H.; Iwata, C.; Sakurai, K.; Tokumoto, K.; Takahashi, H.; Hanada, M.; Yokoyama, Y.; Murakami,




- Y.; *Tetrahedron* **1997**, 53(5), 1593-1606; Suzuki, H.; Yokoyama, Y.; Miyagi, C.; Murakami, Y.; *Chem. Pharm. Bull.* **1991**, 39(8), 2170-2172; Hagen, T.J.; Narayanan, K.; Names, J.; Cook, J.M.; *J. Org. Chem.* **1989**, 54, 2170; Murakami, Y.; Yokoyama, Y.; Watanabe, T.; Aoki, C.; et al.; *Heterocycles* **1987**, 26, 875; and Hagen, T. J.;
- 5 Cook, J.M.; *Tetrahedron Lett.* **1988**, 29(20), 2421. Step F' of Scheme 50 shows the conversion of the phenol to a fluoro, chloro or bromo derivative. Conversion of the phenol to the fluoro derivative could be carried out according to procedures described in Christe, K.O.; Pavlath, A.E.; *J. Org. Chem.* **1965**, 30, 3170; Murakami, Y.; Aoyama, Y.; Nakanishi, S.; *Chem. Lett.* **1976**, 857; Christe, K. O.; Pavlath, A. E.; *J.*
- 10 *Org. Chem.* **1965**, 30, 4104; and Christe, K.O.; Pavlath, A.E.; *J. Org. Chem.* **1966**, 31, 559. Conversion of the phenol to the chloro derivative could be carried out according to procedures described in Wright, S.W.; *Org. Prep. Proc. Int.* **1997**, 29(1), 128-131; Hartmann, H.; Schulze, M.; Guenther, R.; *Dyes Pigm* **1991**, 16(2), 119-136; Bay, E.; Bak, D. A.; Timony, P. E.; Leone-Bay, A.; *J. Org. Chem.* **1990**, 55, 3415;
- 15 Hoffmann, H.; et al.; *Chem. Ber.* **1962**, 95, 523; and Vanallan, J.A.; Reynolds, G.A.; *J. Org. Chem.* **1963**, 28, 1022. Conversion of the phenol to the bromo derivative may be carried out according to procedures described in Katritzky, A.R.; Li, J.; Stevens, C.V.; Ager, D.J.; *Org. Prep. Proc. Int.* **1994**, 26(4), 439-444; Judice, J.K.; Keipert, S.J.; Cram, D.J.; *J. Chem. Soc., Chem. Commun.* **1993**, 17, 1323-1325; Schaeffer,
- 20 J.P.; Higgins, J.; *J. Org. Chem.* **1967**, 32, 1607; Wiley, G.A.; Hershkowitz, R.L.; Rein, R.M.; Chung, B.C.; *J. Am. Chem. Soc.* **1964**, 86, 964; and Tayaka, H.; Akutagawa, S.; Noyori, R.; *Org. Syn.* **1988**, 67, 20.

Scheme 50




Scheme 51 describes methods for the preparation of azaindole acetic acid derivatives by the same methods employed for the preparation of azaindole oxalic acid derivatives as shown and described in Scheme 50 above. The starting material employed in Scheme 51 could be prepared according to *J. Org. Chem.* 1999, 64, 7788-7801. Steps A'', B'', C'', D'', and E'' of Scheme 51 could be carried out in the same fashion as previously described for Steps A', B', C', D', and E' of Scheme 50.

Scheme 51




The remaining schemes provide additional background, examples, and conditions for carrying out this invention. Specific methods for preparing W and modifying A are presented. As shown in Scheme 52, the azaindoles may be treated with oxalyl chloride in either THF or ether to afford the desired glyoxyl chlorides according to literature procedures (Lingens, F.; Lange, J. *Justus Liebigs Ann. Chem.* 1970, 738, 46-53). The intermediate glyoxyl chlorides may be coupled with benzoyl piperazines (Desai, M.; Watthey, J.W. *Org. Prep. Proc. Int.* 1976, 8, 85-86) under basic conditions to afford compounds of Formula I directly.

## Scheme 52




- 5 Alternatively, **Scheme 52** treatment of the azaindole-3-glyoxyl chloride, (Scheme 52) with *tert*-butyl 1-piperazinecarboxylate affords the piperazine coupled product. It is apparent to one skilled in the art that use of an alternative Boc protected piperazine which are synthesized as shown below would provide compounds of formula I with alternative groups of formula W. As discussed earlier, other amine
- 10 protecting groups which do not require acidic deprotection conditions could be utilized if desired. Deprotection of the Boc group is effected with 20% TFA/CH<sub>2</sub>Cl<sub>2</sub> to yield the free piperazine. This product is then coupled with carboxylic acid in the presence of polymer supported 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide (P-EDC) to afford products of Formula I. This sequence provides a general method for
- 15 synthesizing compounds of varied A in formula I.

Scheme 53



An example for preparing compounds of Formula I which possess  
5 substituents in A (or other parts of the molecule) which might interfere with the  
standard reaction schemes reactions is shown in Scheme 53. The piperazine  
derivative (Scheme 53) was treated with Boc-protected aminobenzoic acid in the  
presence of EDC to afford the piperazine diamide. A portion of the resulting product  
was separated and subjected to TFA in order to remove the Boc group, thus yielding  
10 amino derivatives.




## Scheme 54



Similarly, substituents which possess a reactive alcohol can be incorporated as  
5 below. The piperazine derivative (Scheme 54) was treated with acetoxybenzoic acid  
in the presence of EDC to afford the piperazine diamide derivative. A portion of the  
resulting product was separated and subjected to LiOH hydrolysis in order to remove  
the acetate group, thus yielding hydroxy derivatives.


10 Examples containing substituted piperazines are prepared using the general  
procedures outlined in Schemes 55-38. Substituted piperazines are either  
commercially available from Aldrich, Co. or prepared according to literature  
procedures (Behun et al, Ref. 88(a), Scheme 31, eq. 01). Hydrogenation of alkyl  
substituted pyrazines under 40 to 50 psi pressure in EtOH afforded substituted  
15 piperazines. When the substituent was an ester or amide, the pyrazine systems could  
be partially reduced to the tetrahydropyrazine (Rossen et al, Ref. 88(b), Scheme 55,  
eq. 02). The carbonyl substituted piperazines could be obtained under the same  
conditions described above by using commercially available dibenzyl piperazines  
(Scheme 55, eq. 03).

Scheme 55

X = OR, NR<sub>1</sub>R<sub>2</sub>X = OR, NR<sub>1</sub>R<sub>2</sub>


- 2-Trifluoromethylpiperazine (Jenneskens et al., Ref. 88c) was prepared through a four step route (Scheme 56). Using Lewis acid TiCl<sub>4</sub>, N,N'-
- 5 dibenzylethylenediamine reacted with trifluoropyruvates to afford the hemiacetal, which was reduced at room temperature by Et<sub>3</sub>SiH in TFA to afford the lactam. LiAlH<sub>4</sub> treatment then reduced the lactam to 1,4-dibenzyl-2-trifluoromethylpiperazine. Finally, hydrogenation of the dibenzyl-2-trifluoromethylpiperazine in HOAc gave the desired product, 2-
- 10 trifluoromethylpiperazine.

## Scheme 56



Mono-benzoylation of symmetric substituted piperazines could be achieved by using one of the following procedures (Scheme 57). (a) Treatment of a solution of piperazine in acetic acid with acetyl chloride afforded the desired mon-benzoylated piperazine (Desai et al. Ref. 27, Scheme 57, eq. 04). (b) Symmetric piperazines were treated with 2 equivalents of *n*-butyllithium, followed by the addition of benzoyl chloride at room temperature (Wang et al, Ref. 89, Scheme 57, eq. 05).

## Scheme 57




Mono-benzoylation of unsymmetric substituted piperazines could be achieved by using one of the following procedures (Scheme 57), in which all the methods were exemplified by mono-alkyl substituted piperazines. (a) Unsymmetric piperazines were treated with 2 equivalents of *n*-butyllithium, followed by the addition of benzoyl chloride at room temperature to afford a mixture of two regioisomers, which could be separated by chromatography (Wang et al, Ref. 89 and 90(b), Scheme 58 eq. 06); (b) Benzoic acid was converted to its pentafluorophenyl



- ester, and then further reaction with 2-alkylpiperazine to provide the mono-benzoylpiperazines with the benzoyl group at the less hindered nitrogen (Adamczyk et al, Ref. 90(a), Scheme 58, eq. 07); (c) A mixture of piperazine and methyl benzoate was treated with dialkylaluminum chloride in methylene chloride for 2-4 days to yield the mono-benzoylpiperazine with the benzoyl group at the less hindered nitrogen (Scheme 58 eq. 08); (d) Unsymmetric piperazines were treated with 2 equivalents of *n*-butyllithium, followed by subsequent addition of triethylsilyl chloride and benzoyl chloride in THF at room temperature to afford mono-benzoylpiperazines with the benzoyl group at the more hindered nitrogen (Wang et al, Ref. 90(b), Scheme 58, eq. 09). When the substituent at position 2 was a ester or amide, the mono-benzoylation with benzoyl chloride occurred at the less hindered nitrogen of the piperazine with triethylamine as base in THF (Scheme 58, eq. 10).


Scheme 58



15


In the case of tetrahydropyrazines (Scheme 59, eq. 11), mono-benzoylation occurred at the more hindered nitrogen under the same conditions as those in equation 10 of Scheme 58, in the well preceded manner. (Rossen et al, Ref. 88(b)).

### Scheme 59



Furthermore, the ester group can be selectively reduced by  $\text{NaBH}_4$  in the  
5 presence of the benzamide (Masuzawa et al, Ref. 91), which is shown in Scheme 60.


### Scheme 60



10

The ester groups on either the piperazine linkers or on the azaindole nucleus could be hydrolyzed to the corresponding acid under basic conditions such as  $K_2CO_3$  (Scheme 61, eq. 13) or NaOMe (Scheme 61, eq. 14) as bases in MeOH and water.


### Scheme 61



Reaction of an azaindole glyoxyl chloride with substituted benzoyl piperazines or tetrahydropyrazines in  $\text{CH}_2\text{Cl}_2$  using  $\text{I-Pr}_2\text{NEt}$  as base afforded the coupled products as shown in Scheme 62.

5


Scheme 62



In the case of coupling reactions using 3-hydroxymethyl-benzoylpiperazine, the hydroxyl group was temporarily protected as its TMS ether with BSTFA (N,O-bis(trimethylsilyl)trifluoroacetamide) (Furber et al, Ref. 92). The unprotected nitrogen atom can then be reacted with glyoxyl chlorides to form the desired diamides. During workup, the TMS masking group was removed to give free hydroxymethylpiperazine diamides as shown in Scheme 63.

15


Scheme 63



Piperazine intermediates were prepared using standard chemistry as shown in

20 Scheme 64.

Scheme 64




5

Scheme 65

10 Scheme 65 depicts some more specific methodology for preparing 5-azindoles for use in preparation of the claimed compounds. Some reductive cyclizations conditions include Fe in acetic acid, Tin II chloride in aq HCl, or Zinc powder in acetic acid. Hydrogenation conditions or other conditions used in LeimGruber-Batch indole synthesis sequences can also be employed.

**More specific route to 5-azindoles:**




X= chloro or bromo or may be converted to a substituent and then carried through the sequence

15


Tautomers of nitrogen containing heterocycles are covered by this patent application. For example, a hydroxy pyrazine is also known to represent its corresponding tautomer as well as shown in Scheme 66.

Scheme 66




5 Scheme 67-74 provides some nonlimiting methodology for the preparation of substituted pyrazines which can be incorporated into substituents of compounds of claim 1, particularly as part of R<sup>4</sup>. It should be noted that the nomenclature in these schemes does not coincide with that of the claims but rather shows examples of methods which can be used to prepare pieces which make up the compounds of the claims. Thus R<sub>1</sub> and R<sub>2</sub> in these schemes does not refer to the R1 and R2 in the  
10 claims but for example refers to chemically compatible groups which might be envisioned by chemists skilled in the art and which can be utilized to prepare compounds of the claims.


## Scheme 67



## Scheme 68




## Scheme 69






**Scheme 70**  
**Thiazol**


A




B.




C.




## Scheme 71




## Scheme 72



## Scheme 73



## Scheme 74



- Throughout the chemistry discussion, chemical transformations which are well known in the art have been discussed. The average practitioner in the art knows these transformations well and a comprehensive list of useful conditions for nearly all the transformations is available to organic chemists and this list is contained in reference 52 authored by Larock and is incorporated in its entirety for the synthesis of compounds of Formula I.

**Chemistry****General:**

- 5 Additional preparations of starting materials and intermediates are contained in Wang et. al. PCT WO 01/62255 which is incorporated by reference.

**Chemistry**

- 10 All Liquid Chromatography (LC) data were recorded on a Shimadzu LC-10AS liquid chromatograph using a SPD-10AV UV-Vis detector with Mass Spectrometry (MS) data determined using a Micromass Platform for LC in electrospray mode.

15 LC/MS Method (i.e., compound identification)

Column A: YMC ODS-A S7 3.0x50 mm column

Column B: PHX-LUNA C18 4.6x30 mm column

20

Column C: XTERRA ms C18 4.6x30 mm column

Column D: YMC ODS-A C18 4.6x30 mm column

25 Column E: YMC ODS-A C18 4.6x33 mm column

Column F: YMC C18 S5 4.6x50 mm column

Column G: XTERRA C18 S7 3.0x50 mm column

30

Gradient: 100% Solvent A / 0% Solvent B to 0% Solvent A / 100%  
Solvent B

Solvent A = 10% MeOH - 90% H<sub>2</sub>O - 0.1% TFA, Solvent B = 90% MeOH - 10%  
H<sub>2</sub>O - 0.1% TFA; and R<sub>t</sub> in min.

35 Gradient time: 2 minutes

Hold time 1 minute

Flow rate: 5 mL/min

Detector Wavelength: 220 nm

5 Solvent A: 10% MeOH / 90% H<sub>2</sub>O / 0.1% Trifluoroacetic Acid

Solvent B: 10% H<sub>2</sub>O / 90% MeOH / 0.1% Trifluoroacetic Acid

10 Compounds purified by preparative HPLC were diluted in MeOH (1.2 mL) and purified using the following methods on a Shimadzu LC-10A automated preparative HPLC system or on a Shimadzu LC-8A automated preparative HPLC system with detector (SPD-10AV UV-VIS) wavelength and solvent systems (A and B) the same as above.

15 Preparative HPLC Method (i.e., compound purification)

Purification Method: Initial gradient (40% B, 60% A) ramp to final gradient (100% B, 0% A) over 20 minutes, hold for 3 minutes (100% B, 0% A)

20 Solvent A: 10% MeOH / 90% H<sub>2</sub>O / 0.1% Trifluoroacetic Acid

Solvent B: 10% H<sub>2</sub>O / 90% MeOH / 0.1% Trifluoroacetic Acid

Column: YMC C18 S5 20x100 mm column

25


Detector Wavelength: 220 nm

## Typical Procedures and Characterization of Selected Examples:

### Preparation of Intermediates:

5


#### Intermediate 1



- 10 4-Methoxyphenylboronic acid (24.54 g), 4-chloro-3-nitropyridine hydrochloride (26.24 g),  $\text{Pd}(\text{Ph}_3\text{P})_4$  (4 g) and  $\text{K}_2\text{CO}_3$  (111 g) were combined in DME (500 mL). The reaction was heated to reflux for 10 hours. After the mixture cooled down to room temperature, it was poured into saturated aqueous  $\text{NH}_4\text{OAc}$  (500 mL) solution. The aqueous phase was extracted with EtOAc (3 x 200 mL). The combined extract was concentrated to give a residue which was purified using silica
- 15 gel chromatography (10% to 30% EtOAc / PE) to afford 10.6 g of Intermediate 1, 3-Nitro-4-(4-methoxyphenyl)pyridine. MS  $m/z$ :  $(\text{M}+\text{H})^+$  calcd for  $\text{C}_{12}\text{H}_{11}\text{N}_2\text{O}_3$ : 231.08; found 231.02. HPLC retention time: 1.07 minutes (column B).

#### Intermediate 1a

Alternate route to 5-azaindoles:




20

2-methoxy-5-bromo pyridine can be purchased from Aldrich (or others) or prepared. Oxidation with 1.1eq of MCPBA in dichloromethane (20ml per 10.6 mmol bromide) in the presence of anhydrous  $\text{MgSO}_4$  (0.4g per mL dichloromethane) with stirring



from 0° to ambient temperature for approximately 14 h provided the N-oxide after workup and flash chromatographic purification over silica gel using a 5% EtOAc/Hexane gradient of increasing EtOAc. The N-oxide (1.6g) was dissolved in 10mL 98% sulfuric acid and cooled to 0°. 10 mL of 69% nitric acid was added and  
5 then allowed to warm to ambient temp with stirring. The reaction was then heated and stirred at 80° C for 14h and then poured over ice, extracted with dichloromethane, washed with water, and concentrated to give a yellow solid which was purified by flash chromatography over Silica gel using 1:1EtOAc/hexane and then a gradient to provide a yellow crystalline solid: ). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.50  
10 (s, 1H), 7.59 (s, 1H), 4.12 (3H, s). LC MS showed desired M+H. The N-oxide was reduced by dissolving the starting material in dichloromethane (0.147M substrate) and cooling to 0°. A solution of 1.2 eq PCl<sub>3</sub> (0.44M) in dichloromethane was added slowly to keep the reaction at 0°. Warm to ambient temp and stir for 72h. Aqueous workup and concentration provided a yellow solid which could be used in subsequent  
15 reactions or purified by chromatography. Note: a similar sequence could be used with 2-methoxy-5-chloro-pyridine as starting material.

### Intermediate 2a




20

*Typical procedure for preparing azaindole from nitropyridine:* Preparation of 7-chloro-6-azaindole, Intermediate 2a, is an example of Step A of Scheme 1. 2-chloro-3-nitropyridine (5.0g, 31.5mmol) was dissolved in dry THF (200 mL). After the solution was cooled to -78 °C, vinyl magnesium bromide (1.0M in THF, 100 mL)  
25 was added dropwise. The reaction temperature was maintained at -78°C for 1 h, and then at -20 °C for another 12 h before it was quenched by addition of 20% NH<sub>4</sub>Cl aqueous solution (150 mL). The aqueous phase was extracted with EtOAc (3 x 150 mL). The combined organic layer was dried over MgSO<sub>4</sub>, filtered and the filtrate was concentrated *in vacuo* to give a residue which was purified by silica gel column  
30 chromatography (EtOAc / Hexane, 1 / 10) to afford 1.5g (31%) of 7-chloro-6-azaindole, Intermediate 2a. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 7.84 (d, 1H, *J* = 10.7 Hz), 7.55 (dd, 1H, *J* = 10.9, 5.45 Hz), 6.62 (d, 1H, *J* = 5.54 Hz), 4.89 (s, 1H). MS *m/z*:

(M+H)<sup>+</sup> calcd for C<sub>7</sub>H<sub>6</sub>ClN<sub>2</sub>: 153.02; found 152.93. HPLC retention time: 0.43 minutes (column A).

### Intermediate 2b


5



Intermediate 2b, 7-(4-Methoxyphenyl)-4-azaindole, was prepared by the same method as Intermediate 2a starting from 3-Nitro-4-(4-methoxyphenyl)pyridine, Intermediate 1. MS *m/z*: (M+H)<sup>+</sup> calcd for C<sub>14</sub>H<sub>13</sub>N<sub>2</sub>O: 225.10; found 225.02. HPLC retention time: 1.39 minutes (column B).

10

### Intermediate 2c




15

Intermediate 2c, 4-bromo-7-chloro-6-azaindole, was prepared by the same method as Intermediate 2a, starting from 2-Chloro-3-nitro-5-bromo-pyridine (available from Aldrich, Co.). MS *m/z*: (M+H)<sup>+</sup> calcd for C<sub>7</sub>H<sub>5</sub>BrClN<sub>2</sub>: 230.93; found 231.15. HPLC retention time: 1.62 minutes (column B).


20

### Intermediate 2d



25

Intermediate 2d, 4-fluoro-7-chloro-6-azaindole (above), was prepared according to the following scheme:



A) fuming  $\text{HNO}_3$ ,  $\text{H}_2\text{SO}_4$ ;

B)  $\text{POCl}_3/\text{DMF}$ ,  $110^\circ\text{C}$ ;

C) vinylmagnesium bromide, THF,  $-78^\circ\text{C} \sim -20^\circ\text{C}$

5 It should be noted that 2-chloro-5-fluoro-3-nitro pyridine, **zz3'**, may be prepared by the method in example 5B of the reference Marfat, A. ; and Robinson, R. P. ; "Azaoxindole Derivatives" *US. Patent* 5,811,432 1998. The preparation below provides some details which enhance the yields of this route.

10 In Step A, compound **zz1'** (1.2 g, 0.01 mol) was dissolved in sulfuric acid (2.7 mL) at room temperature. Premixed fuming nitric acid (1 mL) and sulfuric acid was added dropwise at  $5\text{--}10^\circ\text{C}$  to the solution of compound **zz1'**. The reaction mixture was then heated at  $85^\circ\text{C}$  for 1 hour, then was cooled to room temperature and poured into ice (20 g). The yellow solid precipitate was collected by filtration, washed with  
15 water and air dried to provide 1.01 g of compound **zz2'**.


In Step B, compound **zz2'** (500 mg, 3.16 mmol) was dissolved in phosphorous oxychloride (1.7 mL, 18.9 mmol) and dimethoxyethane at room temperature. The reaction was heated to  $110^\circ\text{C}$  for 5 hours. The excess  
20 phosphorous oxychloride was then removed by concentrating the reaction mixture in vacuo. The residue was chromatographed on silica gel, eluted with chloroform (100%) to afford 176 mg of product **zz3'**.

In Step C, compound **zz3'** (140 mg, 0.79 mmol) was dissolved in THF (5 mL)  
25 and cooled to  $-78^\circ\text{C}$  under a nitrogen atmosphere. To this solution was added dropwise a solution of vinyl magnesium bromide (1.2 mmol, 1.0 M in diethyl ether, 1.2 mL). The reaction mixture was then kept at  $-20^\circ\text{C}$  for 15 hours. The reaction mixture was then quenched with saturated ammonium chloride, and extracted with

ethyl acetate. The combined organic layers were washed with brine, dried over magnesium sulfate, filtered, and the filtrate was concentrated in vacuo. The residue was chromatographed on silica to provide 130 mg of intermediate 2i. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 7.78 (s, 1H), 7.60 (d, 1H, *J* = 3.0 Hz), 6.71 (d, 1H, *J* = 3.05 Hz). MS *m/z*: (M+H)<sup>+</sup> calcd for C<sub>7</sub>H<sub>5</sub>ClFN<sub>2</sub>: 171.10; found 171.00. HPLC retention time: 1.22 minutes (column A).

Intermediate 2d, 4-fluoro-7-chloro-6-azaindole, was prepared by the same method as Intermediate 2a, starting from 2-Chloro-3-nitro-5-fluoro-pyridine which was prepared according to the procedure above. Experimental details for this preparation are contained in Wang et. al. PCT WO 01/62255. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 7.78 (s, 1H), 7.60 (d, 1H, *J* = 3.0 Hz), 6.71 (d, 1H, *J* = 3.05 Hz). MS *m/z*: (M+H)<sup>+</sup> calcd for C<sub>7</sub>H<sub>5</sub>ClFN<sub>2</sub>: 171.10; found 171.00. HPLC retention time: 1.22 minutes (column A).

#### Intermediate 2e




Intermediate 2e was prepared by either Method A or Method B, below:

Method A: A mixture of 4-bromo-7-chloro-6-azaindole (1 g), CuI (0.65 g) and NaOMe (4 mL, 25% in methanol) in MeOH (16 mL) was heated at 110 – 120 °C for 16 hours in a sealed tube. After cooling to room temperature, the reaction mixture was neutralized with 1N HCl to pH 7. The aqueous solution was extracted with EtOAc (3 x 30 mL). Then the combined organic layer was dried over MgSO<sub>4</sub>, filtered and the filtrate was concentrated *in vacuo* to afford a residue, which was purified by using silica gel chromatography to give 0.3 g of 4-methoxy-7-chloro-6-azaindole, Intermediate 2e. MS *m/z*: (M+H)<sup>+</sup> calcd for C<sub>8</sub>H<sub>8</sub>ClN<sub>2</sub>O: 183.03; found 183.09. HPLC retention time: 1.02 minutes (column B).

Method B :A mixture of 4-bromo-7-chloro-6-azaindole (6 g), CuBr (3.7 g) and NaOMe (30 mL, 5% in MeOH) was heated at 110°C for 24 hours in a sealed

tube. After cooling to room temperature, the reaction mixture was added to saturated aqueous  $\text{NH}_4\text{Cl}$ . The resulting aqueous solution was extracted with EtOAc (3 x 30 mL). The combined organic layer was dried over  $\text{MgSO}_4$ , filtered and the filtrate was concentrated *in vacuo* to afford a residue, which was purified by using silica gel chromatography to give 1.8 g of 4-methoxy-7-chloro-6-azaindole, Intermediate 2e.

#### Intermediate 2f




10

Intermediate 2f, 7-bromo-6-azaindole was prepared by the same method as Intermediate 2a, starting from 2-Bromo-3-nitro-pyridine (available from Aldrich, Co.). MS  $m/z$ :  $(\text{M}+\text{H})^+$  calcd for  $\text{C}_7\text{H}_5\text{BrN}_2$ : 197.97; found 197.01. HPLC retention time: 0.50 minutes (column A).

15

#### Intermediate 2g




20

Intermediate 2g, 7-chloro-4-azaindole was prepared by the same method as Intermediate 2a, starting from 4-Chloro-3-nitro-pyridine (HCl salt, available from Austin Chemical Company, Inc.). MS  $m/z$ :  $(\text{M}+\text{H})^+$  calcd for  $\text{C}_7\text{H}_5\text{ClN}_2$ : 153.02; found 152.90. HPLC retention time: 0.45 minutes (column A).

25

#### Intermediate 2h




Intermediate 2h, 5-chloro-7-methyl-4-azaindole was prepared by the same method as Intermediate 2a, starting from 2-Chloro-4-methyl-5-nitro-pyridine

30

(available from Aldrich, Co.). MS  $m/z$ :  $(M+H)^+$  calcd for  $C_8H_8ClN_2$ : 167.04; found 166.99. HPLC retention time: 1.22 minutes (column B).


### Example 2i

5



Intermediate 2j, 4-fluoro-7-bromo-6-azaindole, was prepared by the same method as Intermediate 2e, using  $POBr_3$  in the step B instead of  $POCl_3$ . MS  $m/z$ :  
10  $(M+H)^+$  calcd for  $C_7H_5BrFN_2$ : 214.96; found 214.97. HPLC retention time: 1.28 minutes (column G).

### Intermediate 2j



15

To a mixture of 5-bromo-2-chloro-3-nitropyridine (10 g, 42 mmol) in 1,4-dioxane (100 ml) was added pyrazole (5.8 g, 85 mmol). The resulting mixture was stirred at 120°C for 26.5 h., and then evaporated after cooling to r.t. The crude material was  
20 purified by flash chromatography (0 to 5% EtOAc/Hexanes) to give the desired product 5-Bromo-3-nitro-2-pyrazol-1-yl-pyridine.  $^1H$  NMR: ( $CD_3OD$ )  $\delta$  8.77 (s, 1H), 8.56 (s, 1H), 8.45 (s, 1H), 7.73 (s, 1H), 6.57 (s, 1H); LC/MS: (ES+)  $m/z$   $(M+H)^+ = 269, 271$ , HPLC  $R_t = 1.223$ .

25 To a 250 mL round bottom flask was charged 5-Bromo-3-nitro-2-pyrazol-1-yl-pyridine (1.02 g, 3.8 mmol) and THF (30 ml). The mixture was then cooled to  $-78^\circ C$ , and added a THF solution of vinylmagnesium bromide (23 mL, 18.4 mmol, 0.8 M). After three minutes, the reaction mixture was warmed to  $-45^\circ C$  and remained


stirring for 1 h. The reaction was then quenched with ammonium chloride, and the resulting mixture extracted with EtOAc. The combined extracts were evaporated *in vacuo*, and the residue purified by flash column chromatography (5% EtOAc/Hexanes) to give compound 2 (which by HPLC contained about 50% of a side product, presumably 3-vinylamino of compound 1); <sup>1</sup>H NMR: (CDCl<sub>3</sub>) δ 10.75 (b s, 1H), 8.73 (s, 1H), 8.10 (s, 1H), 7.82 (s, 1H), 7.52 (s, 1H), 6.67 (s, 1H), 6.53 (s, 1H); LC/MS: (ES+) m/z (M+H) = 262, 264; HPLC R<sub>t</sub> = 1.670.

**Intermediate 2k**

10

To a solution of 2-bromo-5-chloro-3-nitropyridine 5 (20 g, 84 mmol, prepared in 2 steps from 2-amino-5-chloropyridine as described in WO9622990) in THF (300 ml) at -78°C was charged a THF solution of vinylmagnesium bromide (280 ml, 252 mmol, 0.9 M). The resulting mixture was stirred at -78°C for one hour, followed by quenching with aqueous ammonium chloride (500 ml, sat.) and extracted with EtOAc (5 x 500 ml). The combined organic extracts were washed with aqueous ammonium chloride (2 x 500 ml, sat.) and water (3 x 500 ml), dried (MgSO<sub>4</sub>) and evaporated to give a brownish residue. The crude material was triturated with CH<sub>2</sub>Cl<sub>2</sub>, and the solid formed filtered to give compound 6 as a yellow solid (8.0 g, 41%); <sup>1</sup>H NMR: (DMSO-*d*<sub>6</sub>) 12.30 (b s, 1H), 7.99 (s, 1H), 7.80 (d, *J* = 3.0, 1H), 6.71 (d, *J* = 3.0, 1H); LC/MS: (ES+) m/z (M+H)<sup>+</sup> = 231, 233, 235; HPLC R<sub>t</sub> = 1.833.


20

**Intermediate 2m**


25

4-Fluoro-7-Bromo-6-azaindole (500 mg, 1.74 mmol) was dissolved in THF (5ml) and cooled to -78°C and *n*-BuLi (2.5 M, 2.1 ml) was added dropwise. The reaction mixture was stirred at -78°C for 15 min, then stirred at 0°C for 30 min. The reaction

was cooled to  $-78^{\circ}\text{C}$  again, and DMF (0.7 ml, 8.7 mmol) was added. After stirring for 30 min, water was added to quench the reaction. The reaction mixture was extracted with ethylacetate. The organic layer was dried over  $\text{MgSO}_4$ , filtered, concentrated and chromatographed to afford 208 mg of intermediate 2m. LC/MS:  $(\text{ES}^+) m/z (M+H)^+ =$   
5 164.98.  $R_t = 0.44$  min.


**Intermediate 2n**

10 A mixture of intermediate 2m (50 mg, 0.30 mmol), potassium carbonate (42 mg, 0.30 mmol) and tosylmethyl isocyanide (60 mg, 0.30 mmol) in MeOH (3 ml) was heated to reflux for about 2 hr. The solvent was removed in vacuum and the residue was treated with ice water and extracted with ether. The organic layer was washed with an aqueous solution of HCl (2%), water and dried over magnesium sulfate. After  
15 filtration and evaporation of the solvent, the residue was purified on silica to afford the title compound (60 mg). LC/MS:  $(\text{ES}^+) m/z (M+H)^+ = 204$ .  $R_t = 0.77$  min.


**Intermediate 2o**

20 4-Fluoro-7-Bromo-6-azaindole (510 mg, 2.39 mmol) in anhydrous DMF (5 mL) was treated with copper cyanide (430 mg, 4.8 mmol) at  $150^{\circ}\text{C}$  in a seal tube for 1 h. An aqueous solution of  $\text{NH}_4\text{OH}$  (10 mL) was added and the reaction was extracted with diethylether (2 x 50 mL) and ethylacetate (2 x 50 mL). The organic phases were  
25 combined and dried over sodium sulfate, filtered, concentrated in vacuum and chromatographed on silica gel (gradient elution AcOEt/Hexanes 0-30%) to afford the title compound as a brownish solid (255 mg, 66%) LC/MS:  $(\text{ES}^+) m/z (M+H)^+ = 162$ .




**Intermediate 2p**

Intermediate 2o (82 mg, 0.51 mmol) was dissolved in absolute ethanol (200% proof, 5 mL) and treated with hydroxylamine hydrochloride (53 mg, 0.76 mmol) and triethylamine (140  $\mu$ L, 1.0 mmol) and the reaction mixture was heated up at 80°C in a seal tube for 2h. The solvent was removed in vacuum and the pale yellow solid residue was washed with water to afford the title compound. LC/MS: (ES<sup>+</sup>) m/z (M+H)<sup>+</sup> = 195. This compound was taken to the next step without further purification.

**Intermediate 2q**


Intermediate 2p was dissolved in trimethylorthoformate (1 mL) and heated at 85°C in a seal tube for 1h, then it was cooled to rt, the solvent was removed in vacuum and the residue was chromatographed on silica gel (AcOEt/Hexanes, gradient elution 10-60%) to afford the title compound (54 mg, LC/MS: (ES<sup>+</sup>) m/z (M+H)<sup>+</sup> = 205).

**Intermediate 2r**

Intermediate 2q (100 mg, 0.62 mmol, crude) in ethanol (5 mL) was treated with an aqueous solution of sodium hydroxide (50%, 2 mL) and the reaction mixture was heated at 110°C overnight in a seal tube. The pH was adjusted to 2 with HCl (6N)


and a brown precipitate was filtered off. The solution was concentrated to dryness to afford the title compound as a pale yellow solid LC/MS: (ES<sup>+</sup>) m/z (M+H)<sup>+</sup> =181. This compound was used without further purification.

5


**Intermediate 2s**

Intermediate 2r (0.62 mmol) was dissolved in DMF (1 mL) and treated with 3-aminopyridine (58.3 mg, 0.62 mmol), DEBT (185 mg, 0.62) and Hunig's base (216  
10  $\mu$ L, 1.26 mmol) and the reaction mixture was stirred at room temperature for 18h. Water was added and the reaction was extracted with AcOEt (2 x 25 mL) and CHCl<sub>3</sub> (2 x 25 mL), dried over sodium sulfate, concentrated and chromatographed on silica gel (AcOEt/Hexanes gradient elution 0-50%) to afford the title compound as a brownish solid LC/MS: (ES<sup>+</sup>) m/z (M+H)<sup>+</sup> =257.


15

**Intermediate 2s**


Intermediate 2h, 4-methoxy-7-bromo-5-azaindole was prepared by the same method  
20 as Intermediate 2a, starting from 2-methoxy-5-bromo-4-nitro-pyridine (intermediate 1a). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  8.52 (s, 1H), 7.84 (s, 1H), 7.12 (t, 1H), 6.68 (d, 1H), 3.99 (s, 3H). LC MS showed desired M+H.

**Intermediate 2t**


- A mixture of aldehyde intermediate 2m (150 mg, 0.91 mmol), sodium cyanide (44mg, 0.091mmol) and tosylmethyl isocyanide (177 mg, 0.91 mmol) in EtOH(3ml) was stirred at room temperature for 30min, then filtered and the crystals were washed with ether-hexane (1:1) and dried. The obtained crystals, and a saturated solution of ammonia in dry methanol (8ml) were heated between 100-110°C for 16hr. The mixture was concentrated and chromatographed to provide 20mg of intermediate 2.
- LC/MS: (ES<sup>+</sup>) m/z(m+H)<sup>+</sup> = 203. Rt = 0.64 min.

**Intermediate 3a**

- Typical procedure for acylation of azaindole: Preparation of Methyl (7-chloro-6-azaindol-3-yl)-oxoacetate, Intermediate 3a is an example of Step B of Scheme 1. 7-Chloro-6-azaindole, Intermediate 2a (0.5 g, 3.3 mmol) was added to a suspension of AlCl<sub>3</sub> (2.2 g, 16.3 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (100 mL). Stirring was continued at rt for 10 minutes before methyl chlorooxoacetate (2.0 g, 16.3 mmol) was added dropwise. The reaction was stirred for 8 h. The reaction was quenched with iced aqueous NH<sub>4</sub>OAc solution (10%, 200 mL). The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 100mL). The combined organic layer was dried over MgSO<sub>4</sub>, filtered and the filtrate was concentrated *in vacuo* to give a residue which was carried to the next step without further purification. Intermediate 2, Methyl (7-chloro-6-azaindol-3-yl)-oxoacetate: MS m/z: (M+H)<sup>+</sup> calcd for C<sub>10</sub>H<sub>5</sub>ClN<sub>2</sub>O<sub>3</sub>; 239.02; found 238.97. HPLC retention time: 1.07 minutes (column A).


**Intermediate 3b**

Intermediate 3b, Methyl (6-azaindol-3-yl)-oxoacetate, was prepared by the same method as Intermediate 3a, starting from 6-azaindole. MS  $m/z$ : (M+H)<sup>+</sup> calcd for C<sub>10</sub>H<sub>7</sub>N<sub>2</sub>O<sub>3</sub>: 205.06; found 205.14. HPLC retention time: 0.49 minutes (column A).


**Intermediate 3c**

Intermediate 3c, Methyl (7-(4-methoxyphenyl)-4-azaindol-3-yl)-oxoacetate, was prepared by the same method as Intermediate 3a, starting from 7-(4-methoxyphenyl)-4-azaindole (Intermediate 2b). MS  $m/z$ : (M+H)<sup>+</sup> calcd for C<sub>17</sub>H<sub>15</sub>N<sub>2</sub>O<sub>4</sub>: 311.10; found 311.04. HPLC retention time: 1.15 minutes (column A).

15


**Intermediate 3d**

Intermediate 3d, methyl (7-chloro-4-methoxy-6-azaindol-3-yl)-oxoacetate was prepared by the same method as Intermediate 3a, starting from Intermediate 2e, 4-methoxy-7-chloro-6-azaindole. MS  $m/z$ : (M+H)<sup>+</sup> calcd for C<sub>12</sub>H<sub>12</sub>ClN<sub>2</sub>O<sub>4</sub>: 283.05; found 283.22. HPLC retention time: 1.37 minutes (column B).

**Intermediate 3e**


5 Intermediate 3e, Methyl (7-chloro-4-fluoro-6-azaindol-3-yl)-oxoacetate was prepared by the same method as Intermediate 3a starting from Intermediate 2d, 4-fluoro-7-chloro-6-azaindole.  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.63 (s, 1H), 8.00 (s, 1H), 3.95 (s, 3H). MS  $m/z$ :  $(\text{M}+\text{H})^+$  calcd for  $\text{C}_{10}\text{H}_7\text{ClFN}_2\text{O}_3$ : 257.01; found 257.00. HPLC retention time: 1.26 minutes (column A).

10

**Intermediate 3f**

15 Intermediate 3f, Methyl (7-chloro-4-azaindol-3-yl)-oxoacetate was prepared by the same method as Intermediate 3a, starting from Intermediate 2g, 7-chloro-4-azaindole. MS  $m/z$ :  $(\text{M}+\text{H})^+$  calcd for  $\text{C}_{10}\text{H}_8\text{ClN}_2\text{O}_3$ : 239.02; found 238.97. HPLC retention time: 0.60 minutes (column A).


20

**Intermediate 3g**


Intermediate 3g, Methyl (5-chloro-7-methyl-4-azaindol-3-yl)-oxoacetate was prepared by the same method as Intermediate 3a, starting from Intermediate 2h, 5-chloro-7-methyl-4-azaindole. MS  $m/z$ :  $(M+H)^+$  calcd for  $C_{11}H_{10}ClN_2O_3$ : 253.04; found 252.97. HPLC retention time: 1.48 minutes (column B).

5

Intermediate 4a




*Typical procedure of hydrolysis of ester:* Preparation of Potassium (7-chloro-6-azaindol-3-yl)-oxoacetate, Intermediate 4a, is an example of Step C of Scheme 1. Crude methyl (7-chloro-6-azaindol-3-yl)-oxoacetate, Intermediate 3a, and an excess of  $K_2CO_3$  (2 g) were dissolved in MeOH (20 mL) and  $H_2O$  (20 mL). After 8 h, the solution was concentrated and the residue was purified by silica gel column chromatography to provide 200 mg of Potassium (7-chloro-6-azaindol-3-yl)-oxoacetate. MS  $m/z$ :  $(M+H)^+$  of the corresponding acid was observed. Calc'd for  $C_9H_6ClN_2O_3$ : 225.01; found 225.05. HPLC retention time: 0.83 minutes (column A).

**Intermediate 4b**


5 Potassium (6-azaindol-3-yl)oxoacetate, Intermediate 4b, was prepared by the same method as Intermediate 4a, starting from Methyl (6-azaindol-3-yl)oxoacetate, Intermediate 3b. MS *m/z*: (M+H)<sup>+</sup> of the corresponding acid was observed. Calc'd for C<sub>9</sub>H<sub>7</sub>N<sub>2</sub>O<sub>3</sub>: 191.05; Found 190.99. HPLC retention time: 0.12 minutes (column A).

10

**Intermediate 4c**


15 Intermediate 4c, Potassium (7-(4-methoxyphenyl)-4-azaindol-3-yl)-oxoacetate, was prepared by the same method as Intermediate 4a, starting from Methyl (7-(4-methoxyphenyl)-4-azaindol-3-yl)-oxoacetate, Intermediate 3c. MS *m/z*: (M-K+H)<sup>+</sup> calcd for C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>4</sub>: 297.07; found 297.04. HPLC retention time: 1.00 minutes (column A).

20

**Intermediate 4d**

Intermediate 4d, Potassium (7-chloro-4-methoxy-6-azaindol-3-yl)-oxoacetate was prepared by the same method as Intermediate 4a starting from Methyl (7-chloro-4-methoxy-6-azaindol-3-yl)-oxoacetate, Intermediate 3d. MS  $m/z$ :  $(M+H)^+$  of the corresponding acid of compound 4d  $(M-K+H)^+$  calcd for  $C_{10}H_8ClN_2O_4$ : 255.02; found 255.07. HPLC retention time: 0.74 minutes (column A).

#### Intermediate 4e




10

Intermediate 4e, Potassium (7-chloro-4-azaindol-3-yl)-oxoacetate was prepared by the same method as Intermediate 4a, starting from Methyl (7-chloro-4-azaindol-3-yl)-oxoacetate, Intermediate 3f. MS  $m/z$ :  $(M+H)^+$  of the corresponding acid of compound 4e  $(M-K+H)^+$  calcd for  $C_9H_6ClN_2O_3$ : 225.01; found 225.27. HPLC retention time: 0.33 minutes (column A).

15

#### Intermediate 4f




20

Intermediate 4f, Potassium (5-chloro-7-methyl-4-azaindol-3-yl)-oxoacetate was prepared by the same method as Intermediate 4a, starting from Methyl (5-chloro-7-methyl-4-azaindol-3-yl)-oxoacetate, Intermediate 3g. MS  $m/z$ :  $(M+H)^+$  of the corresponding acid of compound 4f  $(M-K+H)^+$  calcd for  $C_{10}H_8ClN_2O_3$ : 239.02; found 238.94. HPLC retention time: 1.24 minutes (column B).


25



**Intermediate 4g**

5 Intermediate 4g, Potassium (7-bromo-6-azaindol-3-yl)-oxoacetate was prepared by the same method as Intermediate 4a, starting from Methyl (7-bromo-6-azaindol-3-yl)-oxoacetate (prepared according to the method of Intermediate 3a from 7-Bromo-6-azaindole, Intermediate 2f).  $^1\text{H}$  NMR (500 MHz,  $\text{DMSO-d}_6$ )  $\delta$  8.59 (s, 1H), 8.16 (d, 1H,  $J = 5.3$  Hz), 8.08 (d, 1H,  $J = 5.45$  Hz);  $^{13}\text{C}$  NMR (125 MHz,  $\text{DMSO-d}_6$ )  $\delta$  180.5, 164.0, 141.6, 140.4, 132.4, 125.3, 115.5, 113.0.

10


**Intermediate 4h**

15

Intermediate 4h, Potassium (7-bromo-4-fluoro-6-azaindol-3-yl)-oxoacetate was prepared by the same method as Intermediate 4a, starting from Methyl (7-bromo-4-fluoro-6-azaindol-3-yl)-oxoacetate (prepared according to the method of Intermediate 3a from 7-Bromo-4-fluoro-6-azaindole, Intermediate 2i). MS  $m/z$ : (M+H) $^+$  of the corresponding acid of compound 4g (M-H) $^+$  calcd for  $\text{C}_9\text{H}_5\text{BrFN}_2\text{O}_3$ : 286.95; found 286.94. HPLC retention time: 0.94 minutes (column A).

20


## Intermediate 4I




- 5 1-ethyl-3-methylimidazolium chloride (0.172 g, 1.1 mmol) was added to aluminum chloride (0.560 g, 4.2 mmol), and the mixture vigorously stirred. Upon formation of a liquid, intermediate 2j was added, followed by ethyl chlorooxoacetate (0.12 ml, 1.1 mmol). The mixture was allowed to stir at r.t. for 16 h, after which additional chlorooxoacetate was added (0.12 ml, 1.1 mmol). Following this addition, the
- 10 reaction was allowed to stir at r.t. for another 24 h. The flask was cooled to 0°C and water added, upon which precipitates were formed. The solid material was filtered, washed with water and methanol, and dried under high vacuum to give compound 3; LC/MS: (ES+) m/z (M+H) = 334, 336; HPLC R<sub>t</sub> = 1.390.

15


## Intermediate 4j



- To 1-ethyl-3-methylimidazolium chloride (2.54 g, 17.3 mmol) was added aluminum chloride (6.91 g, 51.8 mmol). The mixture was stirred vigorously at ambient temperature for ten minutes. To the resulting yellow liquid was added intermediate 2k (2.0 g, 8.64 mmol) and ethyl chlorooxoacetate (2.0 ml, 17.3 mmol), and was stirred at ambient temperature for 16 h. The reaction mixture was then added ice/water (300 ml) to give precipitates, which were filtered and washed with water to give the title
- 20 compound as a yellow solid (1.98 g). The aqueous solution was extracted with EtOAc (3 x 300 ml), and the extracts evaporated *in vacuo* to give a second batch of compound 8 as a yellow solid (439 mg, total yield 92%); <sup>1</sup>H NMR: (DMSO-*d*<sub>6</sub>) 14.25 (b s, 1H), 13.37 (s, 1H), 8.56 (s, 1H), 8.18 (s, 1H); LC/MS: (ES+) m/z (M+H)<sup>+</sup> = 303, 305, 307; HPLC R<sub>t</sub> = 1.360.


**Intermediate 4k**

1-Ethyl-3-methylimidazolium chloride (82mg, 0.56 mmol) was added to a flask  
5 which contained intermediate 2n (56 mg, 0.28 mmol) and the mixture was cooled to  
0°C. Aluminum chloride (336 mg, 2.52 mmol) was added in one portion followed by  
ClCOCOOEt (58  $\mu$ L, 0.56 mmol) and the reaction mixture was stirred at room  
temperature for 2 days. Ice water was added to quench the reaction. The reaction  
mixture was filtered. The solid was washed with water and diethylether and dried in  
10 air to afford the title compound (58mg). LC/MS: (ES<sup>+</sup>) m/z (M+H)<sup>+</sup> = 276. Rt = 0.85  
min.


**Intermediate 4m**

15

1-Ethyl-3-methylimidazolium chloride (73mg, 0.52 mmol) and aluminum chloride  
(198 mg, 1.56 mmol) were stirred together under nitrogen for 1h. To this solution  
was added intermediate 2q (54 mg, 0.26 mmol) and ethyloxalylchloride ( 58  $\mu$ L, 0.52  
mmol) and the reaction mixture was stirred at rt for 18h. The reaction was quenched  
15 with water and the mixture was stirred for 15 min. The solid was collected by  
filtration and washed with water and diethylether. LC/MS (ES<sup>+</sup>) m/z (M+H)<sup>+</sup> =276.  
This compound was used without further purification.

**Intermediate 4n**


1-Ethyl-3-methylimidazolium chloride (26mg, 0.18 mmol) was added to a flask  
 5 which contained intermediate 2t (18 mg, 0.09 mmol) and the mixture was cooled to  
 0°C. Aluminum chloride (92 mg, 0.54mmol) was added in one portion followed by  
 ClCOCOOEt (20μL, 0.18 mmol) and the reaction mixture was stirred at room  
 temperature for 2 days. Ice water was added to quench the reaction. The reaction  
 mixture was filtered. The solid was washed with water and diethylether and dried in  
 10 air to afford compound D (18mg). LC/MS: (ES<sup>+</sup>) m/z(m+H)<sup>+</sup> = 275. Rt = 0.49 min.

**Intermediate 5a**

15


*Typical procedure for coupling piperazine derivative and azaindole acid:*

Preparation of 1-benzoyl-3-(R)-methyl-4-[(7-chloro-6-azaindol-3-yl)-  
 oxoacetyl]piperazine, Intermediate 5, is an example of Step D of Scheme 1.  
 Potassium 7-chloro-6-azaindole 3-glyoxylate, Intermediate 4a, (100 mg, 0.44 mmol),  
 20 3-(R)-methyl-1-benzoylpiperazine (107 mg, 0.44 mol), 3-(diethoxyphosphoryloxy)-  
 1,2,3-benzotriazin-4(3H)-one (DEPBT) (101 mg, 0.44 mol) and Hunig's Base  
 (diisopropylethylamine, 0.5 mL) were combined in 5 mL of DMF. The mixture was  
 stirred at rt for 8 h. DMF was removed *via* evaporation at reduced pressure and the  
 residue was purified using a Shimadzu automated preparative HPLC System to give  
 25 1-(benzoyl)-3-(R)-methyl-4-[(7-chloro-6-azaindol-3-yl)-oxoacetyl]-piperazine (70  
 mg, 39%). MS m/z: (M+H)<sup>+</sup> Calc'd for C<sub>21</sub>H<sub>20</sub>ClN<sub>4</sub>O<sub>3</sub>: 411.12; Found 411.06. HPLC  
 retention time: 1.32 minutes (column A).

**Intermediate 5b**


5 Intermediate 5b, 1-benzoyl-4-[(7-chloro-4-methoxy-6-azaindol-3-yl)-oxoacetyl]piperazine was prepared by the same method as Intermediate 5a starting from Potassium (7-chloro-4-methoxy-6-azaindol-3-yl)-oxoacetate, Intermediate 4d, and 1-benzoylpiperazine. MS  $m/z$ :  $(M+H)^+$  calcd for  $C_{21}H_{20}ClN_4O_4$ : 427.12; found 427.12. HPLC retention time: 1.28 minutes (column A).

10

**Intermediate 5c**


15 Intermediate 5c, 1-benzoyl-3-(R)-methyl-4-[(7-chloro-4-methoxy-6-azaindol-3-yl)-oxoacetyl]piperazine was prepared by the same method as Intermediate 5a starting from Potassium (7-chloro-4-methoxy-6-azaindol-3-yl)-oxoacetate, Intermediate 4d, and 1-benzoylpiperazine.  $^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$  8.10 (s, 1H), 7.72 (s, 1H), 7.40 (s, 5H), 3.89 (s, 3H), 3.71 – 3.40 (m, 8H). MS  $m/z$ :  $(M+H)^+$  calcd for  $C_{22}H_{22}ClN_4O_4$ : 441.13; found 441.17. HPLC retention time: 1.33 minutes (column A).

20

**Intermediate 5d**


5 Intermediate 5d, 1-benzoyl-3-(R)-methyl-4-[(7-chloro-4-azaindol-3-yl)-oxoacetyl]piperazine was prepared by the same method as Intermediate 5a, starting from Potassium (7-chloro-4-azaindol-3-yl)-oxoacetate, Intermediate 4e, and 1-benzoyl-3-(R)-methyl piperazine. MS  $m/z$ : (M+H)<sup>+</sup> calcd for C<sub>21</sub>H<sub>20</sub>ClN<sub>4</sub>O<sub>3</sub>, 411.12, found 411.04. HPLC retention time: 1.10 minutes (column A).

10

**Intermediate 5e**


15 Intermediate 5e, 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-methyl-4-azaindol-3-yl)-oxoacetyl]piperazine was prepared by the same method as Intermediate 5a, starting from Potassium (5-chloro-7-methyl-4-azaindol-3-yl)-oxoacetate, Intermediate 4f, and 1-benzoyl-3-(R)-methyl piperazine. MS  $m/z$ : (M+H)<sup>+</sup> calcd for C<sub>22</sub>H<sub>22</sub>ClN<sub>4</sub>O<sub>3</sub>, 425.24, found 425.04. HPLC retention time: 1.72 minutes (column B).

20

**Intermediate 5f**

Intermediate 5f, 1-benzoyl-3-(R)-methyl-4-[(7-bromo-6-azaindol-3-yl)-oxoacetyl]piperazine was prepared by the same method as Intermediate 5a, starting from (7-bromo-6-azaindol-3-yl)-oxoacetic acid potassium salt, Intermediate 4g, and 1-benzoyl-3-(R)-methylpiperazine. MS  $m/z$ :  $(M+H)^+$  calcd for  $C_{21}H_{20}BrN_4O_3$ : 455.07; found 455.14. HPLC retention time: 1.45 minutes (column B).

#### Intermediate 5g




10

Intermediate 5g, 1-benzoyl-4-[(7-bromo-6-azaindol-3-yl)-oxoacetyl]piperazine was prepared by the same method as Intermediate 5a, starting from (7-bromo-6-azaindol-3-yl)-oxoacetic acid potassium salt, Intermediate 4g, and 1-benzoylpiperazine. MS  $m/z$ :  $(M+H)^+$  calcd for  $C_{20}H_{18}BrN_4O_3$ : 441.06; found 441.07. HPLC retention time: 1.43 minutes (column B).

15


#### Intermediate 5h




20

Intermediate 5h, 1-benzoyl-3-(R)-methyl-4-[(6-azaindol-3-yl)-oxoacetyl]piperazine was prepared by the same method as Intermediate 5a starting from Potassium (6-azaindol-3-yl)oxoacetate, Intermediate 4b, and 1-benzoyl-3-(R)-methylpiperazine. MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{21}H_{21}N_4O_3$ : 377.16; Found 377.10. HPLC retention time: 0.88 minutes (column A).

25

Intermediate 5i

5 Addition of intermediate 2d to a solution of aluminum trichloride in dichloromethane stirring at ambient temperature followed 30 minutes later with chloromethyl or chloroethyl oxalate (according to the method described for intermediate 3a) provides either the methyl or ethyl ester, respectively. Hydrolysis with KOH (as in the standard hydrolysis procedure described for intermediate 4a) provided potassium (7-chloro-4-fluoro-6-azaindol-3-yl)oxoacetate. Potassium (7-chloro-4-fluoro-6-azaindol-3-yl)oxoacetate was then reacted with 1-benzoyl  
10 piperazine in the presence of DEPBT under the standard conditions (as described for intermediate 5a) to provide 1-benzoyl-4-[(4-fluoro-7-chloro-6-azaindol-3-yl)-oxoacetyl]piperazine, intermediate 5i. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 8.40 (s, 1H), 8.04 (s, 1H), 7.46 (bs, 5H), 3.80-3.50 (m, 8H); LC/MS (ES<sup>+</sup>) m/z (M+H)<sup>+</sup> 415  
15 observed; retention time 1.247 minutes; LC/MS method: YMC ODS-A C18 S7 3.0 x 50 mm column; Start %B = 0, Final %B = 100, Gradient time = 2 minutes; Flow rate = 5 mL/min; detector wavelength = 220 nm.


Intermediate 5j

20 1-benzoyl-3-(R)-methyl-4-[(4-fluoro-7-chloro-6-azaindol-3-yl)-oxoacetyl]piperazine was prepared by coupling potassium (7-chloro-4-fluoro-6-azaindol-3-yl)oxoacetate, prepared as described above for intermediate 5i, with 1-benzoyl-3-(R)-methylpiperazine in the presence of DEPBT under the standard conditions (as




described for intermediate 5a) to provide 1-benzoyl-3-(R)-methyl-4-[(4-fluoro-7-chloro-6-azaindol-3-yl)-oxoacetyl] piperazine, intermediate 5j. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 8.42, 8.37 (s, s, 1H), 8.03 (s, 1H), 7.71-7.45 (m, 5H), 4.72-3.05 (m, 7H), 1.45-1.28 (m, 3H); LC/MS (ES<sup>+</sup>) m/z (M+H)<sup>+</sup> 429 observed; retention time 1.297 minutes; LC/MS method: YMC ODS-A C18 S7 3.0 x 50 mm column; Start %B = 0, Final %B = 100, Gradient time = 2 minutes; Flow rate = 5 mL/min; detector wavelength = 220 nm.

### Intermediate 5k



Intermediate 5k, 1-benzoyl-4-[(7-chloro-6-azaindol-3-yl)-oxoacetyl]piperazine was prepared by the same method as Intermediate 5a, starting from (7-chloro-6-azaindol-3-yl)-oxoacetic acid potassium salt, Intermediate 4a, and 1-benzoylpiperazine. MS m/z: (M+H)<sup>+</sup> calcd for C<sub>20</sub>H<sub>18</sub>ClN<sub>4</sub>O<sub>3</sub>: 397.11; found 396.97. HPLC retention time: 2.37 minutes (column F, gradient time = 3 min, flow rate = 4 ml/min).

### Intermediate 5l




Intermediate 5l, 1-picolinoyl-4-[(4-methoxy-7-chloro-6-azaindol-3-yl)-oxoacetyl]piperazine was prepared by the same method as Intermediate 5a starting from Potassium (4-methoxy-7-chloro-6-azaindol-3-yl)oxoacetate, Intermediate 4d, and picolinoyl-piperazine. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 8.63 – 7.45 (m, 7 H),

3.94 (s, 3H), 3.82 – 2.50 (m, 8H). MS  $m/z$ : (M+H)<sup>+</sup> Calc'd for C<sub>20</sub>H<sub>19</sub>ClN<sub>5</sub>O<sub>4</sub>: 428.11; Found 428.11. HPLC retention time: 1.09 minutes (column A).

### Intermediate 5m


5



Intermediate 5m, (R)-1-picolinoyl-3-methyl-4-[(7-bromo-6-azaindol-3-yl)-oxoacetyl]piperazine was prepared by the same method as Intermediate 5a starting from Potassium (7-bromo-6-azaindol-3-yl)oxoacetate, Intermediate 4g, and (R)-3-methyl-1-picolinoyl-piperazine. MS  $m/z$ : (M+H)<sup>+</sup> Calc'd for C<sub>20</sub>H<sub>19</sub>BrN<sub>5</sub>O<sub>3</sub>: 456.07; Found 456.11. HPLC retention time: 1.12 minutes (column A).


### Intermediate 5n

15




Intermediate 5n, (S)-1-picolinoyl-3-methyl-4-[(7-bromo-6-azaindol-3-yl)-oxoacetyl]piperazine was prepared by the same method as Intermediate 5a starting from Potassium (7-bromo-6-azaindol-3-yl)oxoacetate, Intermediate 4g, and (S)-3-methyl-1-picolinoyl-piperazine. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 8.63 – 7.36 (m, 7H), 5.02 – 3.06 (m, 7H), 1.42 – 1.26 (m, 3H).

## Intermediate 5o



5 Intermediate 5o, (R)-1-picolinoyl-3-methyl-4-[(7-bromo-4-fluoro-6-azaindol-3-yl)-oxoacetyl]piperazine was prepared by the same method as Intermediate 5a starting from Potassium (7-bromo-4-fluoro-6-azaindol-3-yl)oxoacetate, Intermediate 4h, and (R)-3-methyl-1-picolinoyl-piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ8.68 – 7.52 (m, 6H), 4.94 – 2.69 (m, 7H), 1.48 – 1.24 (m, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>20</sub>H<sub>18</sub>BrFN<sub>5</sub>O<sub>3</sub>: 474.06; Found 474.23. HPLC retention time: 1.20 minutes (column A).

## Intermediate 5p




15

Intermediate 5p, 1-benzoyl-4-[(7-chloro-4-azaindol-3-yl)-oxoacetyl]piperazine was prepared by the same method as Intermediate 5a starting from Potassium (7-chloro-4-fluoro-4-azaindol-3-yl)oxoacetate, Intermediate 4e, and 1-benzoyl-  
 20 piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ8.83 (s, 1H), 8.63 (d, 1H, J = 5.35 Hz), 7.91 (d, 1H, J = 5.75 Hz), 7.47 (m, 5H), 3.80 – 3.30 (m, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>20</sub>H<sub>18</sub>ClN<sub>4</sub>O<sub>3</sub>: 397.11; Found 397.02. HPLC retention time: 1.20 minutes (column A).


25

## Intermediate 5q




5 Intermediate 5q, 1-(4-Benzoyl-piperazin-1-yl)-2-(7-bromo-4-chloro-1H-pyrrolo[2,3-c]pyridin-3-yl)-ethane-1,2-dione

To a solution of acid intermediate 4j (2.4 g, 7.9 mmol) in DMF (40 ml) was added 3-(diethoxyphosphoryloxy)-1,2,3-benzotriazin-4(3H)-one (DEPBT, 5.96 g, 19.9 mmol), benzoylpiperazine hydrochloride (2.71 g, 11.9 mmol), and *N,N*-diisopropylethylamine (14 ml, 80.4 mmol). The mixture was stirred at ambient temperature for 16 h. The reaction mixture was then added water (400 ml) and extracted with EtOAc (4 x 300 ml). The combined extracts were evaporated *in vacuo* to give a brownish residue, which was triturated with MeOH to provide the title compound as a white solid (2.8 g, 74%); <sup>1</sup>H NMR: (DMSO-*d*<sub>6</sub>) 13.41 (s, 1H), 8.48 (s, 1H), 8.19 (s, 1H); 7.45 (b s, 5H), 3.80 – 3.35 (b m, 8H); LC/MS: (ES+) *m/z* (M+H)<sup>+</sup> = 475, 477, 479; HPLC *R*<sub>t</sub> = 1.953.




Intermediate 5r was prepared by procedure used for 5q using mono N-Boc piperazine. <sup>1</sup>H NMR: (CDCl<sub>3</sub>) δ 8.26 (s, 1H), 8.19 (s, 1H), 3.71 (b s, 2H), 3.53 (b m, 6H), 1.48 (s, 9H); LC/MS: (ES+) *m/z* (M+H)<sup>+</sup> = 471, 473, 475; HPLC *R*<sub>t</sub> = 1.543.

**Intermediate 6**

Typical procedure for N-Oxide formation: Preparation of 1-benzoyl-3-(R)-  
 5 methyl-4-[(6-oxide-6-azaindol-3-yl)-oxoacetyl]piperazine, Intermediate 6. 20 mg of  
 1-benzoyl-3-(R)-methyl-4-[(6-azaindol-3-yl)-oxoacetyl]piperazine, Intermediate 5h,  
 (0.053 mmol) was dissolved in  $\text{CH}_2\text{Cl}_2$  (2 mL). 18 mg of mCPBA (0.11 mmol) was  
 then added into the solution and the reaction was stirred for 12 h at rt.  $\text{CH}_2\text{Cl}_2$  was  
 removed *via* evaporation at reduced pressure and the residue was purified using a  
 10 Shimadzu automated preparative HPLC System to give the compound shown above  
 (5.4 mg, 26%). MS  $m/z$ :  $(\text{M}+\text{H})^+$  Calc'd for  $\text{C}_{21}\text{H}_{21}\text{N}_4\text{O}_4$ : 393.16; Found 393.11.  
 HPLC retention time: 0.90 minutes (column A).

**Intermediate 7**

15




Preparation of 1-benzoyl-3-(R)-methyl-4-[(6-methyl-7-azaindol-3-yl)-  
 oxoacetyl]-piperazine or 1-benzoyl-3-(R)-methyl-4-[(4-methyl-7-azaindol-3-yl)-  
 oxoacetyl]-piperazine. An excess of  $\text{MeMgI}$  (3M in THF, 0.21 ml, 0.63 mmol) was  
 20 added into a solution of 1-benzoyl-3-(R)-methyl-4-[(6-oxide-6-azaindol-3-yl)-  
 oxoacetyl]piperazine, Intermediate 6, (25 mg, 0.064 mmol). The reaction mixture  
 was stirred at rt and then quenched with MeOH. The solvents were removed under  
 vacuum, the residue was diluted with MeOH and purified using a Shimadzu

automated preparative HPLC System to give a compound shown above which was a single isomer but regiochemistry was not definitively assigned. (6.7 mg, 27%). MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{22}H_{23}N_4O_3$ : 391.18; Found 391.17. HPLC retention time: 1.35 minutes (column B).


5

### Intermediate 8



- 10 1-benzoyl-3-(R)-methyl-4-[(6-phenyl-7-azaindol-3-yl)-oxoacetyl]piperazine or 1-benzoyl-3-(R)-methyl-4-[(4-phenyl-7-azaindol-3-yl)-oxoacetyl]piperazine (regiochemistry was not definitively assigned) were prepared by the method described for Example 7 starting with 1-benzoyl-3-(R)-methyl-4-[(6-oxide-6-azaindol-3-yl)-oxoacetyl]piperazine, Intermediate 6, and phenyl magnesium bromide
- 15 (phenyl Grignard reagent). MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{27}H_{25}N_4O_3$ : 453.19; Found 454.20. HPLC retention time: 1.46 minutes (column B).

### Intermediate 9




20

A mixture of Pd (10% on carbon, 100 mg), trifluoroacetic acid (1 mL) and 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-methyl-4-azaindol-3-yl)-oxoacetyl]piperazine, Intermediate 5e (1.5 g) in MeOH (50 mL) and EtOAc (50 mL) was shaken in a Parr reactor under a hydrogen atmosphere (45 psi) for 48 hours. After solids were



removed via filtration, the filtrate was concentrated *in vacuo* to afford intermediate 9 (1 g) which was used without further purification. MS *m/z*: (M+H)<sup>+</sup> calcd for C<sub>21</sub>H<sub>21</sub>N<sub>4</sub>O<sub>3</sub> 391.18, found 391.15. HPLC retention time: 1.15 minutes (column A).

5

### Intermediates 10 and 11



- 10 Preparation of Intermediate 10, 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-carbonyl-4-azaindol-3-yl)-oxoacetyl]-piperazine and Intermediate 11, 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-hydroxycarbonyl-4-azaindol-3-yl)-oxoacetyl]-piperazine: A mixture of 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-methyl-4-azaindol-3-yl)-oxoacetyl]piperazine (1.78 g) and SeO<sub>2</sub> (4.7 g) in dioxane/water (100 : 1) was
- 15 refluxed for 10 hours. After cooling to room temperature, the mixture was concentrated *in vacuo* to provide a residue. The residue was purified by using silica gel chromatography with EtOAc and MeOH as eluting solvents to afford intermediate 10 (350 mg) and intermediate 11 (410 mg).
- Intermediate 10, 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-carbonyl-4-azaindol-3-yl)-oxoacetyl]-piperazine: MS *m/z*: (M+H)<sup>+</sup> calcd for C<sub>22</sub>H<sub>20</sub>ClN<sub>4</sub>O<sub>4</sub>: 439.12, found 439.01. HPLC retention time: 1.37 minutes (column A); Intermediate 11, 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-hydroxycarbonyl-4-azaindol-3-yl)-oxoacetyl]-piperazine: MS *m/z*: (M+H)<sup>+</sup> calcd for C<sub>22</sub>H<sub>20</sub>ClN<sub>4</sub>O<sub>5</sub>: 455.11, found 455.10. HPLC retention time: 1.44 minutes (column A).
- 20

**Intermediates 12 and 13**

5

Intermediate 12, 1-benzoyl-3-(R)-methyl-4-[(7-carbonyl-4-azaindol-3-yl)-oxoacetyl]-piperazine and Intermediate 13, 1-benzoyl-3-(R)-methyl-4-[(7-hydroxycarbonyl-4-azaindol-3-yl)-oxoacetyl]-piperazine were made according to the same procedure of preparing Intermediates 10 and 11, by using Intermediate 9 as a starting material. Intermediate 12, 1-benzoyl-3-(R)-methyl-4-[(7-carbonyl-4-azaindol-3-yl)-oxoacetyl]-piperazine: MS  $m/z$ :  $(M+H)^+$  calcd for  $C_{22}H_{21}N_4O_4$ : 405.16, found 405.14. HPLC retention time: 0.91 minutes (column A); Intermediate 13, 1-benzoyl-3-(R)-methyl-4-[(7-hydroxycarbonyl-4-azaindol-3-yl)-oxoacetyl]-piperazine: MS  $m/z$ :  $(M+H)^+$  calcd for  $C_{22}H_{21}N_4O_5$ : 421.15, found 421.09. HPLC retention time: 1.02 minutes (column A).

15












**Intermediates 14a-1 - 14a-21**











20

The following tin agents and boron agents can be purchased from commercial resources and used without any further treatment (Table 2).



**Table 2**

Intermediate Number	Structure	Company
14a-1		Frontier Scientific, Inc.
14a-2		Maybridge Chem. Co.
14a-3		Frontier Scientific, Inc.
14a-4		Matrix Scientific
14a-5		Matrix Scientific
14a-6		Aldrich, Co.
14a-7		Aldrich, Co.
14a-8		Aldrich, Co.
14a-9		Aldrich, Co.
14a-10		Aldrich, Co.
14a-11		Lancaster









14a-12		Aldrich, Co.
14a-13		Aldrich, Co.
14a-14		Frontier Scientific, Inc.
14a-15		Matrix Scientific
14a-16		Frontier Scientific, Inc.
14a-17		Riedel-de Haen AG
14a-18		Lancaster
14a-19		Lancaster
14a-20		Aldrich, Co.
14a-21		Frontier Scientific, Inc.












Preparation of Tin Agents:Intermediates 14-1 – 14-65









- 5           The following known tin agents and boron agents could be prepared according to the documented procedures indicated without any modification (Table 3):









Table 3










10







Intermediate Number	Structure	Reference
14-1		Dondoni, A., et al Synthesis, 1987, 693
14-2		Aldous, D. J., et al US – 5,453,433
14-3		Sandosham, J., et al Tetrahedron 1994, 50, 275.
14-4		Lehn, L. M., et al. Chem. Eur. J. 2000, 6, 4133.
14-5		Jutzi, P., et al J. Organometallic Chem. 1983, 246, 163.
14-6		Jutzi, P., et al J. Organometallic Chem. 1983, 246, 163.
14-7		Graybill, T. L., et al Bioorg. Med. Chem. Lett. 1995, 5 (4), 387.
14-8		Heldmann, D. K., et al Tetrahedron Lett. 1997, 38, 5791.

14-9	 <chem>CCCC[Sn](CCCC)(CCCC)c1ccncn1</chem>	Kennedy, G., et al Tetrahedron Lett. 1996, 37, 7611.
14-10	 <chem>CCCC[Sn](CCCC)(CCCC)c1cc(C)nco1</chem>	Kondo, Y., et al Tetrahedron Lett. 1989,30, 4249
14-11	 <chem>CCCC[Sn](CCCC)(CCCC)c1cc(C)c(C(=O)OCC)nco1</chem>	Kondo, Y., et al Tetrahedron Lett. 1989,30, 4249
14-12	 <chem>CCCC[Sn](CCCC)(CCCC)c1ccsc1</chem>	Or, Y. S., et al US-6,054,435
14-13	 <chem>CCCC[Sn](CCCC)(CCCC)c1ccsc1</chem>	Or, Y. S., et al US-6,054,435
14-14	 <chem>CCCC[Sn](CCCC)(CCCC)c1ccnnc1</chem>	Okada, T., et al WO-0123383
14-15	 <chem>CCCC[Sn](CCCC)(CCCC)c1ccnnc1</chem>	Okada, T., et al WO-0123383
14-16	 <chem>CCCC[Sn](CCCC)(CCCC)c1cc(C)nnc1</chem>	Sandosham, J., et al Tetrahedron 1994, 50, 275
14-17	 <chem>CCCC[Sn](CCCC)(CCCC)c1cc(C)nnc1</chem>	Sandosham, J., et al Acta Chem. Scand. 1989, 43, 684.
14-18	 <chem>CCCC[Sn](CCCC)(CCCC)c1cc(OC)nsc1</chem>	Nicolaou, K. C., et al WO-9967252
14-19	 <chem>CCCC[Sn](CCCC)(CCCC)c1cc(OCC)nsc1</chem>	Nicolaou, K. C., et al WO-9967252








14-20		Nicolaou, K. C., et al WO-9967252
14-21		Benheda, R., et al Tetrahedron Lett. 1999, 40, 5701.
14-22		Collins, I., et al Tetrahedron Lett. 1999, 40, 4069.
14-23		Fuss, R. W., et al DE-19502178
14-24		Bunnage, M.E.et.al PCT Int. Appl. WO 0024745 A1 (2000); and Sandosham, J. et. Al Tetrahedron (1994), 50(1), 275-84.
14-25		From 5-iodo2-chloro-1,3 pyrimidine. Fluoropyrimidines are obtained by fluorination of chloropyrimidines with CsF in N-methyl-2- pyrrolidinone or DMF 2.5-63 h at 80-150 °C. The iodo is then converted to the lithium reagent with tBuLi and trapped with Bu <sub>3</sub> SnCl. See Sandosham above.
14-26		Arukwe, J.; Benneche, T.; Undheim, K. J. Chem. Soc., Perkin Trans. 1 (1989), (2), 255-9.
14-27		Fruit, C.; et.al. Heterocycles (1999), 51(10), 2349-2365.








14-28		Ziener, U.; et.al. Chem.— Eur. J. (2000), 6(22), 4132-4139.
14-29		Turck, A.; et.al Lab.. J. Organomet. Chem. (1991), 412(3), 301-10. Metallation of 2,6- dicloropyrazine and quench with Bu <sub>3</sub> SnCl.
14-30		Ueno, K.; Sasaki, A.; Kawano, K.; Okabe, T.; Kitazawa, N.; Takahashi, K.; Yamamoto, N.; Suzuki, Y.; Matsunaga, M.; Kubota, A. PCT Int. Appl. WO 9918077 A1 (1999).
14-31		Fensome, A.; Miller, L. L.; Ullrich, J.W.; Bender, R.H.W.; Zhang, P.; Wrobel, J.E.; Zhi, L.; Jones, T.K.; Marschke, K.B.; Tegley, C.M. PCT Int. Appl. WO 0066556 A1 (2000).
14-32		Maw, G.N.; Middleton, D.S. Jpn. Kokai Tokkyo Koho JP 2000016984 A2 (2000).
14-33		Chem. Pharm. Bull. (1998), 46(3), 400-412.
14-34		Hayashi, K.; Kito, T.; Mitsuyama, J.; Yamakawa, T.; Kuroda, H.; Kawafuchi, H. PCT Int. Appl. WO 9951588 A1 (1999).
14-35		Brown, A.D.; Dickinson, R.P.; Wythes, M.J. PCT Int. Appl. WO 9321178 A1 (1993).


14-36		Brown, A.D.; Dickinson, R.P.; Wythes, M.J. PCT Int. Appl. WO 9321178 A1 (1993).
14-37		Zalutsky, M.R. PCT Int. Appl. WO 0032240 A2 (2000).
14-38		Brown, A.D.; Dickinson, R.P.; Wythes, M. J. PCT Int. Appl. WO 9321178 A1 (1993).
14-39		North, P.C.; Wadman, S.N. PCT Int. Appl. WO 9408993 A1 (1994).
14-40		North, P.C.; Wadman, S.N. PCT Int. Appl. WO 9408993 A1 (1994).
14-41		Achab, S.; Guyot, M.; Potier, P. Tetrahedron Lett. (1993), 34(13), 2127-30.
14-42		Muratake, H.; Tonegawa, M.; Natsume, M... Chem. Pharm. Bull. (1998), 46(3), 400-412. Dehmlow, E.V.; Slegers, A. Liebigs Ann. Chem. (1992), (9), 953-9.
14-43		Proudfoot, J.R.; Hargrave, K.; Kapadia, S. PCT Int. Appl. WO 9907379 A1 (1999); and Chem. Pharm. Bull. (1998), 46(3), 400-412.
14-44		Cruskie, M.P. Jr.; Zoltewicz, J.A.; Abboud, K.A. J. Org. Chem. (1995), 60(23), 7491-5.


14-45		Muratake, H.; et.al Chem. Pharm. Bull. (1998), 46(3), 400-412.
14-46		Muratake, H.; Tonegawa, M.; Natsume, M. Chem. Pharm. Bull. (1998), 46(3), 400-412. Dolle, R.E.; Graybill, T.L.; Osifo, I.K.; Harris, A.L.; Miller, M.S.; Gregory, J.S. U.S. US 5622967 (1997).
14-47		Henze, O.; Lehmann, U.; Schlueter, A.D. Synthesis (1999), (4), 683-687.
14-48		Hayashi, K.; Kito, T.; Mitsuyama, J.; Yamakawa, T.; Kuroda, H.; Kawafuchi, H. PCT Int. Appl. WO 9951588 A1 (1999); Reuman, M.; Daum, S.J.; Singh, B.; Wentland, M.P.; Perni, R.B.; Pennock, P.; Carabateas, P.M.; Gruett, M.D.; Saindane, M.T.; et al. J. Med. Chem. (1995), 38(14), 2531-40.
14-49		Barros, M.T.; Maycock, C.D.; Ventura, M.R. Tetrahedron Lett. (1999), 40(3), 557-560. Sirisoma, N.S.; Johnson, C.R. Tetrahedron Lett. (1998), 39(15), 2059- 2062. Trost, B.M.; Cook, G.R. Tetrahedron Lett. (1996), 37(42), 7485- 7488.
14-50		Bunnage, M.E.; Maw, G.N.; Rawson, D.J.; Wood, A.; Mathias, J.P.; Street, S.D.A. PCT Int. Appl. WO 0024745 A1 (2000).




14-51		Bunnage, M.E.; Maw, G.N.; Rawson, D.J.; Wood, A.; Mathias, J. P.; Street, S.D.A. PCT Int. Appl. WO 0024745 A1 (2000).
14-52		Hayashi, K.; Kito, T.; Mitsuyama, J.; Yamakawa, T.; Kuroda, H.; Kawafuchi, H. PCT Int. Appl. WO 9951588 A1 (1999); and Sirisoma, N.S.; Johnson, C.R. Tetrahedron Lett. (1998), 39(15), 2059-2062.
14-53		Schnatterer, S.; Kern, M.; Sanft, U. PCT Int. Appl. WO 9965901 A1 (1999).
14-54		Hayashi, K.; Kito, T.; Mitsuyama, J.; Yamakawa, T.; Kuroda, H.; Kawafuchi, H. PCT Int. Appl. WO 9951588 A1 (1999).
14-55		Betageri, R.; Breitfelder, S.; Cirillo, P.F.; Gilmore, T.A.; Hickey, E.R.; Kirrane, T.M.; Moriak, M.H.; Moss, N.; Patel, U.R.; Proudfoot, J.R.; Regan, J.R.; Sharma, R.; Sun, S.; Swinamer, A.D.; Takahashi, H. PCT Int. Appl. WO 0055139 A2 (2000).
14-56		Ueno, K.; Sasaki, A.; Kawano, K.; Okabe, T.; Kitazawa, N.; Takahashi, K.; Yamamoto, N.; Suzuki, Y.; Matsunaga, M.; Kubota, A. PCT Int. Appl. WO 9918077 A1 (1999).
14-57		Calderwood, D.; Arnold, L.D.; Mazdiyasni, H.; Hirst, G.; Deng, B.B. PCT Int. Appl. WO 0017202 A1 (2000).

14-58		Hayashi, K.; Kito, T.; Mitsuyama, J.; Yamakawa, T.; Kuroda, H.; Kawafuchi, H. PCT Int. Appl. WO 9951588 A1 (1999).
14-59		Saji, H.; Watanabe, A.; Magata, Y.; Ohmono, Y.; Kiyono, Y.; Yamada, Y.; Iida, Y.; Yonekura, H.; Konishi, J.; Yokoyama, A. Chem. Pharm. Bull. (1997), 45(2), 284-290.
14-60		Hayashi, K.; Kito, T.; Mitsuyama, J.; Yamakawa, T.; Kuroda, H.; Kawafuchi, H. PCT Int. Appl. WO 9951588 A1 (1999); Reuman, M.; Daum, S.J.; Singh, B.; Wentland, M.P.; Perni, R.B.; Pennock, P.; Carabateas, P.M.; Gruett, M.D.; Saindane, M.T.; et al. J. Med. Chem. (1995), 38(14), 2531-40.
14-61		Iino, Y.; Fujita, K.; Kodaira, A.; Hatanaka, T.; Takehana, K.; Kobayashi, T.; Konishi, A.; Yamamoto, T. PCT Int. Appl. WO 0102359 A1 (2001).
14-62		Iino, Y.; Fujita, K.; Kodaira, A.; Hatanaka, T.; Takehana, K.; Kobayashi, T.; Konishi, A.; Yamamoto, T. PCT Int. Appl. WO 0102359 A1 (2001).
14-63		Torrado, A.; Imperiali, B. J. Org. Chem. (1996), 61(25), 8940-8948.
14-64		Iino, Y.; Fujita, K.; Kodaira, A.; Hatanaka, T.; Takehana, K.; Kobayashi, T.; Konishi, A.; Yamamoto, T. PCT Int. Appl. WO 0102359 A1 (2001).

14-65		Gros, P.; Fort, Y. <i>Synthesis</i> (1999), (5), 754-756 and Gros, P.; Fort, Y.; Caubere, P. J. <i>Chem. Soc., Perkin Trans. 1</i> (1997), (20), 3071-3080.
-------	---	---

Intermediate 14-66

- 5 Preparation of 2,3-dichloro-5-(tri-n-butylstannyl)pyrazine (An example of general procedure Tin-01, below): TMP-Li (2,2,6,6-tetramethylpiperidinyllithium) was prepared by addition of n-butyl lithium (1.6 M, 6.25 mL) to a solution of 2,2,4,4-tetramethylpiperidine (1.4 g) in dry THF (180 mL) at  $-78^{\circ}\text{C}$ . The solution was then allowed to warm to  $0^{\circ}\text{C}$ , was stirred at  $0^{\circ}\text{C}$  for 15 minutes, then was
- 10 cooled to  $-78^{\circ}\text{C}$ . To the solution was added 2,3-dichloropyrazine (1.35 g), and followed by an addition of tri-n-butyltin chloride (3.25 g) in another 2 hours. The reaction was quenched with aqueous ammonium chloride solution. The organic layer was separated, and aqueous layer was extracted with ethyl acetate (3 x 100 mL). The combined organic extract was dried over magnesium sulfate, filtered and the filtrate
- 15 concentrated *in vacuo*. The residue was purified by silica gel chromatography to afford 2,3-dichloro-5-(tri-n-butylstannyl)pyrazine (1 g).


Intermediate 14-67

20

Preparation of 2-(tri-n-butylstannyl)-pyrimidine: (Example of the general procedure Tin-03, below) Tri-n-butylstannyl lithium was prepared at  $0^{\circ}\text{C}$  in dry THF (20 mL) from tri-butyltin hydride (2.2 mL) and LDA (lithium diisopropylamide, 2M,


4.09 mL). The tri-*n*-butylstannyl lithium solution was then cooled to  $-78^{\circ}\text{C}$  and to it was added 2-bromopyrimidine (1 g). The reaction mixture was then allowed to warm up to room temperature over 8 hours. The reaction was then quenched with aqueous ammonium chloride solution. The organic layer was separated, and aqueous layer was extracted with ethyl acetate (3 x 20 mL). The combined organic layer was dried over magnesium sulfate, filtered and the filtrate concentrated *in vacuo*. The residue was purified by silica gel chromatography to afford 2-(tri-*n*-butylstannyl)-pyrimidine (190 mg).

10

Intermediate 14-68

Preparation of 2-amino-6-(tri-*n*-butylstannyl)pyrazine (Example of the general procedure Tin-04, below): To a sealed tube, 2-amino-6-chloro-pyrazine (1 g), bis(tri-*n*-butyltin) (3.92 mL) and tetrakis-triphenylphosphine palladium, Pd(Ph<sub>3</sub>P)<sub>4</sub> (100 mg) were combined in dioxane (10 mL). The reaction was heated at 110-120  $^{\circ}\text{C}$  for 10 h. After the mixture cooled down to room temperature, it was poured into 20 mL of water. The solution was extracted with EtOAc (4 x 20 mL). The combined extract was concentrated *in vacuo* to give a residue which was purified by silica gel chromatography to afford 2-amino-6-(tri-*n*-butylstannyl)pyrazine (0.5 g).

20

Intermediate 14-69

25

Preparation of 2-methylsulfonylamino-5-(tri-*n*-butylstannyl)pyrazine (Example of general procedure Tin-05, below): NaH (60%, 20 mg) was added into a solution of 2-amino-5-(tri-*n*-butylstannyl)pyrazine (0.2 g) in THF (30 mL) at room


temperature. After the mixture stirred at room temperature for 30 minutes, to it was added methylsulfonyl chloride (63 mg). The reaction mixture was stirred at room temperature over 8 hours. The reaction was quenched with aqueous ammonium chloride solution. The organic layer was separated, and the aqueous layer was  
 5 extracted with ethyl acetate (3 x 100 mL). The combined organic extract was dried over magnesium sulfate, filtered and the filtrate was concentrated *in vacuo*. The residue was purified by silica gel chromatography to afford 2-methylsulfonylamino-5-(tri-n-butylstannyl)pyrazine (20 mg).

10

### Intermediates 14-70 – 14-129

The intermediates 14-70 – 14-129 were prepared according to the following general procedures designated Tin-01 through Tin-05.


#### 15 General Procedure Tin-01:



Base = LDA, TMP-Li, n-BuLi, S-BuLi or t-BuLi;  
 Solvent = THF, diethyl ether or DME;  
 R = methyl or butyl


To a solution of a base (1.1 equivalents) selected from lithium diisopropylamide, 2,2,6,6-tetramethylpiperidinyllithium, n-butyl lithium, sec-butyl lithium or tert-butyl lithium in a solvent selected from tetrahydrofuran, diethyl ether or dimethoxyethane (concentration of approximately 0.05 mmol base/mL of solvent) at -78 °C was added an appropriate aryl or heteroaryl substrate (1.0 equivalents) followed by an addition of tri-n-butyltin chloride or trimethyltin chloride (1.1  
 20 equivalents) in another 2 hours. The reaction was quenched with aqueous ammonium chloride solution. The organic layer was separated, and aqueous layer was extracted with ethyl acetate. The combined organic extract was dried over magnesium sulfate, filtered and the filtrate concentrated *in vacuo*. The residue was purified by silica gel chromatography to afford the desired stannane.  
 25

30

General Procedure Tin-02:

LG = Br or I; Base = n-BuLi, S-BuLi, or t-BuLi;  
 Solvent = THF, ether or DME;  
 R = methyl or butyl

- 5 To a solution of a base (1.1 equivalents) selected from n-butyl lithium, sec-butyl lithium or tert-butyl lithium in a solvent selected from tetrahydrofuran, diethyl ether or dimethoxyethane (concentration of approximately 0.05 mmol base/mL of solvent) at  $-78^\circ\text{C}$  was added an appropriate aryl or heteroaryl bromide or aryl or heteroaryl iodide substrate (1.0 equivalents). The reaction mixture was stirred at  $-78^\circ\text{C}$  for a period suitable to generate the anion via metal-halogen exchange then to it was added tri-n-butylin chloride or trimethyltin chloride (1.1 equivalents). The reaction was quenched with aqueous ammonium chloride solution. The organic layer was separated, and aqueous layer was extracted with ethyl acetate. The combined organic extract was dried over magnesium sulfate, filtered and the filtrate concentrated *in vacuo*. The residue was purified by silica gel chromatography to afford the desired stannane.


General Procedure Tin-03:

- 20 LG = F, Cl, Br, I; Solvent = THF, diethyl ether or DME; R = methyl or butyl

- 25 Tri-n-butylin lithium or trimethylstannyl lithium (1.3 equivalents) was prepared at  $0^\circ\text{C}$  in dry solvent selected from THF, diethyl ether or dimethoxyethane (20 mL) from tri-n-butylin hydride or trimethyltin hydride, respectively (1.3 equivalents) and LDA (lithium diisopropylamide, 1.3 equivalents) at a concentration of approximately 0.4 mmol of alkylstannyl lithium/mL of solvent. The tri-n-butylin lithium or trimethylstannyl lithium solution was then cooled to  $-78^\circ\text{C}$  and to it was added an appropriate haloaryl or haloheteroaryl substrate (1.0 equivalent). The reaction mixture was then allowed to warm up to room temperature

over 8 hours. The reaction was then quenched with aqueous ammonium chloride solution. The organic layer was separated, and aqueous layer was extracted with ethyl acetate (3 x 20 mL). The combined organic layer was dried over magnesium sulfate, filtered and the filtrate concentrated *in vacuo*. The residue was purified by silica gel chromatography to afford the desired stannane intermediate.

General Procedure Tin-04:



LG = Cl, Br, I or OTf; Solvent = Dioxane or Toluene; R = methyl or butyl

10

To a sealed tube, an appropriate aryl or heteroaryl substrate (1.0 equivalent), bis(tri-butyltin) or hexamethylditin (1.0 equivalent) and tetrakis-triphenylphosphine palladium, Pd(Ph<sub>3</sub>P)<sub>4</sub> (1.0 mol%) were combined in dioxane or toluene (10 mL). The reaction was heated at 110-120 °C for 10 h. After the mixture cooled down to room temperature, it was poured into water. The solution was extracted with ethyl acetate and the combined extracts were concentrated *in vacuo* to give a residue which was purified by silica gel chromatography to afford the desired stannane product.


15

General Procedure Tin-05:

20

The following general reaction scheme depicts the derivatization of stannane intermediates in which the stannane has a reactive ring NH group or reactive exocyclic amino, hydroxy or thiol group. The starting stannane is treated with base in an appropriate solvent then is reacted with suitable electrophiles such as alkyl halides, acid chlorides, sulfonyl chlorides, isocyanates and the like.

25



**Electrophile = R'-halide, R'C(O)Cl, R'OC(O)Cl, R'R''NCOCl, R'SO<sub>2</sub>Cl, R'NCO, R'NSO, R'NCNR''**

**E = R'-, R'C(O)-, R'OC(O)-, R'R''NC(O)-, R'SO<sub>2</sub>-, R'NC(O)-, R'NS(O)-, R'NCNR''**

**Solvent = CH<sub>2</sub>Cl<sub>2</sub>, THF, diethyl ether, DMF**


**R = methyl or butyl; X = NH, O or S**

**Base = NaH, BuLi, LDA, K<sub>2</sub>CO<sub>3</sub>, Et<sub>3</sub>N, DBU, DMAP, NaHMDS**

- An appropriate base selected from sodium hydride, n-butyl lithium, lithium diisopropylamide, potassium carbonate, triethylamine, DBU, DMAP or sodium
- 5 hexamethyldisilazide (1.0 equivalent) was added into a solution of an appropriate stannane substrate (as depicted above, 1.0 equivalent) in an appropriate solvent selected from dichloromethane, THF, diethyl ether or N,N-dimethylformamide at a temperature between -78 °C and room temperature. After the mixture stirred for a period sufficient to allow deprotonation, typically for 5 to 30 minutes, then to it was
- 10 added an appropriate electrophile such as an alkyl halide, acid chloride, sulfonyl (1.0 equivalent). The reaction mixture was stirred, typically at room temperature, over a period of 2 to 8 hours. The reaction was quenched with aqueous ammonium chloride solution. The organic layer was separated, and the aqueous layer was extracted with ethyl acetate (3 x 100 mL). The combined organic extract was dried over
- 15 magnesium sulfate, filtered and the filtrate was concentrated in vacuo. The residue was purified by silica gel chromatography to afford the desired stannane intermediate.



## General procedure Tin-06

































- 5 An aryl hilide stannane agent was dissolved in appropriate alcohol, either methanol or ethanol. After a cataylst (pt or pd) was added into the solvent, the reaction mixture is placed in an environment of hydrogen under normal or raised pressure. After reaction finishes, the catalyst is filtered, and, concentration of the mother solution provides a residue which is used in the further reactions without any
- 10 purification.

















R<sub>f</sub> = retention time















Intermed. Number	Structure	Starting Material	Method Applied	Identification
14-70			Tin-04	R <sub>f</sub> = 2.33 min (Column A) <sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 4.00 (s, 6H), 1.63 – 0.85 (m, 27H)
14-71			Tin-01	R <sub>f</sub> = 2.52 min (Column A) <sup>1</sup> H NMR (300 MHz, CDCl <sub>3</sub> ) δ 7.02 (s, 1H), 4.44 (q, 2H, J = 7.02 Hz), 1.63 – 0.85 (m, 30H)
14-72			Tin-01	R <sub>f</sub> = 2.84 min (Column B) <sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 9.48 (s, 1H), 8.45 (s, 1H), 2.03 – 0.88 (m, 36H)
14-73			Tin-05	R <sub>f</sub> = 2.27 min (Column A) <sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 7.53 (m, 1H), 6.29 (m, 1H), 3.94 (s, 3H), 1.56 – 0.87 (m, 27H)













14-74			Tin-05	$R_f = 2.22$ min (Column A)
14-75			Tin-01	$R_f = 2.44$ min (Column B) $^1\text{H NMR}$ (500 MHz, $\text{CDCl}_3$ ) $\delta$ 8.89 (s, 1H), 8.34 (s, 1H), 1.61 – 0.85 (m, 27H)
14-76			Tin-01	$R_f = 3.41$ min (Column A, flow rate = 4 ml/min) $^1\text{H NMR}$ (300 MHz, $\text{CDCl}_3$ ) $\delta$ 8.58 (d, 1H, $J = 2.52$ Hz), 8.13 (d, 1H, $J = 2.52$ Hz), 1.63 – 0.85 (m, 27H)
14-77			Tin-01	$R_f = 3.89$ min (Column A, flow rate = 4 ml/min) $^1\text{H NMR}$ (300 MHz, $\text{CDCl}_3$ ) $\delta$ 8.63 (s, 1H), 1.61 – 0.85 (m, 27H)
14-78			Tin-01	$R_f = 3.86$ min (Column A, flow rate = 4 ml/min) $^1\text{H NMR}$ (300 MHz, $\text{CDCl}_3$ ) $\delta$ 8.24 (s, 1H), 1.61 – 0.85 (m, 27H)
14-79			Tin-04	$R_f = 2.10$ min (Column B) $^1\text{H NMR}$ (500 MHz, $\text{CDCl}_3$ ) $\delta$ 7.90 (s, 1H), 7.26 (s, 1H), 1.58 – 0.87 (m, 27H)
14-80			Tin-04	$R_f = 1.83$ min (Column A)















14-81			Tin-04	$R_f = 1.84$ min (Column A)
14-82			Tin-04	$R_f = 1.84$ min (Column A)
14-83			Tin-04	$R_f = 1.90$ min (Column A)
14-84			Tin-01	$R_f = 2.23$ min (Column A)
14-85			Tin-04	$R_f = 1.92$ min (Column A)
14-86			Tin-03	$R_f = 2.01$ min (Column A)
14-87			Tin-01	$R_f = 2.45$ min (Column A)
14-88			Tin-01	$R_f = 2.67$ min (Column C)
14-89			Tin-01	$R_f = 2.31$ min (Column C)

14-90	 <chem>CCCC[Sn](CCCC)(CCCC)c1ccncc1</chem>	 <chem>Brc1ccncc1</chem>	Tin-04	$R_f = 2.71$ min (Column D)
14-91	 <chem>CCCC[Sn](CCCC)(CCCC)c1cc(s1)NC(=O)OC(C)(C)C</chem>	 <chem>CC1(C)OC(=O)NC2=CN=CSC2</chem>	Tin-01	$R_f = 2.49$ min (Column C)
14-92	 <chem>CCCC[Sn](CCCC)(CCCC)c1cc(s1)n2c(Cl)ncn2</chem>	 <chem>Clc1cc2nncn2s1</chem>	Tin-01	$R_f = 2.42$ min (Column C)
14-93	 <chem>CCCC[Sn](CCCC)(CCCC)c1cc(s1)n2c3nncn3n2</chem>	 <chem>CN1C=NC2=NNC2=N1</chem>	Tin-01	$R_f = 3.49$ min (Column C) Flow Rate = 4 ml/min
14-94	 <chem>CCCC[Sn](CCCC)(CCCC)c1cc(s1)n2c(=O)ccn2</chem>	 <chem>O=C1CCCN1c2cc(s2)</chem>	Tin-01	$R_f = 2.46$ min (Column C)
14-95	 <chem>CCCC[Sn](CCCC)(CCCC)c1ccncc1C</chem>	 <chem>CCCC[Sn](CCCC)(CCCC)c1ccc(N)cc1</chem>	Tin-05	$R_f = 2.15$ min (Column A)













14-96			Tin-01	$R_f = 2.28$ min (Column C)
14-97			Tin-01	$R_f = 2.60$ min (Column C)
14-98			Tin-01	$R_f = 2.37$ min (Column A)
14-99			Tin-01	$R_f = 2.59$ min (Column A)
14-100			Tin-01	$R_f = 2.49$ min (Column C)
14-101			Tin-04	$R_f = 2.41$ min (Column A)
14-102			Tin-04	$R_f = 1.88$ min (Column E)
14-103			Tin-04	$R_f = 1.92$ min (Column E)

14-104			Tin-04	R <sub>f</sub> = 2.01 min (Column E)
14-105			Tin-04	R <sub>f</sub> = 2.15 min (Column E)
14-106			Tin-04	R <sub>f</sub> = 1.91 min (Column E)
14-107			Tin-04	R <sub>f</sub> = 1.95 min (Column A)
14-108			Tin-04	R <sub>f</sub> = 1.93 min (Column A)
12-109			Tin-01	R <sub>f</sub> = 1.95 min (Column A)
14-110			Tin-01	R <sub>f</sub> = 1.83 min (Column A) <sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 9.03 (d, 1H, J = 5.15 Hz), 7.49 (d, 1H, J = 7.95 Hz), 7.26 (m, 1H), 1.61 – 0.86 (m, 27H); <sup>13</sup> C NMR (125 MHz, CDCl <sub>3</sub> )

				$\delta$ 175.3, 149.8, 133.2, 123.7, 29.0, 27.3, 13.6, 10.1.
14-111			Tin-01	$R_f$ = 2.18 min (Column E) $^1\text{H}$ NMR (500 MHz, $\text{CDCl}_3$ ) $\delta$ 9.22 (s, 1H), 8.46 (d, 1H, $J$ = 4.80 Hz), 7.42 (d, 1H, $J$ = 4.75 Hz), 1.56 – 0.86 (m, 27H); $^{13}\text{C}$ NMR (125 MHz, $\text{CDCl}_3$ ) $\delta$ 185.4, 158.0, 153.2, 130.6, 28.9, 27.2, 13.5, 9.9.
14-112			Tin-04	$R_f$ = 1.96 min (Column A)
14-113			Tin-01	$R_f$ = 2.61 min (Column A)
14-114			Tin-01	$R_f$ = 2.85 min (Column A)
14-115			Tin-05	$R_f$ = 2.09 min (Column A) $^1\text{H}$ NMR (500 MHz, $\text{CDCl}_3$ ) $\delta$ 8.12 (s, 1H), 7.95 (s, 1H), 4.11 (s, 1H), 2.95 (s, 3H), 2.03 – 0.85 (m, 27H)
14-116			Tin-05	$R_f$ = 2.16 min (Column A) $^1\text{H}$ NMR (500 MHz, $\text{CDCl}_3$ ) $\delta$ 8.08 (s, 1H), 7.92 (s, 1H), 4.49 (s, 1H), 3.35 (m, 2H),








				1.63 – 0.85 (m, 30H)
14-117			Tin-04	Rf = 2.19 min (Column A)
14-118			Tin-04	Rf = 2.18 min (Column A)
14-119			Tin-04	Rf = 2.47 min (Column A) <sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 7.85 (s, 1H), 4.91 (s, 2H), 2.16 – 0.87 (m, 27H)
14-120			Tin-04	Rf = 2.61 min (Column A)
14-121			Tin-04	Rf = 2.92 min (Column A)
14-122			Tin-04	Rf = 1.93 min (Column A)
14-123			Tin-01	Rf = 2.20 min (Column A)




















14-124			Tin-01	R <sub>f</sub> = 2.50 min (Column A) <sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 9.07 (s, 1H), 7.87 (s, 1H), 1.59 – 0.85 (m, 27H)
12-125			Tin-04	R <sub>f</sub> = 1.97 min (Column A)
14-126			Tin-04	R <sub>f</sub> = 1.97 min (Column A)
14-127			Tin-01	R <sub>f</sub> = 2.70 min (Column E) <sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 8.11 (d, 1H, J = 5.2 Hz), 7.41 (d, 1H, J = 5.2 Hz), 6.94 (s, 1H), 1.62 – 0.89 (m, 27H)
14-128			Tin-06	<sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 8.12 (d, 1H, J = 5.2 Hz), 7.78 (s, 1H), 7.46 (d, 1H, J = 5.2 Hz), 6.84 (s, 1H), 1.98 – 0.85 (m, 27H)
14-129			Tin-01	R <sub>f</sub> = 1.86 min (Column A)









The following table contains novel stannane reagents which can be prepared by the methodology described above and then could be used to prepare compounds of formula I










**Table 3**










Intermediate Number	Structure	Reference
		<p>From 5-iodo-2-chloro-1,3 pyrimidine. Fluoropyrimidines are obtained by fluorination of chloropyrimidines with CsF in N-methyl-2-pyrrolidinone or DMF 2.5-63 h at 80-150°. The iodo is then converted to the lithium reagent with tBuLi and trapped with Bu<sub>3</sub>SnCl. See Sandosham above.</p>
		
		
		
		
		
		









	 <chem>Fc1nc(Sn(CCCC)CCC)nc1</chem>	
	 <chem>Fc1nc(Sn(CCCC)CCC)nc1</chem>	
	 <chem>COC(=O)c1nc(C)c(Sn(CCCC)CCC)n1</chem>	
	 <chem>Cc1nc(C)c(Sn(CCCC)CCC)n1</chem>	
	 <chem>Fc1nc(Sn(CCCC)CCC)cc1</chem>	
	 <chem>Fc1nc(Sn(CCCC)CCC)nc1</chem>	
	 <chem>Fc1nc(Sn(CCCC)CCC)cc1</chem>	
	 <chem>CC(=O)Nc1nc(Sn(C)C)cc1</chem>	
	 <chem>CCOC(=O)Nc1nc(Sn(C)C)cc1</chem>	

	 <chem>CCCC[Sn](CCCC)(CCCC)c1ccnc(=O)n1</chem>	
	 <chem>CCCC[Sn](CCCC)(CCCC)c1ccnc(C(=O)N)cn1</chem>	
	 <chem>CCCC[Sn](CCCC)(CCCC)c1ccnc(C(=O)N(C)C)cn1</chem>	
	 <chem>CCCC[Sn](CCCC)(CCCC)c1cc2nc(C(=O)N3CCCC3)cnc2n1</chem>	
	 <chem>CCCC[Sn](CCCC)(CCCC)c1ccnc(C(=O)NCC)cn1</chem>	
	 <chem>C[Sn](C)(C)c1cc2nc(F)c(=O)[nH]2n1</chem>	
	 <chem>C[Sn](C)(C)c1cc2nc(OC)c(=O)[nH]2n1</chem>	
	 <chem>C[Sn](C)(C)c1cc2nc(O)c(=O)[nH]2n1</chem>	






		
		
		Turck, A.; et.al Lab.. J. Organomet. Chem. (1991), 412(3), 301- 10. Metallation of 2,6- dicloropyrazine and quench with Bu3SnCl
		Analogous to Lehn, L. M., et al. Chem. Eur. J. 2000, 6, 4133.
		
		
		
		






		
		
		
		
		
		
		
		<p>Metallation of 1-trityl-4-iodo imidazole (prepared in Takahashi, Kazuyuki; Kirk, Kenneth L.; Cohen, Louis A. Lab. Chem., Natl. Inst. Arthritis Diabetes Dig. Kidney Dis., Bethesda, MD, USA. J. Labelled Compd. Radiopharm. (1986), 23(1), 1-8)</p>













		using $t\text{BuLi}$ in THF at $-78$ and quenching with $\text{Bu}_3\text{SnCl}$ . Detritylate with TFA or aq HCl after coupling to azaindole core.
		Metallation of 1-methyl-4-iodoimidazole (prepared in Takahashi, Kazuyuki; Kirk, Kenneth L.; Cohen, Louis A. Lab. Chem., Natl. Inst. Arthritis Diabetes Dig. Kidney Dis., Bethesda, MD, USA. J. Labelled Compd. Radiopharm. (1986), 23(1), 1-8) using $t\text{BuLi}$ in THF at $-78$ and quenching with $\text{Bu}_3\text{SnCl}$ . El Borai, M.; Moustafa, A. H.; Anwar, M.; Abdel Hay, F. I. The bromo derivative is described in Pol. J. Chem. (1981), 55(7-8), 1659-65 and can be used to generate the tin reagent via transmetallation.
		
		4,5difluoroimidazole prepared as in Dolensky, Bohumil;et.al, USA. J. Fluorine Chem. (2001), 107(1), 147-148.
		Dolensky, Bohumil;et.al, USA. J. Fluorine Chem. (2001), 107(1), 147-148.
		











Select general procedures, via  $S_NAr$  reactions, for the preparation of starting materials for Tin agents









- 5 a. Preparation of 2-bromo-5-substituted-pyrazine, 5-bromo-2-substituted-thiazole, 2-substituted-thiazole, 4-chloro-6-substituted-pyrimidine and 5-bromo-2-substituted-pyrimidine

















20 To a flask, an appropriate pyrazine, pyrimidine or thiazole (1.0 equivalent) and a nucleophile, such as amine, alcohol or thio-derivatives in one equivalent or an excess amount were combined in a solvent such as THF, DMF or alcohol, with or without an addition of NaH. The reaction was either stirred at room temperature or under heating for one to three days. After all the solvents were removed, the residue was partitioned between saturated  $\text{NaHCO}_3$  and EtOAc. The aqueous layer was extracted with ethyl acetate and the combined extracts were concentrated *in vacuo* to give a residue, which was purified by silica gel chromatography to afford the desired product.

Starting Material	Product	Reaction Condition	Rf (minutes)	MS (M+H)+ Cald.	MS (M+H)+ Obsv.
 SM-01		SM-01 (2g) Piperazine (10 g), THF (50 ml), r.t.	0.56 (column G)	243.02	243.03
 SM-01		SM-01 (1 g), MeNH <sub>2</sub> (2M in THF, 100 ml), r.t.	0.89 (column E)	187.93	187.98
 SM-01		SM-01 (1g), Me <sub>2</sub> NH (2M in THF, 100 ml), r.t.	1.19 (column E)	201.92	202.00
 SM-01		SM-01 (1 g), MeONa (0.5M in MeOH, 100 ml), r.t.	1.05 (column E)	188.91	188.97
 SM-01		SM-01 (50 mg), NaH (17 mg), 2-amino-1,3,4-thiadiazole (25 mg), DMF 5 ml) r.t.	1.21 (column E)	257.94	257.89


 SM-01		SM-01 (50 mg), NaH (17 mg), N-benzylpiperazine (25 mg), DMF (5 ml) r.t.	1.04 (column E)	333.07	332.99
 SM-01		SM-01 (50 mg), NaH (17 mg), N,N-diethylamino-ethanol (0.033 ml), DMF (5 ml) r.t.	0.72 (column E)	274.06	273.97
 SM-02		SM-02 (2 g) Piperazine (10 g), THF (50 ml), r.t.	0.89 (column E)	247.99	247.97
 SM-02		SM-05 (1 g), Me <sub>2</sub> NH (2M in THF, 100 ml), r.t.	0.65 (column E)	206.89	206.96
 SM-02		SM-02 (1 g), MeONa (0.5M in MeOH, 100 ml), r.t.	1.35 (column E)	193.93	193.84

 SM-02	 SM-03	SM-03 (50 mg), NaH (16 mg), imidazole (77 mg), DMF 5 ml) r.t.	0.89 (column E)	229.94	229.83
 SM-02	 SM-02	SM-02 (50 mg), NaH (16 mg), N-benzylpiperazine (30 mg), DMF 5 ml) r.t.	1.02 (column E)	338.03	337.98
 SM-02	 SM-02	SM-02 (50 mg), NaH (16 mg), N,N-diethylamino-ethanol (0.033 ml), DMF 5 ml) r.t.	0.83 (column E)	279.02	278.95
 SM-03	 SM-03	SM-03 (50 mg), NaH (25 mg), imidazole (25 mg), DMF 5 ml) r.t.	0.31 (column E)	151.91	152.03

 SM-03		SM-03 (50 mg), NaH (25 mg), N-benzylpiperazine (37 mg), DMF (5 ml) r.t.	0.66 (column E)	260.07	260.12
 SM-03		SM-03 (50 mg), NaH (25 mg), N,N-diethylamino-ethanol (0.05 ml), DMF (5 ml) r.t.	0.46 (column E)	201.11	201.02
 SM-04		SM-04 (1g), MeONa (0.5M in MeOH, 13.52 ml), r.t.	0.86 (column E)	145.02	144.99
 SM-04		SM-04 (1g), MeNH <sub>2</sub> (2M in THF, 100ml), r.t.	0.46 (column E),	144.03	143.96
 SM-05		SM-05 (1g), MeONa (0.5M in MeOH, 100ml), 1 day, r.t.	0.91 (column E)	188.97	188.91

 SM-05	 SM-05	SM-05 (1g), MeNH <sub>2</sub> (2M in THF, 100 ml), r.t.	0.84 (column E)	187.99	187.94
 SM-05	 SM-05	SM-05 (1g), Me <sub>2</sub> NH (2M in THF, 100 ml), r.t.	1.24 (column E)	202.00	201.98

b. Preparation of 2-bromo-5,6-disubstituted-pyrazine













5

*Step one*

To a flask, an appropriate pyrazine (1.0 equivalent) and a nucleophile, such as amine or sodium alkoxide in an excess amount were combined in a solvent such as water or THF or without solvent. The reaction was either stirred at room temperature or under heating for one to three days. After all the solvents were removed, a residue was collected and used in the further steps without any purification.

10







Starting Material	Product	Reaction Condition	Rf (minutes)	MS (M+H)+ Cald.	MS (M+H)+ Obsv.
 SM-06		SM-06 (100 mg), propylamine (2 ml), r.t.	1.28 (column C)	172.06	172.09
 SM-06		SM-06 (100 mg), Me <sub>2</sub> NH (2M in THF, 10 ml) or Me <sub>2</sub> NH (40% in water, 10 ml), r.t.	1.21 (column C)	158.05	158.07
 SM-06		SM-06 (100 mg), Me <sub>2</sub> NH (40% in water, 10 ml), 100°C	0.49 (column C)	167.13	167.19
 SM-06		SM-06 (100 mg), MeNH <sub>2</sub> (2M in THF, 10 ml), r.t.	0.72 (column C)	144.03	144.07
 SM-06		SM-06 (100 mg), NH <sub>4</sub> OH (10 ml), 100°C	0.41 (column C)	162.04 (M+Me OH+H) <sup>+</sup>	162.06 (M+MeO H+H) <sup>+</sup>

*Step two*

- 5 To a flask, the crude pyrazine derivative obtained from the step one (1.0 equivalent) and a nucleophile, such as amine or sodium alkoxide in an excess amount were combined in a solvent such as water or THF or without solvent. The reaction was either stirred at room temperature or under heating for one to three days. After









all the solvents were removed, a residue was collected and used in the further steps without any purification.


Starting Material	Product	Reaction Condition	Rf (minutes)	MS (M+H)+ Cald.	MS (M+H)+ Obsv.
 SM-07		SM-07 (2 g), MeONa (12.5 wt%, 100ml, 100°C	0.28 (column C)	140.08	140.14
 SM-08		SM-08 (2 g), MeONa (12.5 wt%, 20ml), 100°C	0.28 (column C)	158.13	158.09
 SM-07		SM-07 (2 g), MeNH <sub>2</sub> (40% in water, 100 ml), 110°C	0.34 (column C)	139.10	139.13

### 5 Step three

To a flask, the crude pyrazine derivative obtained from the step two (1.0 equivalent) was dissolved in methylene chloride. A slightly excess of bromine was then added into the mixed solution. The reaction was stirred at room temperature for ten hours. After all the solvents were removed, a residue was collected and purified by silica gel chromatography to afford the desired product.

Starting Material	Product	Reaction Condition	Rf (minutes)	MS (M+H) <sup>+</sup> Cald.	MS (M+H) <sup>+</sup> Obsv.
 SM-09		SM-09 (5 g), bromine (1.34 ml), CH <sub>2</sub> Cl <sub>2</sub> (100 ml)	1.77 (column C)	249.97	250.02
 SM-10		SM-10 (2 g), bromine (0.72 ml), CH <sub>2</sub> Cl <sub>2</sub> (20 ml)	1.13 (column C)	217.99	217.98
 SM-11		SM-11 (2 g), bromine (0.72 ml), CH <sub>2</sub> Cl <sub>2</sub> (20 ml)	0.98 (column C)	203.98	203.99



General procedure of the preparation of 2-alkyl-5-bromo-pyrimidine:




5

To a sealed tube, 5-bromo-2-iodopyrimidine (1.0 equivalent), tri-alkylaluminum (1.5 equivalent) and tetrakis-triphenylphosphine palladium, Pd(Ph<sub>3</sub>P)<sub>4</sub> (1.0 mol%) were combined in dioxane (10 mL). The reaction was heated at 110-120 °C for 10 h. After the mixture cooled down to room temperature, it was poured into water. The solution was extracted with ethyl acetate and the combined extracts were concentrated *in vacuo* to give a residue which was purified by silica gel chromatography to afford the desired 2-alkyl-5-bromopyrimidine product.

10

R <sub>3</sub> Al	Product	R <sub>f</sub> (minutes)	MS (M+H) <sup>+</sup> Cald.	MS (M+H) <sup>+</sup> Obsv.
Me <sub>3</sub> Al		0.90 (column E)	172.94	172.97
(i-Bu) <sub>3</sub> Al		1.45 (column E)	215.02	214.99

Prep of triazine stannane for Stille coupling to prepare examples of claim 1. (the sulfur can then be removed with Raney Nickel to give additional desulfurized triazines)




2,2,6,6-tetramethylpiperidine (2.0ml, 11.81mmol) in 30ml of THF was cooled to -78°C and treated with n-butyllithium (4.7ml, 11.81mmol, 2.5M in hexane). After stirring 30min at 0°C, the reaction was cooled to -78°C again and 3-methylthio-1,2,4-triazine (1.0g, 7.87mmol) was added. The resulting solution was stirred at -78°C for 30min before tributyltin chloride (2.1ml, 7.87mmol) was added. The reaction was kept at -78°C for 1hr, then quenched with water. The THF solvent was removed on rotary evaporator and the remaining solution was extracted with ethylacetate. The organic layer was dried over MgSO<sub>4</sub>, filtered and the filtrate was concentrated. The residue was chromatographed to afford 96mg of 3-methylthio-6-tributyltin-1,2,4-triazine.

<sup>1</sup>H NMR (300Hz, CHCl<sub>3</sub>): 8.83 (s, 1H); 2.62 (s, 3H); 2.04 - 0.79 (m, 27H).

LC/MS: (ES<sup>+</sup>) M/Z (M+H)<sup>+</sup> = 418, RT = 2.29min.


## Intermediate 15



5

To a mixture of **5q** (50 mg, 105  $\mu\text{mol}$ ) and  $\text{Pd}(\text{PPh}_3)_4$  (25 mg, 21  $\mu\text{mol}$ ) was added 1,4-dioxane (1 ml) and vinyl tributylstannane (50 mg, 158  $\mu\text{mol}$ ). The reaction mixture was heated in a sealed tube at 145°C for 3 hours. After cooling to ambient temperature, the reaction mixture was added MeOH (4 ml) and then filtered. The filtrate was purified by preparative reverse phase HPLC to give the TFA salt of **Intermediate 13** using the method: Start %B = 30, Final %B = 75, Gradient time = 20 min, Flow Rate = 25 ml/min, Column : YMC C18 5 $\mu\text{m}$  20 x 100mm, Fraction Collection: 7.92 – 8.58 min.  $^1\text{H}$  NMR: ( $\text{CD}_3\text{OD}$ )  $\delta$  8.61 (s, 1H), 8.37 (s, 1H), 7.47 (b s, 5H), 7.31 (dd,  $J$  = 17.3, 11.3, 1H), 6.50 (d,  $J$  = 17.3, 1H), 5.97 (d,  $J$  = 11.3, 1H), 3.97 – 3.38 (b m, 8H); LC/MS: ( $\text{ES}^+$ )  $m/z$  ( $\text{M}+\text{H}$ ) $^+$  = 423, 425; HPLC  $R_t$  = 1.887.

## Intermediate 14




20

To a mixture of **intermediate 5q** (30 mg, 63  $\mu\text{mol}$ ) and  $\text{Pd}(\text{PPh}_3)_4$  (20 mg, 17  $\mu\text{mol}$ ) was added 1,4-dioxane (1 ml) and 1-tributylstannyl propyne (40 mg, 122  $\mu\text{mol}$ ). The reaction mixture was heated in a sealed tube at 145°C for 2 hours. After cooling to ambient temperature, the reaction mixture was added MeOH (4 ml) and then filtered. The filtrate was purified by preparative reverse phase HPLC to give the TFA salt of **intermediate 14** (1-(4-Benzoyl-piperazin-1-yl)-2-(4-chloro-7-prop-1-ynyl-1H-pyrrolo[2,3-c]pyridin-3-yl)-ethane-1,2-dione) using the method: Start %B = 20, Final %B = 80, Gradient time = 20 min, Flow Rate = 25 ml/min, Column : YMC C18 5 $\mu\text{m}$


20 x 100mm, Fraction Collection: 8.74 – 9.00 min.  $^1\text{H}$  NMR: ( $\text{CD}_3\text{OD}$ )  $\delta$  8.47 (s, 1H), 8.27 (s, 1H), 7.46 (b s, 5H), 3.82 – 3.34 (b m, 8H), 2.26 (s, 3H); LC/MS: (ES+)  $m/z$  ( $\text{M}+\text{H}$ ) $^+$  = 435, 437; HPLC  $R_t$  = 2.123.

5

**Intermediate 15**

- To a solution of intermediate 5q (50 mg, 0.11 mmol) in DMF (1 ml) was added CuCN (30 mg, 0.335 mmol). The reaction mixture was heated at 170°C for 30 min.
- 10 After cooling to ambient temperature, the reaction mixture was diluted with MeOH (15 ml), filtered under gravity, and the filtrate evaporated *in vacuo* to afford a brownish residue. To the residue in EtOH (3 ml) at ambient temperature was bubbled hydrogen chloride gas for 10 minutes to give a yellow solution, which was purified by preparative reverse phase HPLC using the method: Start %B = 15, Final %B = 85,
- 15 Gradient time = 15 min, Flow Rate = 40 ml/min, Column : XTERRA C18 5  $\mu\text{m}$  30 x 100 mm, Fraction Collection: 10.40 – 10.85 min;  $^1\text{H}$  NMR: ( $\text{CD}_3\text{OD}$ ) 8.35 (s, 1H), 8.33 (s, 1H), 7.42 (b s, 5H), 3.95 – 3.41 (b m, 8H); LC/MS: (ES+)  $m/z$  ( $\text{M}+\text{H}$ ) $^+$  = 440, 442; HPLC  $R_t$  = 1.820.

20

**Intermediate 16****Preparation of intermediate 16:**

- 25 To a suspension of intermediate 15 (6 mg, 13  $\mu\text{mol}$ ) in a mixture of AcOH (0.5 ml) and  $\text{Ac}_2\text{O}$  (1.0 ml) at 0°C was charged with sodium nitrite (17 mg, 246  $\mu\text{mol}$ ). The reaction mixture was stirred at 0°C for 30 min. and then at ambient temperature for 1 hour. After addition of MeOH (4 ml), the reaction mixture was purified by preparative reverse phase HPLC to give the TFA solvate of the title compound using
- 30 the method: Start %B = 15, Final %B = 80, Gradient time = 15 min, Flow Rate = 25


ml/min, Column : YMC C18 5 $\mu$ m 20 x 100mm, Fraction Collection: 9.48 – 10.03 min. <sup>1</sup>H NMR: (DMSO-*d*<sub>6</sub>)  $\delta$  12.76 (s, 1H), 8.48 (s, 1H), 8.32 (d, *J* = 3.0, 1H), 7.44 (b s, 5H), 3.97 – 3.47 (b m, overlapping with water peak, 8H); LC/MS: (ES+) *m/z* (M+H)<sup>+</sup> = 441, 443; HPLC *R*<sub>t</sub> = 1.530.

- 5    Ref: Amide hydrolysis: Evans, D. A.; Carter, P. H.; Dinsmore, C. J.; Barrow, J. C.; Katz, J. L.; Kung, D. W. *Tetrahedron Lett.* 1997, 38, 4535 and references cited therein.

### Preparation of Compounds of Formula I

10

#### EXAMPLE 1



- Typical procedure for coupling azaindole with aromatic boron reagent (An example of the general procedure described below for examples 2-14): Preparation of 1-benzoyl-3-(R)-methyl-4-[(7-(4-fluorophenyl)-6-azaindol-3-yl)-oxoacetyl]piperazine is an example of Step E as described in Scheme 15. To a sealed tube, 1-(benzoyl)-3-(R)-methyl-4-[(7-chloro-6-azaindol-3-yl)-oxoacetyl]piperazine, Intermediate 5a, (20 mg, 0.049 mmol), 4-fluorophenylboronic acid, Intermediate 14a-9, (8.2 mg, 0.059 mmol), Pd(Ph<sub>3</sub>P)<sub>4</sub> (5 mg) and K<sub>2</sub>CO<sub>3</sub> (20 mg, 0.14 mmol) were combined in 1.5 mL of DMF and 1.5 mL of water. The reaction was heated at 110-120 °C for 10 h. After the mixture cooled down to rt, it was poured into 20 mL of water. The solution was extracted with EtOAc (4 x 20 mL). The combined extract was concentrated to give a residue which was purified using a Shimadzu automated preparative HPLC System to give compound 1-benzoyl-3-(R)-methyl-4-[(7-(4-fluorophenyl)-6-azaindol-3-yl)-oxoacetyl]piperazine (1.8 mg, 7.9%). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>24</sub>FN<sub>4</sub>O<sub>3</sub>: 471.18; found 471.08. HPLC retention time: 1.12 minutes (column A).
- 15  
20  
25

30

### EXAMPLES 2-14


Examples 2-14 were prepared according to the following general method in a manner analogous to the preparation of Example 1.

5

*Typical procedure for coupling azaindole with aromatic boron reagent:* To a sealed tube, an appropriately substituted azaindole intermediate (0.049 mmol), an appropriate boronic acid derivative (0.059 mmol), Pd(PH<sub>3</sub>P)<sub>4</sub> (5 mg) and K<sub>2</sub>CO<sub>3</sub> (20 mg, 0.14 mmol) were combined in 1.5 mL of DMF and 1.5 mL of water. The reaction was heated at 110-120 °C for 10 h. After the mixture cooled down to rt, it was poured into 20 mL of water. The solution was extracted with EtOAc (4 x 20 mL). The combined extract was concentrated *in vacuo* to give a residue which was purified using a Shimadzu automated preparative HPLC System to provide the desired compound.


15

### EXAMPLE 2




20

Example 2, was prepared according to the general method described above starting from Intermediate 5g and 4-chlorophenyl boronic acid, Intermediate 14a-10, to provide 1-benzoyl-4-[(7-(4-chlorophenyl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>24</sub>FN<sub>4</sub>O<sub>3</sub>: 473.14; found 473.13. HPLC retention time: 1.43 minutes (column B).

**EXAMPLE 3**


Example 3, was prepared according to the general method described above  
5 starting from Intermediate 5a and 3-amino-4-methylphenyl boronic acid, Intermediate 14a-11, to provide 1-benzoyl-3-methyl-4-[(7-(3-amino-4-methylphenyl)-6-azaindol-3-yl)-oxoacetyl] piperazine. MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{27}H_{24}ClN_4O_3$ : 482.22; found 482.25. HPLC retention time: 1.35 minutes (column B).

10


**EXAMPLE 4**

Example 4, was prepared according to the general method described above  
15 starting from Intermediate 5g and 4-hydroxycarbonylphenyl boronic acid, Intermediate 14a-12, to provide 1-benzoyl-4-[(7-(4-carboxyphenyl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{27}H_{24}ClN_4O_3$ : 483.17; found 483.10. HPLC retention time: 1.00 minutes (column A).




**EXAMPLE 5**

5        Example 5, was prepared according to the general method described above  
from 1-benzoyl-3-methyl-4-[(7-chloro-6-azaindol-3-yl)-oxoacetyl]piperazine and  
3,4-methylenedioxyphenyl boronic acid, Intermediate 14a-13, to provide 1-benzoyl-  
3-methyl-4-[(7-(3,4-methylenedioxyphenyl)-6-azaindol-3-yl)-oxoacetyl]piperazine.  
MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>28</sub>H<sub>23</sub>N<sub>4</sub>O<sub>3</sub>: 497.18; found 497.03. HPLC retention time:  
10    1.41 minutes (column B).


**EXAMPLE 6**

15        Example 6, was prepared according to the general method described above  
starting from Intermediate 5a and furan-2-yl boronic acid to provide 1-benzoyl-3-  
methyl-4-[(7-(furan-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup>  
Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>4</sub>O<sub>4</sub>: 443.17; found 443.12. HPLC retention time: 1.20 minutes  
20    (column A).

**EXAMPLE 7**

5        Example 7, was prepared according to the general method described above starting from Intermediate 5g and furan-2-yl boronic acid to provide 1-benzoyl-4-[(7-(furan-2-yl)-6-azaindol-3-yl)-oxoacetyl] piperazine MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>24</sub>H<sub>21</sub>N<sub>4</sub>O<sub>4</sub>: 429.16; found 428.98. HPLC retention time: 1.36 minutes (column A).


10

**EXAMPLE 8**

15        Example 8, was prepared according to the general method described above starting from Intermediate 5g and benzofuran-2-yl boronic acid to provide 1-benzoyl-4-[(7-(benzofuran-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>28</sub>H<sub>23</sub>N<sub>4</sub>O<sub>4</sub>: 479.17; found 479.09. HPLC retention time: 1.67 minutes (column B).


20

25


**EXAMPLE 9**

5            Example 9, was prepared according to the general method described above starting from Intermediate 5a and thien-2-yl boronic acid to provide 1-(benzoyl)-3-methyl-4-[(7-(thien-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>4</sub>O<sub>3</sub>S: 459.15; found 459.10. HPLC retention time: 1.20 minutes (column A).

10


**EXAMPLE 10**

15            Example 10, was prepared according to the general method described above starting from Intermediate 5g and pyridin-4-yl boronic acid to provide 1-(benzoyl)-4-[(7-(pyridin-4-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>22</sub>N<sub>5</sub>O<sub>3</sub>: 440.17; found 440.10. HPLC retention time: 0.97 minutes (column A).

**EXAMPLE 11**

5            Example 11, was prepared according to the general method described above starting from Intermediate 5g and quinolin-8-yl boronic acid, Intermediate 14a-14, to provide 1-benzoyl-4-[(7-(quinolin-8-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS  $m/z$ : (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>22</sub>N<sub>5</sub>O<sub>3</sub>: 490.19; found 490.09. HPLC retention time: 1.34 minutes (column B).


10

**EXAMPLE 12**

15            Example 12, was prepared according to the general method described above starting from Intermediate 5a and 2,4-dimethoxypyrimidin-5-yl boronic acid, Intermediate 14a-4, to provide 1-benzoyl-3-methyl-4-[(7-(2,4-dimethoxy-pyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS  $m/z$ : (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>27</sub>N<sub>6</sub>O<sub>5</sub>: 515.20; found 515.28. HPLC retention time: 1.17 minutes (column B).


20

### EXAMPLE 13




Example 13, was prepared according to the general method described above starting from Intermediate 5b and 2,4-dimethoxypyrimidin-5-yl boronic acid, Intermediate 14a-4, to provide 1-benzoyl-4-[(4-methoxy-7-(2,4-dimethoxypyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 8.71 (s, 1H), 8.64 (s, 1H), 8.21 (s, 1H), 7.48 (s, 5H), 4.15 (s, 3H), 4.13 (s, 3H), 3.84 (s, 3H), 3.64 – 3.34 (m, 8H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>29</sub>H<sub>33</sub>N<sub>6</sub>O<sub>6</sub>: 531.20; found 531.26. HPLC retention time: 1.09 minutes (column A).

### EXAMPLE 14



Example 14, was prepared according to the general method described above starting from Intermediate 5b and pyridin-4-yl boronic acid to provide 1-benzoyl-4-[(4-methoxy-7-(pyridin-4-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>24</sub>N<sub>5</sub>O<sub>4</sub>: 470.18; found 470.32. HPLC retention time: 1.02 minutes (column A).

**EXAMPLE 15**

5            *Typical procedure for coupling azaindole with aromatic tin reagent (An example of the general procedure described below for examples 16-53):* Preparation of Example 15, 1-benzoyl-4-[(4-methoxy-7-(2-(1,1-dimethylethylaminocarbonyl)-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine is an example of Step E as described in Scheme 15. To a sealed tube, 1-benzoyl-4-[(7-chloro-4-methoxy-6-azaindol-3-yl)-oxoacetyl]piperazine, Intermediate 5b, (20 mg), 2-(1,1-

10            dimethylethylaminocarbonyl)-5-tributylstannyl-pyrazine (1.2 equivalents, 27 mg.) and Pd(Ph<sub>3</sub>P)<sub>4</sub> (1 mg) were combined in 1.5 mL of dioxane. The reaction was heated at 110-120 °C for 10 h. After the mixture cooled down to room temperature, it was poured into 5 mL of water. The solution was extracted with EtOAc (4 x 5 mL). The

15            combined extract was concentrated *in vacuo* to give a residue which was purified using a Shimadzu automated preparative HPLC System to give compound 1-benzoyl-4-[(4-methoxy-7-(2-(1,1-dimethylethylaminocarbonyl)-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine (1 mg); MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>30</sub>H<sub>32</sub>N<sub>7</sub>O<sub>5</sub>; 570.25; found 570.43. HPLC retention time: 1.83 minutes (column B).

20

**EXAMPLES 16-54**


Examples 16-54 were prepared according to the following general procedure by a method analogous to the method described for the preparation of Example 15.

25

*Typical procedure for coupling azaindole with aromatic tin reagent:* To a sealed tube, an appropriate azaindole (0.049 mmol), an appropriate stannane (0.059 mmol) and Pd(Ph<sub>3</sub>P)<sub>4</sub> (1 mg) were combined in 1.5 mL of dioxane. The reaction was heated at 110-120 °C for 10 h. After the mixture cooled down to rt, it was poured


into 5 mL of water. The solution was extracted with EtOAc (4 x 5 mL). The combined extract was concentrated to give a residue which was purified using a Shimadzu automated preparative HPLC System to provide the desired compound.

5


**EXAMPLE 16**

Example 16, was prepared according to the general method described above  
10 starting from Intermediate 5a and pyrimidin-5-yl tributyltin, Intermediate 14-9, to provide 1-benzoyl-3-(R)-methyl-4-[(7-(pyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{25}H_{23}N_6O_3$ : 455.18; found 455.17. HPLC retention time: 1.33 minutes (column B).

15


**EXAMPLE 17**

Example 17, was prepared according to the general method described above  
20 starting from Intermediate 5g and pyrimidin-5-yl tributyltin, Intermediate 14-9, to provide 1-benzoyl-4-[(7-(pyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{25}H_{23}N_6O_3$ : 441.17; found 441.07. HPLC retention time: 1.30 minutes (column B).

**EXAMPLE 18**


5            Example 18, was prepared according to the general method described above starting from Intermediate 5a and pyridin-3-yl tributyltin, Intermediate 14a-2, to provide 1-benzoyl-3-(R)-methyl-4-[(7-(pyridin-3-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>24</sub>N<sub>5</sub>O<sub>3</sub>; 454.19; found 454.17. HPLC retention time: 1.11 minutes (column A).

10

**EXAMPLE 19**

15            Example 19, was prepared according to the general method described above starting from Intermediate 5g and pyridin-2-yl tributyltin, Intermediate 14a-19, to provide 1-benzoyl-4-[(7-(pyridin-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>22</sub>N<sub>5</sub>O<sub>3</sub>; 440.17; found 440.07. HPLC retention time: 1.40 minutes (column B).


20

**EXAMPLE 20**



Example 20, was prepared according to the general method described above starting from Intermediate 5a and thiazol-2-yl tributyltin, Intermediate 14a-21, to provide 1-benzoyl-3-(R)-methyl-4-[(7-(thiazol-2-yl)-6-azaindol-3-yl)oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>24</sub>H<sub>22</sub>N<sub>5</sub>O<sub>3</sub>S: 460.14; found 460.15. HPLC retention time: 1.48 minutes (column B).

#### EXAMPLE 21




10

Example 21, was prepared according to the general method described above starting from Intermediate 5g and thiazol-2-yl tributyltin, Intermediate 14a-21, to provide 1-benzoyl-4-[(7-(thiazol-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>23</sub>H<sub>20</sub>N<sub>5</sub>O<sub>3</sub>S: 446.13; found 446.03. HPLC retention time: 1.44 minutes (column B).

15


#### EXAMPLE 22



20


Example 22, was prepared according to the general method described above starting from Intermediate 5b and 1-methylpyrazol-3-yl tributyltin, to provide 1-benzoyl-4-[(4-methoxy-7-(1-methyl-pyrazol-3-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>25</sub>N<sub>6</sub>O<sub>4</sub>: 473.19; found 473.28. HPLC retention time: 1.18 minutes (column B).

25

**EXAMPLE 23**


Example 23, was prepared according to the general method described above  
5 starting from Intermediate 5b and Intermediate 14-9 to provide 1-benzoyl-4-[(4-methoxy-7-(pyridazin-4-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>6</sub>O<sub>4</sub>: 471.18; found 471.26. HPLC retention time: 1.20 minutes (column B).

10

**EXAMPLE 24**


Example 24, was prepared according to the general method described above  
15 starting from Intermediate 5b and 2-aminopyrimidin-5-yl tributyltin, to provide 1-benzoyl-4-[(4-methoxy-7-(2-amino-pyrimidin-5-yl))-6-azaindol-3-yl]-oxoacetyl]piperazine MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>24</sub>N<sub>7</sub>O<sub>4</sub>: 486.19; found 486.24. HPLC retention time: 1.19 minutes (column A).

20

**EXAMPLE 25**

5            Example 25, was prepared according to the general method described above starting from Intermediate 5b and pyridin-3-yl tributyltin, Intermediate 14a-2, to provide 1-benzoyl-4-[(4-methoxy-7-(pyridin-3-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>24</sub>N<sub>5</sub>O<sub>4</sub>: 470.18; found 470.19. HPLC retention time: 1.04 minutes (column A).


10

**EXAMPLE 26**

15            Example 26, was prepared according to the general method described above starting from Intermediate 5b and 2-aminopyrazin-5-yl trimethyltin, Intermediate 14-28, to provide 1-benzoyl-4-[(4-methoxy-7-(2-amino-pyrazin-5-yl))-6-azaindol-3-yl]-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>24</sub>N<sub>7</sub>O<sub>4</sub>: 486.19; found 470.19. HPLC retention time: 1.13 minutes (column B).


20

25

**EXAMPLE 27**

5            Example 27, was prepared according to the general method described above starting from Intermediate 5b and 1-methylimidazol-2-yl trimethyltin, Intermediate 14-5, to provide 1-benzoyl-4-[(4-methoxy-7-(1-methyl-imidazol-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>25</sub>N<sub>6</sub>O<sub>4</sub>: 473.18; found 473.27. HPLC retention time: 1.07 minutes (column B).


10

**EXAMPLE 28**

15            Example 28, was prepared according to the general method described above starting from Intermediate 5b and 1-methylpyrrol-2-yl tributyltin, Intermediate 14a-15, to provide 1-benzoyl-4-[(4-methoxy-7-(1-methyl-pyrrol-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>26</sub>N<sub>5</sub>O<sub>4</sub>: 472.20; found 470.26. HPLC retention time: 1.11 minutes (column A).


20

25

**EXAMPLE 29**

5            Example 29, was prepared according to the general method described above starting from Intermediate 5i and 1-methylpyrazol-3-yl tributyltin, to provide 1-benzoyl-4-[(4-fluoro-7-(1-methylpyrazol-3-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{24}H_{22}FN_6O_3$ : 461.17; found 461.24. HPLC retention time: 1.36 minutes (column A).


10

**EXAMPLE 30**

15            Example 30, was prepared according to the general method described above starting from Intermediate 5i and pyridazin-4-yl tributyltin, Intermediate 14-8, to provide 1-benzoyl-4-[(4-fluoro-7-(pyridazin-4-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine  $^1H$  NMR (500 MHz,  $CD_3OD$ )  $\delta$  9.72 (s, 1H), 9.39 (s, 1H), 8.42 (m, 2H), 8.22 (s, 1H), 7.47 (s, 5H), 3.84 – 3.38 (m, 8H). MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{24}H_{20}FN_6O_3$ : 459.16; found 459.25. HPLC retention time: 1.26 minutes (column A).


20

25

**EXAMPLE 32**

5            Example 32, was prepared according to the general method described above starting from Intermediate 5b and pyrazin-2-yl tributyltin, Intermediate 14a-1, to provide 1-benzoyl-4-[(4-methoxy-7-(pyrazin-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>6</sub>O<sub>3</sub>: 471.18; found 471.17. HPLC retention time: 1.35 minutes (column A).


10

**EXAMPLE 33**

15            Example 33, was prepared according to the general method described above starting from Intermediate 5a and pyrazin-2-yl tributyltin, Intermediate 14a-1, to provide 1-benzoyl-3-(R)-methyl-4-[(7-(pyrazin-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>6</sub>O<sub>3</sub>: 455.18; found 455.26. HPLC retention time: 1.46 minutes (column A).


20

25

**EXAMPLE 34**

5            Example 34, was prepared according to the general method described above starting from Intermediate 5g and pyrazin-2-yl tributyltin, Intermediate 14a-1, to provide 1-benzoyl-4-[(7-(pyrazin-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{24}H_{21}N_6O_3$ : 441.17; found 441.22. HPLC retention time: 1.22 minutes (column A).


10

**EXAMPLE 35**

15            Example 35, was prepared according to the general method described above starting from Intermediate 5b and thiazol-2-yl tributyltin, Intermediate 14a-21, to provide 1-(benzoyl)-4-[(4-methoxy-7-(thiazol-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{24}H_{22}N_5O_3S$ : 476.14; found 476.20. HPLC retention time: 1.25 minutes (column B).


20

25

EXAMPLE 36

5            Example 36, was prepared according to the general method described above starting from Intermediate 5b and pyridin-2-yl tributyltin, Intermediate 14a-19, to provide 1-benzoyl-4-[(4-methoxy-7-(pyridin-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>24</sub>N<sub>5</sub>O<sub>4</sub>: 470.18; found 470.17. HPLC retention time: 1.04 minutes (column A).


10

EXAMPLE 37

15            Example 37, was prepared according to the general method described above starting from Intermediate 5j and thiazol-2-yl tributyltin, Intermediate 14a-21, to provide 1-benzoyl-3-(R)-methyl-4-[(4-fluoro-7-(thiazol-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>24</sub>H<sub>21</sub>FN<sub>5</sub>O<sub>3</sub>S: 478.13; found 478.13. HPLC retention time: 1.34 minutes (column A).


20



EXAMPLE 38

5            Example 38, was prepared according to the general method described above starting from Intermediate 5i and pyrazol-3-yl tributyltin, to provide 1-benzoyl-4-[(4-fluoro-7-(pyrazol-3-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>23</sub>H<sub>20</sub>FN<sub>6</sub>O<sub>3</sub>: 447.16; found 447.15. HPLC retention time: 1.26 minutes (column A).


10

EXAMPLE 39

15            Example 39, was prepared according to the general method described above starting from Intermediate 5b and pyrazol-3-yl tributyltin, to provide 1-benzoyl-4-[(4-methoxy-7-(pyrazol-3-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>24</sub>H<sub>23</sub>N<sub>6</sub>O<sub>4</sub>: 459.18; found 459.21. HPLC retention time: 1.11 minutes (column A).


20

25

**EXAMPLE 40**

5            Example 40, was prepared according to the general method described above starting from Intermediate 5b and pyrimidin-5-yl tributyltin, Intermediate 14-9, to provide 1-benzoyl-4-[(4-methoxy-7-(pyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>6</sub>O<sub>4</sub>: 471.18; found 471.20. HPLC retention time: 1.61 minutes (column A).


10

**EXAMPLE 41**

15            Example 41, was prepared according to the general method described above starting from Intermediate 5j and pyrimidin-5-yl tributyltin, Intermediate 14-9, to provide 1-benzoyl-3-(R)-methyl-4-[(4-fluoro-7-(pyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 9.26 (m, 3H), 8.39 (m, 2H), 7.56 (m, 5H), 4.72 – 3.12 (m, 7H), 1.40 – 0.91 (m, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>22</sub>FN<sub>6</sub>O<sub>3</sub>: 473.17; found 473.17. HPLC retention time: 1.34 minutes (column A).


20

25

**EXAMPLE 42**

5            Example 42, was prepared according to the general method described above starting from Intermediate 5i and pyrimidin-5-yl tributyltin, Intermediate 14-9, to provide 1-benzoyl-4-[(4-fluoro-7-(pyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>24</sub>H<sub>20</sub>FN<sub>6</sub>O<sub>3</sub>: 459.16; found 459.14. HPLC retention time: 1.28 minutes (column A).


10

**Example 43**

15            Example 43, (R)-1-(benzoyl)-3-methyl-4-[(7-(2,4-dimethoxy-pyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>27</sub>N<sub>6</sub>O<sub>5</sub>: 515.20; found 515.28. HPLC retention time: 1.17 minutes (column B).


20

25

**EXAMPLE 44**


5            Example 44, was prepared according to the general method described above starting from Intermediate 5a and 2,3-dichloropyrazin-5-yl tributyltin, Intermediate 14-66, to provide 1-benzoyl-3-(R)-methyl-4-[(7-(2,3-dichloro-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+Na)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>20</sub>Cl<sub>2</sub>NaN<sub>6</sub>O<sub>3</sub>: 545.09; found 545.29. HPLC retention time: 1.87 minutes (column B).

10

**EXAMPLE 45**


15            Example 45, was prepared according to the general method described above starting from Intermediate 5b and 2-ethoxythiazol-5-yl tributyltin, Intermediate 14-71, to provide 1-benzoyl-4-[(4-methoxy-7-(2-ethoxy-thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>26</sub>N<sub>5</sub>O<sub>5</sub>S: 520.17; found 520.24. HPLC retention time: 1.32 minutes (column A).

20


**EXAMPLE 46**

5            Example 46, was prepared according to the general method described above starting from Intermediate 5b and the 2-amino-pyrazin-6-yl stannane, Intermediate 14-68, to provide 1-benzoyl-4-[(4-methoxy-7-(2-amino-pyrazin-6-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>22</sub>H<sub>24</sub>N<sub>4</sub>O<sub>4</sub>: 486.19; found 486.31. HPLC retention time: 1.22 minutes (column B).

10


**EXAMPLE 47**

15            Example 47, was prepared according to the general method described above starting from Intermediate 5b and 2-methylsulfonylamino-5-(tri-n-butylstannyl)pyrazine, Intermediate 14-69, to provide 1-benzoyl-4-[(7-(2-methylsulfonylamino-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>26</sub>N<sub>7</sub>O<sub>6</sub>S: 564.17; found 564.21. HPLC retention time: 1.24  
20    minutes (column A).

**EXAMPLE 48**

5            Example 48, was prepared according to the general method described above starting from Intermediate 5b and 2,4-dimethoxy-1,3,5-triazin-6-yl tributyltin, Intermediate 14-70, to provide 1-benzoyl-4-[(7-(2,4-dimethoxy-1,3,5-triazin-6-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>26</sub>N<sub>7</sub>O<sub>6</sub>: 532.19; found 532.12. HPLC retention time: 1.28 minutes (column A).


10

**EXAMPLE 49**


15            Example 49, was prepared according to the general method described above starting from Intermediate 5b and pyrimidin-2-yl tributyltin, Intermediate 14-67, to provide 1-benzoyl-4-[(4-methoxy-7-(pyrimidin-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>6</sub>O<sub>4</sub>: 471.18; found 471.29. HPLC retention time: 1.21 minutes (column A).

20


25

**EXAMPLE 50**

Example 50, was prepared from 1-(pyridin-2-yl)-4-[(4-methoxy-7-chloro-6-azaindol-3-yl)-oxoacetyl]piperazine and thiazol-2-yl tributyltin, Intermediate 14a-21, according to the general method above to provide 1-(pyridin-2-yl)-4-[(4-methoxy-7-(thiazol-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{24}H_{25}N_6O_4S$ : 477.13; found 477.22. HPLC retention time: 0.98 minutes (column A).


**EXAMPLE 51**

Example 51, was prepared according to the general method described above starting from Intermediate 5d and pyrimidin-5-yl tributyltin, Intermediate 14-9, to provide 1-benzoyl-3-(R)-methyl-4-[(7-(pyrimidin-5-yl)-4-azaindol-3-yl)-oxoacetyl]piperazine; MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{25}H_{23}N_6O_3$ : 455.18; found 455.16. HPLC retention time: 0.98 minutes (column A).

**EXAMPLE 52**

5        Example 52, was prepared according to the general method described above  
starting from Intermediate 5d and pyrimidin-2-yl tributyltin, Intermediate 14a-1, to  
provide 1-benzoyl-3-(R)-methyl-4-[(7-(pyrazin-2-yl)-4-azaindol-3-yl)-  
oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>6</sub>O<sub>3</sub>: 455.18; found 455.16.  
HPLC retention time: 1.09 minutes (column A).

10


**EXAMPLE 53**

15        Example 53, was prepared according to the general method described above  
starting from Intermediate 5d and thiazol-2-yl tributyltin, Intermediate 14a-21, to  
provide 1-benzoyl-3-(R)-methyl-4-[(7-(thiazol-2-yl)-4-azaindol-3-yl)-  
oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>24</sub>H<sub>22</sub>N<sub>5</sub>O<sub>3</sub>S: 460.14; found  
460.26. HPLC retention time: 1.02 minutes (column A).

20


25




**EXAMPLE 54**

5        Example 54, was prepared according to the general method described above starting from Intermediate 5d and 2-ethoxythiazol-5-yl tributyltin, Intermediate 14-71, to provide 1-benzoyl-3-(R)-methyl-4-[(7-(2-ethoxy-thiazol-5-yl)-4-azaindol-3-yl)-oxoacetyl]piperazine; MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>26</sub>N<sub>5</sub>O<sub>4</sub>S: 504.17; found 4504.18. HPLC retention time: 1.26 minutes (column A).

10

**EXAMPLE 55**


15        The compound of Example 15, 1-benzoyl-4-[(4-methoxy-7-(2-(1,1-dimethylethylaminocarbonyl)-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine (20 mg) was dissolved in 1 drop of concentrated sulfuric acid. After 30 minutes, the mixture was diluted with 2 mL of methanol. The resulting solution was injected into a Shimadzu automated preparative HPLC System and the HPLC purification afforded  
20        the compound of Example 55, 1-benzoyl-4-[(4-methoxy-7-(2-aminocarbonyl-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine (1 mg); MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>24</sub>N<sub>7</sub>O<sub>5</sub>: 514.78; found 514.22. HPLC retention time: 1.44 minutes (column B).

**EXAMPLE 56**

5           An excess of  $\text{NH}_4\text{Cl}$  (27mg) was added into a solution of 1-(benzoyl)-3-(R)-methyl-4-[(6-cyano-7-azaindol-3-yl)-oxoacetyl]piperazine (20 mg) and  $\text{NaN}_3$  (16 mg) in DMF. The reaction was heated to reflux for 12 h. After cooling down, the mixture was concentrated under reduced pressure and the residue was purified using Shimadzu automated preparative HPLC System to give 1-benzoyl-3-(R)-methyl-4-  
10 [(6-(tetrazol-1-yl)-7-azaindol-3-yl)-oxoacetyl]piperazine (6.3mg). MS  $m/z$ :  $(\text{M}+\text{H})^+$  Calc'd for  $\text{C}_{22}\text{H}_{21}\text{N}_8\text{O}_3$ : 445.17; Found 3445.16. HPLC retention time: 1.42 minutes (column B); Column B: PHX-LUNA C18 4.6 x 30 mm.

**EXAMPLE 57**

15




Preparation of 1-benzoyl-3-(R)-methyl-4-[(7-(methoxymethylamino)carbonyl)-4-azaindol-3-yl)-oxoacetyl]piperazine: A mixture  
20 of Intermediate 13 (267 mg), N,O-dimethylhydroxylamine hydrogen chloride (248 mg), carbon tetrabromide (844 mg), pyridine (202 mg) and triphenylphosphine (668 mg) in dichloromethane (10 mL) was stirred at room temperature for 10 hours. After solvent was removed under vacuum, the residue was purified by using silica gel chromatography to afford 1-(benzoyl)-3-(R)-methyl-4-[(7-(methoxymethylamino)carbonyl)-4-azaindol-3-yl)-oxoacetyl]piperazine (56 mg); MS  
25

$m/z$ : (M+H)<sup>+</sup> Calc'd for C<sub>24</sub>H<sub>26</sub>N<sub>5</sub>O<sub>5</sub>: 464.19; found 464.25. HPLC retention time: 1.02 minutes (column A).

### EXAMPLE 58


5



Example 58 was prepared according to the same procedure used in preparing Example 57 with the exception of using Intermediate 11 as a starting material instead of Intermediate 13. The procedure provided 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-(methoxymethylamino)carbonyl)-4-azaindol-3-yl]-oxoacetyl]piperazine; MS  $m/z$ : (M+H)<sup>+</sup> Calc'd for C<sub>24</sub>H<sub>25</sub>ClN<sub>5</sub>O<sub>5</sub>: 498.15; found 498.12. HPLC retention time: 1.39 minutes (column A).

15 General procedure A to prepare CO-NR1R2 from COOH

### EXAMPLE 59



20


Preparation of 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-(methylamino)carbonyl)-4-azaindol-3-yl]-oxoacetyl]piperazine: A mixture of Intermediate 11 (25 mg), methylamine (2M in THF, 0.08 mL), EDC (26 mg), HOBT (11.2 mg) and diisopropylethylamine (43 mg) in tetrahydrofuran (5 mL) was stirred at room temperature for 10 hours. After the solvent was removed under vacuum, the residue was purified by using silica gel chromatography to afford 1-benzoyl-3-(R)-

25

methyl-4-[(5-chloro-7-(methylamino)carbonyl)-4-azaindol-3-yl]-oxoacetyl]piperazine (13.6 mg); MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{23}H_{23}ClN_5O_4$ : 468.14; found 468.03. HPLC retention time: 1.33 minutes (column A).

- 5 This general procedure A is applied to prepare examples 94 and 135:


#### Example 94




- 10 Example 94, 1-benzoyl-4-[(4-methoxy-7-(2-methylaminocarbonyl-furan-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine.  $^1H$  NMR (500 MHz,  $CD_3OD$ )  $\delta$  8.37 (s, 1H), 8.06 (s, 1H), 7.48 – 7.26 (m, 7H), 4.08 (s, 3H), 3.83 – 3.44 (m, 8H), 2.96 (s, 3H). MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{29}H_{26}N_5O_6$ : 516.19; found 516.14. HPLC retention time: 1.35 minutes (column A).

15

#### Example 135



- 20 Example 135, (R)-1-benzoyl-3-methyl-4-[(7-(4-trifluoromethylbenzylamino)carbonyl)-4-azaindol-3-yl]-oxoacetyl]piperazine. MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{30}H_{27}F_3N_5O_4$ : 578.20; found 578.39. HPLC retention time: 1.47 minutes (column G).

**General procedure B to prepare CO-NR1R2 from COOH**

5                    **Preparation of Example 136, (R)-1-benzoyl-3-methyl-4-[(7-(4-methylthiazol-2-yl)aminocarbonyl-4-azaindol-3-yl)-oxoacetyl]piperazine:**

To a solution of (R)-1-benzoyl-3-methyl-4-[(7-hydroxycarbonyl-4-azaindol-3-yl)-oxoacetyl]piperazine (146mg) in DMF (5ml) at room temperature was added  
 10 pentafluorophenyl (70.3mg) followed by EDC (73.23mg). The reaction mixture was stirred at room temperature for 8 hours. The crude product was diluted with methylene chloride and was washed with water, 0.1N HCl and brine. The organic phase was dried over MgSO<sub>4</sub>, filtered and concentrated. The pentafluorophenyl ester was used in the following reaction without further purification.

15


To a stirred solution of 4-methyl-2-amino-thiazole (39.6mg) and Hunig's base (49.4mg) in DMF (5ml) at room temperature was added a solution of pentafluorophenyl ester (1/3 of the product obtained in the previous step described above) in DMF (2 ml). The reaction mixture was stirred at room temperature for 16  
 20 hours. The crude product was diluted with methylene chloride and was washed with Na<sub>2</sub>CO<sub>3</sub> (sat.) and brine. The organic phase was dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was purified using Shimadzu automated preparative HPLC System to give (R)-1-benzoyl-3-methyl-4-[(7-(4-methylthiazol-2-yl)aminocarbonyl-4-azaindol-3-yl)-oxoacetyl]piperazine (3.6mg). MS *m/z*: (M+H)<sup>+</sup> Calc'd for

$C_{26}H_{25}N_6O_4S$ : 517.17; found 517.41. HPLC retention time: 1.25 minutes (column A).

This general procedure B is applied to prepare example 137:

5


### Example 137



10 Example 137, (R)-1-benzoyl-3-methyl-4-[(7-(thiazol-2-yl)aminocarbonyl-4-azaindol-3-yl)-oxoacetyl]piperazine. MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{25}H_{23}N_6O_4S$ : 503.15; found 503.29. HPLC retention time: 1.33 minutes (column A).

### EXAMPLE 60


15



Preparation of 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-(imidazol-2-yl)-4-azaindol-3-yl)-oxoacetyl]piperazine: A mixture of Intermediate 10 (34 mg), glyoxal (40% in water, 0.2 mL) and ammonia acetate (139 mg) in methanol was heated up to reflux for 10 hours. After cooling down, the mixture was concentrated under reduced pressure and the residue was purified using Shimadzu automated preparative HPLC System to provide 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-(imidazol-2-yl)-4-azaindol-3-yl)-oxoacetyl]piperazine (1.8 mg); MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{24}H_{22}ClN_6O_3$ : 477.14; found 477.13. HPLC retention time: 1.17 minutes (column A).


20

25


**EXAMPLE 61**

5            Example 61 was prepared according to the same procedure used for preparing Example 60 with the exception of using methylglyoxal as a starting material instead of glyoxal to provide 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-(4-methyl-imidazol-2-yl)-4-azaindol-3-yl)-oxoacetyl]piperazine MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{25}H_{24}ClN_6O_3$ : 491.16; found 491.13. HPLC retention time: 1.26 minutes (column A).

10

**EXAMPLE 62**


15            Example 62 was prepared according to the same procedure used for preparing Example 60 with the exception of using dimethylglyoxal as a starting material instead of glyoxal to provide 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-(4,5-dimethyl-imidazol-2-yl)-4-azaindol-3-yl)-oxoacetyl]piperazine; MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{26}H_{26}ClN_6O_3$ : 505.18; found 505.10. HPLC retention time: 1.24 minutes (column  
20    A).

**EXAMPLE 63**

- 5 Preparation of 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-(oxazol-5-yl)-4-azaindol-3-yl)-oxoacetyl]piperazine: A mixture of Intermediate 10 (27.6 mg), tosylmethyl isocyanide (12.3 mg) and  $K_2CO_3$  (8.7 mg) in MeOH was heated up to reflux for 10 hours. After cooling down, the mixture was concentrated under reduced pressure and the residue was purified using Shimadzu automated preparative HPLC
- 10 System to provide 1-(benzoyl)-3-(R)-methyl-4-[(5-chloro-7-(oxazol-5-yl)-4-azaindol-3-yl)-oxoacetyl]piperazine (17.7 mg); MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{24}H_{21}ClN_5O_4$ : 478.13; found 478.03. HPLC retention time: 1.48 minutes (column A).

**EXAMPLE 64**

15




I-64

- Step 1: Preparation of I-64, 1-benzoyl-3-(R)-methyl-4-[(7-(2-propynyl)carbonyl)-4-azaindol-3-yl)-oxoacetyl]piperazine: Propynyllithium (21 mg)
- 20 was added to a solution of Example 52 (41 mg) in tetrahydrofuran (5 ml) at  $-78^\circ C$ . The reaction was quenched with methanol at  $-25^\circ C$  in 2 hours. After solvents were removed under vacuum, the residue was carried to the further reactions without any purification.



I-64, 1-benzoyl-3-(R)-methyl-4-[(7-(2-propynyl)carbonyl-4-azaindol-3-yl)-oxoacetyl]piperazine MS  $m/z$ : (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>22</sub>ClN<sub>4</sub>O<sub>4</sub>: 477.13; found 477.17. HPLC retention time: 1.46 minutes (column A).

5 Step 2: Preparation of Example 64:



Example 64

Preparation of Example 64, 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-(3-methyl-pyrazol-5-yl)-4-azaindol-3-yl)-oxoacetyl]piperazine: A mixture of I-64 (crude product from Step 1) and hydrazine (0.22 mL) in EtOAc (2 mL) and water (2 mL) was stirred at room temperature for 24 hours. Then solvents were removed under vacuum, and the residue was purified using Shimadzu automated preparative HPLC System to give 1-benzoyl-3-(R)-methyl-4-[(5-chloro-7-(3-methyl-pyrazol-5-yl)-4-azaindol-3-yl)-oxoacetyl]piperazine (9 mg); MS  $m/z$ : (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>24</sub>ClN<sub>6</sub>O<sub>3</sub>: 491.16; found 491.19. HPLC retention time: 1.42 minutes (column A).


EXAMPLES 65-67

20

The procedure for the preparation of Examples 65-67 is the same as that described previously for the preparation of Intermediate 5a and is as follows: Potassium 7-(4-methoxyphenyl)-4-azaindole-3-glyoxylate, Intermediate 4c (147 mg, 0.44 mmol), an appropriate 1-benzoylpiperazine derivative (0.44 mmol), 3-(diethoxyphosphoryloxy)-1,2,3-benzotriazin-4(3H)-one (DEPBT) (101 mg, 0.44 mol) and Hunig's Base (0.5 mL) were combined in 5 mL of DMF. The mixture was stirred at rt for 8 h. DMF was removed *via* evaporation at reduced pressure and the residue was purified using a Shimadzu automated preparative HPLC System to give the corresponding 1-benzoyl-4-[(7-(4-methoxyphenyl)-4-azaindol-3-yl)-oxoacetyl]-piperazine derivative.


25

30

EXAMPLE 65


5            Example 19, 1-(benzoyl)-4-[(7-(4-methoxy)-4-azaindol-3-yl)-oxoacetyl]piperazine was prepared from potassium 7-(4-methoxyphenyl)-4-azaindole-3-glyoxylate and 1-(benzoyl)piperazine according to the above general procedure. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>25</sub>N<sub>4</sub>O<sub>4</sub>: 469.19; found 469.16. HPLC retention time: 1.26 minutes (column A).

10

EXAMPLE 66

15            Example 66, 1-benzoyl-3-(S)-methyl-4-[(7-(4-methoxy)-4-azaindol-3-yl)-oxoacetyl]piperazine was prepared from potassium 7-(4-methoxyphenyl)-4-azaindole-3-glyoxylate and the corresponding 1-(benzoyl)-3-methylpiperazine according to the above general procedure. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>28</sub>H<sub>27</sub>N<sub>4</sub>O<sub>4</sub>: 483.20; found 483.17. HPLC retention time: 1.30 minutes (column A).

20


EXAMPLE 67

25            Example 67, 1-benzoyl-3-(R)-methyl-4-[(7-(4-methoxyphenyl)-4-azaindol-3-yl)oxoacetyl]piperazine was prepared from potassium 7-(4-methoxyphenyl)-4-azaindole-3-glyoxylate and the corresponding 1-benzoyl-3-methylpiperazine according to the above general procedure. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>28</sub>H<sub>27</sub>N<sub>4</sub>O<sub>4</sub>: 483.20; found 483.16. HPLC retention time: 1.28 minutes (column A).

**EXAMPLES 68-79 and 81**


Examples 68-79 and 81 were prepared according to the same general method as previously described for Examples 16-54.

5

**EXAMPLE 68**


Example 68, was prepared from Intermediate 5b and the 2,4-dimethoxypyrimidin-6-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2,6-dimethoxy-pyrimidin-4-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 8.20 (s, 1H), 8.13 (s, 1H), 7.52 (s, 1H), 7.42 (m, 5H), 4.11 (s, 3H), 4.06 (s, 3H), 4.00 – 3.40 (m, 8H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>27</sub>N<sub>6</sub>O<sub>6</sub>: 531.20; found 531.24. HPLC retention time: 1.54 minutes (column A).

15


**EXAMPLE 69**

Example 69, was prepared from Intermediate 5b and the 6-methoxypyridin-3-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(6-methoxy-pyridin-3-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 8.69 (s, 1H), 8.63 (s, 1H), 8.11 (m, 2H), 7.49 (m, 5H), 7.10 (d, 1H, J = 8.65 Hz), 4.16 (s, 3H), 4.06 (s, 3H), 4.00 – 3.40 (m, 8H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>26</sub>N<sub>5</sub>O<sub>5</sub>: 500.09; found 500.20. HPLC retention time: 1.11 minutes (column A).


25

**EXAMPLE 70**

5            Example 70, was prepared from Intermediate 5b and the 2-diethylamino-thiazol-4-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-diethylamino-thiazol-4-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 8.47 (s, 1H), 7.97 (m, 2H), 7.49 (m, 5H), 4.08 (s, 3H), 3.64 (m, 12H), 1.35 (m, 6H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>28</sub>H<sub>31</sub>N<sub>6</sub>O<sub>4</sub>S: 547.21; found 547.22. HPLC retention time: 1.35  
10   minutes (column A).


**EXAMPLE 71**

15            Example 71, was prepared from Intermediate 5b and the thiazol-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 9.19 (s, 1H), 8.64 (s, 1H), 8.34 (s, 1H), 8.11 (s, 1H), 7.46 (m, 5H), 4.00 (s, 3H), 3.55 (m, 8H). MS *m/z*: (M+H)<sup>+</sup>  
20   Calc'd for C<sub>24</sub>H<sub>22</sub>N<sub>5</sub>O<sub>4</sub>S: 476.14; found 476.17. HPLC retention time: 1.13 minutes (column A).

**EXAMPLE 72**


5            Example 72, was prepared from Intermediate 5b and the 2-dimethylamino-pyrazin-5-yl stannane to provide 1-(benzoyl)-4-[(4-methoxy-7-(2-dimethylamino-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>28</sub>N<sub>7</sub>O<sub>4</sub>: 514.22; found 514.29. HPLC retention time: 1.27 minutes (column A).

10

**EXAMPLE 73**


15            Example 73, was prepared from Intermediate 5b and the furan-2-yl stannane to provide 1-(benzoyl)-4-[(4-methoxy-7-(furan-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>4</sub>O<sub>5</sub>: 459.17; found 459.25. HPLC retention time: 1.15 minutes (column A).

20

**EXAMPLE 74**

Example 74, was prepared from Intermediate 5b and the oxazol-2-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(oxazol-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine.  $^1\text{H}$  NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.19 (s, 1H), 8.64 (s, 1H), 8.34 (s, 1H), 8.11 (s, 1H), 7.46 (m, 5H), 4.00 (s, 3H), 3.55 (m, 8H). MS  $m/z$ : (M+H) $^+$  Calc'd for  $\text{C}_{24}\text{H}_{22}\text{N}_5\text{O}_5$ : 460.16; found 460.23. HPLC retention time: 1.22 minutes (column A).

#### EXAMPLE 75




10

Example 75, was prepared from Intermediate 5b and the 6-aminopyridin-2-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-aminopyridin-6-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS  $m/z$ : (M+H) $^+$  Calc'd for  $\text{C}_{26}\text{H}_{25}\text{N}_6\text{O}_4$ : 485.19; found 485.24. HPLC retention time: 1.15 minutes (column A).

15

#### EXAMPLE 76




20

Example 76, was prepared from Intermediate 5b and the 6-methylpyridin-2-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-methylpyridin-6-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS  $m/z$ : (M+H) $^+$  Calc'd for  $\text{C}_{27}\text{H}_{26}\text{N}_5\text{O}_4$ : 484.20; found 484.22. HPLC retention time: 1.24 minutes (column A).

25


### EXAMPLE 77



Example 77, was prepared from Intermediate 5b and the 6-methoxypyridin-2-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-methoxy-pyridin-6-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>26</sub>N<sub>5</sub>O<sub>5</sub>: 500.19; found 500.23. HPLC retention time: 1.26 minutes (column A).

## 10


### EXAMPLE 78



Example 78, was prepared from Intermediate 5b and the 2-acetylaminothiazol-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-acetylaminothiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>23</sub>N<sub>6</sub>O<sub>5</sub>S: 533.16; found 533.18. HPLC retention time: 1.21 minutes (column A).


20

25

**EXAMPLE 79**

5            Example 79, was prepared from Intermediate 5b and the 2-ethylamino-pyrazin-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-ethylamino-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>28</sub>N<sub>7</sub>O<sub>4</sub>: 514.22; found 514.18. HPLC retention time: 1.31 minutes (column A).


10

**EXAMPLE 88**

15            Example 88, was prepared from Intermediate 5b and the 2-ethyl-thiazol-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-ethyl-thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>26</sub>N<sub>5</sub>O<sub>4</sub>S: 504.17; found 504.32. HPLC retention time: 1.50 minutes (column A).


20



**EXAMPLE 89**


5            Example 89, was prepared from Intermediate 5k and the 2-isobutyl-thiazol-5-yl stannane to provide 1-benzoyl-4-[(7-(2-isobutyl-thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>28</sub>N<sub>5</sub>O<sub>3</sub>S: 502.19; found 502.26. HPLC retention time: 1.56 minutes (column E).

10

**Example 90**


15            Example 90, was prepared from Intermediate 5b and the 2-isobutyl-thiazol-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-isobutyl-thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>28</sub>H<sub>30</sub>N<sub>5</sub>O<sub>4</sub>S: 532.20; found 532.27. HPLC retention time: 1.57 minutes (column E).

20

**Example 91**


5            Example 91, was prepared from Intermediate 5b and the 2-(2-butyl)-thiazol-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-(2-butyl)-thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>28</sub>H<sub>30</sub>N<sub>5</sub>O<sub>4</sub>S: 532.20; found 532.27. HPLC retention time: 1.57 minutes (column E).

10

**Example 92**


15            Example 92, was prepared from Intermediate 5b and the 2-(thiazol-2-yl)-thiazol-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-(thiazol-2-yl)-thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>23</sub>N<sub>6</sub>O<sub>4</sub>S<sub>2</sub>: 559.12; found 559.18. HPLC retention time: 1.55 minutes (column E).

20

**Example 93**

5        Example 93, was prepared from Intermediate 5b and the 2-methylthio-thiazol-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-methylthio-thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>24</sub>N<sub>5</sub>O<sub>4</sub>S<sub>2</sub>: 522.13; found 522.17. HPLC retention time: 1.45 minutes (column E).

10


**Example 95**

15        Example 95, was prepared from Intermediate 5i and the pyrazin-2-yl stannane to provide 1-benzoyl-4-[(4-fluoro-7-(pyrazin-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 9.89 (s, 1H), 8.70 – 8.34 (m, 4H), 7.46 (m, 5H), 3.80 – 3.50 (m, 8H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>24</sub>H<sub>20</sub>FN<sub>6</sub>O<sub>3</sub>: 459.16; found 459.33. HPLC retention time: 1.46 minutes (column G).

20

25


## Example 100



5            Example 100, was prepared from Intermediate 5b and the 2-methylamino-3-methoxy-pyrazin-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-methylamino-3-methoxy-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ8.65 (s, 1H), 8.43 (s, 1H), 7.95 (s, 1H), 7.45 (m, 5H), 4.21 (s, 3H), 4.12 (s, 3H), 3.89 – 3.32 (m, 8H), 3.06 (s, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>28</sub>N<sub>7</sub>O<sub>5</sub>: 530.22; found 530.19. HPLC retention time: 1.31 minutes (column A).

10


## Example 101



15


          Example 101, was prepared from Intermediate 5b and the 2-amino-3-methoxy-pyrazin-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-amino-3-methoxy-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ8.67 (s, 1H), 8.34 (s, 1H), 7.96 (s, 1H), 7.48 (m, 5H), 4.22 (s, 3H), 4.12 (s, 3H), 3.92 – 3.32 (m, 8H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>26</sub>N<sub>7</sub>O<sub>5</sub>: 516.20; found 516.23. HPLC retention time: 1.27 minutes (column A).

20

**Example 102**


5            Example 102, was prepared from Intermediate 51 and the pyrazin-2-yl stannane to provide 1-picolinoyl-4-[(4-methoxy-7-(pyrazin-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 9.59 (s, 1H), 8.79 – 7.51 (m, 8H), 4.13 (s, 3H), 3.95 – 3.34 (m, 8H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>24</sub>H<sub>22</sub>N<sub>7</sub>O<sub>4</sub>: 472.17; found 472.25. HPLC retention time: 1.15 minutes (column A).

10

**Example 103**


15            Example 103, was prepared from Intermediate 51 and the 2-dimethylamino-pyrazin-5-yl stannane to provide 1-picolinoyl-4-[(4-methoxy-7-(2-dimethylamino-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>27</sub>N<sub>8</sub>O<sub>4</sub>: 515.22; found 515.16. HPLC retention time: 1.29 minutes (column A).

## Example 104




5            Example 104, was prepared from Intermediate 5b and the 6-aza-benzofuran-2-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(6-aza-benzofuran-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 8.48 (d, 1H, *J* = 8.5 Hz), 8.36 (s, 1H), 8.30 (s, 1H), 8.02 (s, 1H), 7.64 (d, 1H, *J* = 8.55 Hz), 7.41 (m, 4H), 6.92 (s, 1H), 4.12 (s, 3H), 3.87 – 3.38 (m, 8H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for  
10   C<sub>28</sub>H<sub>24</sub>N<sub>5</sub>O<sub>5</sub>: 510.18; found 510.33. HPLC retention time: 1.33 minutes (column A).

## Example 105




15


Example 105, was prepared from Intermediate 5m and the 2-dimethylamino-pyrazin-5-yl stannane to provide (R)-1-picolinoyl-3-methyl-4-[(7-(2-dimethylamino-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>27</sub>N<sub>8</sub>O<sub>3</sub>: 499.22; found 499.27. HPLC retention time: 1.17 minutes (column A).

**Example 106**

5            Example 106, was prepared from Intermediate 5n and the 2-dimethylamino-pyrazin-5-yl stannane to provide (S)-1-picolinoyl-3-methyl-4-[(7-(2-dimethylamino-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ9.08 – 7.49 (m, 9H), 5.00 – 3.15 (m, 13H), 1.44 – 1.27 (m, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>27</sub>N<sub>8</sub>O<sub>3</sub>: 499.22; found 499.27. HPLC retention time: 1.19 minutes  
10 (column A).


**Example 109**

15            Example 109, was prepared from Intermediate 5m and the thiazol-5-yl stannane to provide (R)-1-picolinoyl-3-methyl-4-[(7-(thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ9.42 – 7.49 (m, 9H), 4.98 – 3.14 (m, 7H), 1.43 – 1.26 (m, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>23</sub>H<sub>21</sub>N<sub>6</sub>O<sub>3</sub>S: 461.14; found 461.28. HPLC retention time: 1.11 minutes (column A).  
20

**Example 110**


5           Example 110, was prepared from Intermediate 5n and the thiazol-5-yl stannane to provide (S)-1-picolinoyl-3-methyl-4-[(7-(thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 9.44 – 7.48 (m, 9H), 4.98 – 3.15 (m, 7H), 1.43 – 1.26 (m, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>23</sub>H<sub>21</sub>N<sub>6</sub>O<sub>3</sub>S: 461.14; found 461.27. HPLC retention time: 1.13 minutes (column A).

10


**Example 111**

15           Example 111, was prepared from Intermediate 5f and the 2-amino-pyrazin-6-yl stannane to provide (R)-1-benzoyl-3-methyl-4-[(7-(2-amino-pyrazin-6-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 8.68 – 7.45 (m, 10H), 4.89 – 3.13 (m, 7H), 1.39 – 0.99 (m, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>24</sub>N<sub>7</sub>O<sub>3</sub>: 470.19; found 470.31. HPLC retention time: 1.30 minutes (column A).




**Example 112**

- 5            Example 112, was prepared from Intermediate 5f and the 2-amino-pyridin-6-yl stannane to provide (R)-1-benzoyl-3-methyl-4-[(7-(2-amino-pyridin-6-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 8.65 – 6.89 (m, 11H), 4.90 – 3.12 (m, 7H), 1.39 – 0.99 (m, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>25</sub>N<sub>6</sub>O<sub>3</sub>: 469.20; found 469.32. HPLC retention time: 1.26 minutes (column A).
- 10.

**Example 113**

- 15            Example 113, was prepared from Intermediate 5f and the 2-amino-pyridin-5-yl stannane to provide (R)-1-benzoyl-3-methyl-4-[(7-(2-amino-pyridin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 8.75 – 7.19 (m, 11H), 4.91 – 3.12 (m, 7H), 1.38 – 1.25 (m, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>25</sub>N<sub>6</sub>O<sub>3</sub>: 469.20; found 469.34. HPLC retention time: 1.05 minutes (column A).


## Example 114



5            Example 114, was prepared from Intermediate 5f and the 5-amino-pyridin-2-yl stannane to provide (R)-1-benzoyl-3-methyl-4-[(7-(5-amino-pyridin-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 8.67 – 7.20 (m, 11H), 4.88 – 3.13 (m, 7H), 1.39 – 1.25 (m, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>25</sub>N<sub>6</sub>O<sub>3</sub>: 469.20; found 469.33. HPLC retention time: 1.22 minutes (column A).


10

## Example 115




15            Example 115, was prepared from Intermediate 5b and the 2-methylamino-pyrazin-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-methylamino-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 8.90 (s, 1H), 8.61 (s, 1H), 8.18 (s, 1H), 7.92 (s, 1H), 7.46 (m, 5H), 4.12 (s, 3H), 3.85 – 3.40 (m, 8H), 3.02 (s, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>26</sub>N<sub>7</sub>O<sub>4</sub>: 500.20; found 500.23. HPLC retention time: 1.24 minutes (column A).

20


**Example 116**

5            Example 116, was prepared from Intermediate 5b and the 2-(2-pyrrolidinon-1-yl)-thiazol-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-((2-pyrrolidinon-1-yl)-thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>28</sub>H<sub>27</sub>N<sub>6</sub>O<sub>5</sub>S<sub>2</sub>: 559.18; found 559.11. HPLC retention time: 1.39 minutes (column E).

10


**Example 117**

15            Example 117, was prepared from Intermediate 5b and the 2-methoxy-pyrimidin-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-methoxypyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>25</sub>N<sub>6</sub>O<sub>5</sub>: 501.19; found 501.12. HPLC retention time: 1.21 minutes (column E).


**Example 118**

5            Example 118, was prepared from Intermediate 5b and the 2-(pyrrol-1-yl)-pyrimidin-5-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(2-(pyrrol-1-yl)-pyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>29</sub>H<sub>26</sub>N<sub>7</sub>O<sub>4</sub>: 536.20; found 536.33. HPLC retention time: 1.44 minutes (column C).


10

**Example 119**


15            Example 119, was prepared from Intermediate 5b and the pyrimidin-4-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(pyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 9.29 (s, 1H), 8.88 (d, 1H, *J* = 5.4 Hz), 8.48 (d, 1H, *J* = 5.25 Hz), 8.26 (s, 1H), 8.18 (s, 1H), 7.43 (m, 5H), 4.13 (s, 3H), 3.85 – 3.47 (m, 8H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>6</sub>O<sub>4</sub>: 471.18; found 471.32. HPLC retention time: 1.35 minutes (column G).

**Example 120**

5            Example 119, was prepared from Intermediate 5b and the pyridazin-3-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(pyridazin-3-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ9.16 (s, 1H), 8.77 (d, 1H, *J* = 8.5 Hz), 8.26 (d, 1H, *J* = 3.05 Hz), 8.18 (s, 1H), 7.68 (m, 1H), 7.43 (m, 5H), 4.13 (s, 3H), 3.85 – 3.47 (m, 8H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>6</sub>O<sub>4</sub>: 471.18; found  
10    471.16. HPLC retention time: 1.35 minutes (column G).


**Example 125**

15            Example 125, was prepared from Intermediate 5i and the pyrimidin-4-yl stannane to provide 1-benzoyl-4-[(4-fluoro-7-(pyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ9.36 (s, 1H), 8.96 (d, 1H, *J* = 5.35 Hz), 8.58 (d, 1H, *J* = 5.10 Hz), 8.43 (s, 1H), 8.38 (s, 1H), 7.43 (m, 5H), 3.85 –  
20    3.47 (m, 8H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>24</sub>H<sub>20</sub>FN<sub>6</sub>O<sub>2</sub>: 459.16; found 459.15. HPLC retention time: 1.48 minutes (column A).

**Example 126**

5            Example 126, was prepared from Intermediate 5i and the oxazol-2-yl stannane to provide (R)-1-benzoyl-3-Methyl-4-[7-(oxazol-2-yl)-4-azaindol-3-yl]-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>24</sub>H<sub>22</sub>N<sub>5</sub>O<sub>4</sub>: 444.17; found 444.25. HPLC retention time: 1.13 minutes (column A).


10

**Example 131**


15            Example 131, was prepared from Intermediate 5p and the thiazol-2-yl stannane to provide 1-benzoyl-4-[7-(thiazol-2-yl)-4-azaindol-3-yl]-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>23</sub>H<sub>20</sub>N<sub>5</sub>O<sub>3</sub>S: 446.13; found 446.04. HPLC retention time: 1.12 minutes (column A).

20

25


**EXAMPLE 80**

- 5 Preparation of **Example 80**, 1-benzoyl-4-[(4-methoxy-7-(2-amino-thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine: A mixture of **Example 78** (9 mg), TFA (3 mL) and water (1 mL) was stirred at 80 °C for 10 hours. After solvent was removed under vacuum, the residue was purified by using silica gel chromatography to afford 1-benzoyl-4-[(4-methoxy-7-(2-amino-thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine (3 mg); MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>24</sub>H<sub>23</sub>N<sub>6</sub>O<sub>3</sub>S: 491.15; found 491.21. HPLC retention time: 1.20 minutes (column A).
- 10

**EXAMPLE 81**


15

- Example 81, was prepared from Intermediate 5b and the furan-3-yl stannane to provide 1-benzoyl-4-[(4-methoxy-7-(furan-3-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>4</sub>O<sub>5</sub>: 459.17; found 459.24.
- 20 HPLC retention time: 1.13 minutes (column A).

**Example 150**

5            Example 150, was prepared from Intermediate 5f and the 5-amino-pyrazin-2-yl stannane to provide (R)-1-benzoyl-3-methyl-4-[(7-(5-amino-pyrazin-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>24</sub>N<sub>7</sub>O<sub>3</sub>: 470.19; found 470.19. HPLC retention time: 1.14 minutes (column G).


10

**Example 153**

15            Example 153, was prepared from Intermediate 5f and the 2-amino-pyrimidin-5-yl stannane to provide (R)-1-benzoyl-3-methyl-4-[(7-(2-amino-pyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>24</sub>N<sub>7</sub>O<sub>3</sub>: 470.19; found 470.22. HPLC retention time: 1.07 minutes (column G).




## Example 147




- 5 Intermediate 5i (16.5 mg, 0.05 mmol) in DMF (1 mL) was treated with *N*-benzoylpiperazine hydrochloride, DEBPT (15 mg, 0.05 mmol) and Hunig's base (34  $\mu$ L, 0.2 mmol) at rt for 18h. The solvent was removed in vacuum and the residue was purified by reverse phase preparative HPLC. The fractions showing the right LC/MS(ES<sup>+</sup>)  $m/z$  (M+H)<sup>+</sup> = 501 were collected, concentrated and purified again  
10 using a preparative TLC (5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to afford the title compound as a white solid. <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  11.2 (s, 1H), 10.0 (s, 1H), 9.21 (s, 1 H), 8.51 (s, 1H), 8.41 (s, 1H), 8.40 (m, 1 H), 8.32 (s, 1H), 7.62 (m, 1H), 7.45 (m, 5H), 3.90-3.50 (bm, 8H).

15

## Example 156



- Example 156, was prepared from Intermediate 5b and the 4,4-dimethyloxazolin-2-yl stannane to provide 1-benzoyl-4-[(7-(4,4-dimethyloxazolin-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS  $m/z$ : (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>28</sub>N<sub>5</sub>O<sub>5</sub>:  
20 490.21; found 490.22. HPLC retention time: 1.20 minutes (column C).

**Example 169**


5            Example 169, was prepared from Intermediate 5b and the 2-(4-pyridinecarboxamido)-thiazol-5-yl stannane to provide 1-benzoyl-4-[(7-(2-(4-pyridinecarboxamido)-thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>30</sub>H<sub>26</sub>N<sub>7</sub>O<sub>5</sub>S: 596.17; found 596.14. HPLC retention time: 1.32 minutes (column C).

10

**EXAMPLES 82-86, 98, 107, 108, 129, 130, 132, 133, 134**

          Examples 82-86, 98, 107, 108, 127, 128, 129, 130, 132, 133 and 134 were prepared according to the general procedure as previously described for Examples 2-14.


15

**EXAMPLE 82**

20


          Example 82, was prepared from Intermediate 5b and thien-2-yl boronic acid to provide 1-benzoyl-4-[(4-methoxy-7-(thiophen-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>4</sub>O<sub>4</sub>S: 475.14; found 475.31. HPLC retention time: 1.14 minutes (column A).

25


**EXAMPLE 83**

5            Example 83, was prepared from Intermediate 5b and thien-2-yl boronic acid to provide 1-benzoyl-4-[(4-methoxy-7-(thiophen-3-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>4</sub>O<sub>4</sub>S: 475.14; found 475.33. HPLC retention time: 1.16 minutes (column A).

10


**EXAMPLE 84**

15            Example 84, was prepared from Intermediate 5b and 5-carbonylthien-2-yl boronic acid to provide 1-benzoyl-4-[(4-methoxy-7-(5-carbonylthiophen-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>23</sub>N<sub>4</sub>O<sub>5</sub>S: 503.14; found 503.23. HPLC retention time: 1.31 minutes (column A).


**EXAMPLE 85**

5            Example 76, was prepared from Intermediate 5b and 5-carboxylfuran-2-yl boronic acid to provide 1-(benzoyl)-4-[(4-methoxy-7-(5-carboxylfuran-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>23</sub>N<sub>4</sub>O<sub>6</sub>: 487.16; found 487.28. HPLC retention time: 1.44 minutes (column A).

10


**EXAMPLE 86**

15            Example 86, was prepared from Intermediate 5d and 4-methylthien-2-yl boronic acid to provide 1-benzoyl-3-(R)-methyl-4-[(7-(4-methylthiophen-2-yl)-4-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>23</sub>N<sub>4</sub>O<sub>3</sub>S: 473.16; found 473.26. HPLC retention time: 1.28 minutes (column A).

**Example 98**


5            Example 98, was prepared from Intermediate 5d and 2-benzofuranyl boronic acid to provide 1-benzoyl-3-(R)-methyl-4-[(7-(benzofuran-2-yl)-4-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 8.24 (s, 1H), 8.09 (s, 1H), 7.70-7.26 (m, 10H), 4.03 (s, 3H), 3.97 – 3.49 (m, 8H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>29</sub>H<sub>25</sub>N<sub>4</sub>O<sub>5</sub>: 509.18; found 509.18. HPLC retention time: 1.50 minutes (column A).

10

**Example 107**


15            Example 107, was prepared from Intermediate 5m and 2-benzofuranyl boronic acid to provide (R)-1-picolinoyl-3-methyl-4-[(7-(benzofuran-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 8.77 – 7.38 (m, 12H), 4.99 – 3.16 (m, 7H), 1.44 – 1.27 (m, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>28</sub>H<sub>24</sub>N<sub>5</sub>O<sub>4</sub>: 494.18; found 494.24. HPLC retention time: 1.35 minutes (column A).

20


**Example 108**

5            Example 108, was prepared from Intermediate 5n and 2-benzofuranyl boronic acid to provide (S)-1-picolinoyl-3-methyl-4-[(7-(benzofuran-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>28</sub>H<sub>24</sub>N<sub>5</sub>O<sub>4</sub>: 494.18; found 494.23. HPLC retention time: 1.37 minutes (column A).

10


**Example 127**

15            Example 127, was prepared from Intermediate 5i and the benzothiophen-2-yl boronic acid to provide (R)-1-benzoyl-3-Methyl-4-[7-(benzothiophen-2-yl)-4-azaindol-3-yl]-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>29</sub>H<sub>25</sub>N<sub>4</sub>O<sub>3</sub>S: 509.16; found 509.21. HPLC retention time: 1.42 minutes (column A).

**Example 128**


5            Example 128, was prepared from Intermediate 5i and the thiophen-2-yl boronic acid to provide (R)-1-benzoyl-3-Methyl-4-[7-(thiophen-2-yl)-4-azaindol-3-yl]-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>4</sub>O<sub>3</sub>S: 459.15; found 459.27. HPLC retention time: 1.22 minutes (column A).

10

**Example 129**

15            Example 129, was prepared from Intermediate 5i and the thiophen-3-yl boronic acid to provide (R)-1-benzoyl-3-Methyl-4-[7-(thiophen-3-yl)-4-azaindol-3-yl]-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>4</sub>O<sub>3</sub>S: 459.15; found 459.34. HPLC retention time: 1.31 minutes (column A).


20

**Example 130**

Example 130, was prepared from Intermediate 5i and the 2,5-dimethyl-isoxazol-4-yl boronic acid to provide (R)-1-benzoyl-3-Methyl-4-[7-(2,5-dimethyl-isoxazol-4-yl)-4-azaindol-3-yl]-oxoacetyl]piperazine. MS  $m/z$ : (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>26</sub>N<sub>5</sub>O<sub>4</sub>: 472.20; found 472.28. HPLC retention time: 1.14 minutes (column A).

5


### Example 132



Example 132, was prepared from Intermediate 5p and the 2-methylcarbonyl-thiophen-5-yl boronic acid to provide 1-benzoyl-4-[7-(2-methylcarbonyl-thiophen-5-yl)-4-azaindol-3-yl]-oxoacetyl]piperazine. MS  $m/z$ : (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>23</sub>N<sub>4</sub>O<sub>4</sub>S: 487.14; found 487.20. HPLC retention time: 1.14 minutes (column A).

15


### Example 133



Example 133, was prepared from Intermediate 5p and the 2-carbonyl-thiophen-5-yl boronic acid to provide 1-benzoyl-4-[7-(2-carbonyl-thiophen-5-yl)-4-azaindol-3-yl]-oxoacetyl]piperazine. MS  $m/z$ : (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>21</sub>N<sub>4</sub>O<sub>4</sub>S: 473.13; found 473.11. HPLC retention time: 1.14 minutes (column A).


25



**Example 134**

5            Example 134, was prepared from Intermediate 5p and the 4-methyl-thiophen-2-yl boronic acid to provide 1-benzoyl-4-[7-(4-methyl-thiophen-2-yl)-4-azaindol-3-yl]-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>4</sub>O<sub>3</sub>S: 459.15; found 459.08. HPLC retention time: 1.26 minutes (column G).


10

**Example 152****Preparation of Example 152:**


To a mixture of **acid intermediate 16** (30 mg, 68  $\mu$ mol), 3-aminopyridine (26 mg, 0.27 mmol) and DMAP (50 mg, 0.41 mmol) was added THF (2 ml), and then EDC (60 mg, 0.31 mmol). The reaction mixture was stirred at ambient temperature for 16 hours. The LC/MS analysis indicated that the major product was the activated ester. The reaction mixture was then added into a DMF (2 ml) solution of 3-aminopyridine (400 mg, 4.25 mmol) and stirred at ambient temperature for 16 hours. After addition of MeOH (4 ml), the reaction mixture was purified by preparative reverse phase HPLC to give the TFA salt of the title compound using the method: Start %B = 30, Final %B = 75, Gradient time = 25 min, Flow Rate = 25 ml/min, Column : YMC C18

Sum 20 x 100mm, Fraction Collection: 10.41 – 11.08 min.  $^1\text{H}$  NMR: ( $\text{DMSO}-d_6$ )  $\delta$  13.04 (s, 1H), 11.17 (s, 1H), 9.17 (s, 1H), 8.53 (s, 1H), 8.35 (m, 3H), 7.44 (b s, 6H), 3.75 – 3.37 (b m, 8H); LC/MS: (ES+)  $m/z$  ( $M+H$ ) $^+$  = 517, 519; HPLC  $R_t$  = 1.653.

5

**Example 143****10 Prep of Example 143:**

To a mixture of intermediate 5q (31 mg, 65  $\mu\text{mol}$ ) and  $\text{Pd}(\text{PPh}_3)_4$  (20 mg, 17  $\mu\text{mol}$ ) was added 1,4-dioxane (1 ml) and ii (30 mg, 78  $\mu\text{mol}$ ). The reaction mixture was heated in a sealed tube at  $145^\circ\text{C}$  for 4 hours. After cooling to ambient temperature, the reaction mixture was added MeOH (4 ml) and then filtered. The filtrate was purified by preparative reverse phase HPLC to give the TFA salt of the title compound using the method: Start %B = 25, Final %B = 90, Gradient time = 20 min, Flow Rate = 25 ml/min, Column : YMC C18 Sum 20 x 100mm, Fraction Collection: 11.14 – 11.92 min.  $^1\text{H}$  NMR: ( $\text{DMSO}-d_6$ )  $\delta$  12.71 (s, 1H), 9.01 (s, 1H), 8.36 (s, 1H), 8.27 (s, 1H), 8.08 (s, 1H), 7.44 (b s, 5H), 7.44 (b s, 2H), 3.75 – 3.37 (b m, 8H); LC/MS: (ES+)  $m/z$  ( $M+H$ ) $^+$  = 490, 492; HPLC  $R_t$  = 2.250.


**Example 149**

25

## Preparation of Example 49:

To a suspension of compound of Example 143 (12 mg, 24  $\mu$ mol) in sulfuric acid (5%, 2 ml), was charged sodium nitrite (22 mg, 0.32 mol) at 0°C. The reaction mixture was stirred at 0°C for 30 minutes and then at ambient temperature for 1 hour. After addition of MeOH (4 ml), the reaction mixture was purified by preparative reverse phase HPLC to give a TFA solvate of title compound using the method: Start %B = 20, Final %B = 85, Gradient time = 15 min, Flow Rate = 25 ml/min, Column : YMC C18 5 $\mu$ m 20 x 100mm, Fraction Collection: 10.67- 11.36 min. <sup>1</sup>H NMR: (DMSO-*d*<sub>6</sub>)  $\delta$  12.62 (s, 1H), 8.45 (s, 1H), 8.35 (s, 1H), 8.29 (s, 1H), 8.18 (s, 1H), 7.44 (b s, 5H), 3.80 – 3.30 (b m, 8H); LC/MS: (ES+) *m/z* (M+H)<sup>+</sup> = 491, 493; HPLC *R*<sub>t</sub> = 2.193.


## Example 144



## Preparation of Example 144:

To a mixture of intermediate 5q (50 mg, 105  $\mu$ mol) and Pd(PPh<sub>3</sub>)<sub>4</sub> (50 mg, 43  $\mu$ mol) was added 1,4-dioxane (1 ml) and iii (77 mg, 210  $\mu$ mol). The reaction mixture was heated in a sealed tube at 145°C for 16 hours. After cooling to ambient temperature, the reaction mixture was added MeOH (4 ml) and then filtered. The filtrate was purified by reverse phase HPLC to give the TFA salt of the title compound of using the method: Start %B = 15, Final %B = 100, Gradient time = 20 min, Flow Rate = 25 ml/min, Column : YMC C18 5 $\mu$ m 20 x 100mm, Fraction Collection: 11.80 – 12.31 min. <sup>1</sup>H NMR: (CD<sub>3</sub>OD)  $\delta$  9.32 (s, 1H), 9.25 (s, 2H), 8.50 (s, 1H), 8.44 (s, 1H), 7.47 (b s, 5H), 4.00 – 3.44 (b m, 8H); LC/MS: (ES+) *m/z* (M+H)<sup>+</sup> = 475, 477; HPLC *R*<sub>t</sub> = 1.833.

## EXAMPLE 87




5 Preparation of **Example 87**, 1-benzoyl-4-[(4-methoxy-7-(2-hydroxycarbonyl-furan-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine: A mixture of the compound of **Example 85** (19 mg), NaClO<sub>2</sub> (9.2 mg) in a mixed solution of CH<sub>3</sub>CN (3 mL) and water (0.5 mL) was stirred at room temperature for 24 hours. After the reaction was quenched by 1N NaOH solution (1 ml), the mixture was extracted with diethyl ether (3 x 10 mL). The aqueous phase was acidified with 1N HCl to give a yellow solid precipitate (5mg) which was the product shown. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>23</sub>N<sub>6</sub>O<sub>7</sub>: 503.16; found 503.19. HPLC retention time: 1.37 minutes (column A).

#### General Procedure of Converting -NH<sub>2</sub> Group to -NHCOR Group

15 Preparation of **Example 99**, 1-(benzoyl)-4-[(4-methoxy-7-(2-acetylaminopyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine: 1-(benzoyl)-4-[(4-methoxy-7-(2-amino-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine (4mg) and acetic anhydride (20mg) were dissolved in pyridine (0.5ml). The reaction was stirred for three hours at room temperature. After reaction was quenched with MeOH (1ml), solvents were concentrated to give a residue which was purified using a Shimadzu automated preparative HPLC System to provide 3.0 mg of the desired compound, 1-(benzoyl)-4-[(4-methoxy-7-(2-acetylaminopyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 9.58 (s, 1H), 9.25 (s, 1H), 8.45 (s, 1H), 8.10 (s, 1H), 7.49 (m, 5H), 4.12 (s, 3H), 3.84 – 3.35 (m, 8H), 2.27 (s, 3H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>27</sub>H<sub>26</sub>N<sub>7</sub>O<sub>5</sub>: 528.20; found 528.22. HPLC retention time: 1.33 minutes (column A).


### General Procedure of Converting -NH<sub>2</sub> Gr up to -OH Group

Preparation of Example 97, 1-(benzoyl)-4-[(4-methoxy-7-(2-hydroxyl-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine: 1-(benzoyl)-4-[(4-methoxy-7-(2-amino-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine (15 mg) and NaNO<sub>2</sub> (10 mg) was added into a H<sub>2</sub>SO<sub>4</sub> solution (0.1ml of concentrated H<sub>2</sub>SO<sub>4</sub> diluted with 0.3 ml of water). The reaction was stirred at room temperature for one hour. Then, the reaction mixture was neutralized with a saturated Na<sub>2</sub>CO<sub>3</sub> solution (10 ml). The solvents were concentrated to give a residue which was purified using a Shimadzu automated preparative HPLC System to provide 4.2mg of the desired compound, 1-(benzoyl)-4-[(4-methoxy-7-(2-hydroxyl-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ8.55 (s, 1H), 8.44 (s, 1H), 8.31 (s, 1H), 8.01 (s, 1H), 7.49 (m, 5H), 4.12 (s, 3H), 3.84 – 3.64 (m, 8H). MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>25</sub>H<sub>23</sub>N<sub>6</sub>O<sub>5</sub>: 487.17; found 487.22. HPLC retention time: 1.13 minutes (column A).




This general procedure is applied to prepare examples 121, 122, 123, 124, 155, 157, and 162.

### Example 121




Example 121, (R)-1-(benzoyl)-3-methyl-4-[(4-methoxy-7-(2-hydroxyl-pyrazin-6-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{25}H_{23}N_6O_4$ : 471.18; found 471.17. HPLC retention time: 1.39 minutes (column G).

5

**Example 121-2**


Example 121-2, (R)-1-(benzoyl)-3-methyl-4-[(4-methoxy-7-(2-hydroxyl-4-oxo-pyrazin-6-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine was isolated during the preparation of Example 121. MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{25}H_{23}N_6O_5$ : 487.17; found 487.17. HPLC retention time: 1.08 minutes (column G).

15

**Example 122**

Example 122, (R)-1-(benzoyl)-3-methyl-4-[(4-methoxy-7-(2-hydroxyl-pyridin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{26}H_{24}N_5O_4$ : 470.18; found 470.17. HPLC retention time: 1.03 minutes (column G).


20

**Example 123**


5        Example 123, (R)-1-(benzoyl)-3-methyl-4-[(4-methoxy-7-(2-hydroxypyridin-6-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>24</sub>N<sub>5</sub>O<sub>4</sub>: 470.18; found 470.14. HPLC retention time: 1.28 minutes (column G).

**Example 124**


10



15        Example 124, (R)-1-(benzoyl)-3-methyl-4-[(4-methoxy-7-(5-hydroxypyridin-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>24</sub>N<sub>5</sub>O<sub>4</sub>: 470.18; found 470.13. HPLC retention time: 1.21 minutes (column G).




## Preparation of Example 138



- 5 Preparation of Example 138, 1-(benzoyl)-4-[(4-methoxy-7-(1-methylpyrazin-2-on-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine: 1-(benzoyl)-4-[(4-methoxy-7-(2-hydroxyl-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine (6mg), MeI (5mg) and K<sub>2</sub>CO<sub>3</sub> (4 mg) were dissolved in acetone (5 ml). The reaction was stirred for four hours at room temperature. After solid was filtered away, the mother liquid was
- 10 concentrated to give a residue which was purified using a Shimadzu automated preparative HPLC System to provide 3.0 mg of the desired compound, 1-(benzoyl)-4-[(4-methoxy-7-(1-methylpyrazin-2-on-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS *m/z*: (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>25</sub>N<sub>6</sub>O<sub>3</sub>; 501.19; found 501.14. HPLC retention time: 1.08 minutes (column G).

15


## Example 139





- Intermediate 4i was dissolved in DMF (2 ml), and to which *N*-benzoyl-(*R*)-methylpiperazine hydrochloride (0.092 g, 0.45 mmol) and 3-(diethoxyphosphoryloxy)-1,2,3-benzotriazin-4(3*H*)-one (DEPBT, 0.180 g, 0.60 mmol) were added, followed by *N,N*-diisopropylethylamine (0.15 ml, 0.87 mmol).
- 5 The reaction mixture was stirred for 2 h at r.t., and then the volatile evaporated under high vacuum. Water was added to the mixture to induce precipitation, and the solids were filtered and dried *in vacuo*. Purification of the crude solid by preparative thin layer chromatography (5% MeOH/ CH<sub>2</sub>Cl<sub>2</sub>), and subsequent washing with ether gave
- the title compound; <sup>1</sup>H NMR: (CDCl<sub>3</sub>) δ 8.78 (s, 1H), 8.32 (s, 1H); 8.28 (s, 1H)
- 10 7.84 (s, 1H), 7.44 (b s, 5H), 6.56 (s, 1H), 5.00–3.00 (b m, 7H), 1.45–1.20 (b s, 3H); LC/MS: (ES<sup>+</sup>) *m/z* (M+H)<sup>+</sup> = 521, 523; HPLC R<sub>t</sub> = 1.677

#### Example 140




15

The title compound was prepared according to general procedures described before (Sn-coupling). <sup>1</sup>H NMR: 8.41(m, 1H); 8.33(m, 1H); 8.16(m, 1H); 7.53(m, 1H); 7.47(bs, 5H); 3.97-3.54(m, 8H). LC/MS: (ES<sup>+</sup>) *m/z*(m+H)<sup>+</sup> = 448, R<sub>t</sub> = 1.28min.

20


#### Example 141



- 25 The title compound was prepared according to general procedures described before (Sn-coupling). <sup>1</sup>H-NMR: 9.71-9.70(m, 1H); 8.80-8.79(m, 1H); 8.66-8.42(m, 2H);

8.41-8.35(m, 2H); 7.99-7.92(m, 1H), 7.69-7.53(m, 1H); 7.48-7.44(m, 1H); 5.05-3.15(m, 8H). LC/MS: (ES<sup>+</sup>) m/z (M+H)<sup>+</sup> = 474. Rt = 1.26min.

### Example 144




5

#### Preparation of Example 144:

To a mixture of intermediate 5q (50 mg, 105  $\mu$ mol) and Pd(PPh<sub>3</sub>)<sub>4</sub> (50 mg, 43  $\mu$ mol) was added 1,4-dioxane (1 ml) and iii (77 mg, 210  $\mu$ mol). The reaction mixture was heated in a sealed tube at 145°C for 16 hours. After cooling to ambient temperature, the reaction mixture was added MeOH (4 ml) and then filtered. The filtrate was purified by reverse phase HPLC to give the TFA salt of the title compound of using the method: Start %B = 15, Final %B = 100, Gradient time = 20 min, Flow Rate = 25 ml/min, Column : YMC C18 5 $\mu$ m 20 x 100mm, Fraction Collection: 11.80 – 12.31 min. <sup>1</sup>H NMR: (CD<sub>3</sub>OD)  $\delta$  9.32 (s, 1H), 9.25 (s, 2H), 8.50 (s, 1H), 8.44 (s, 1H), 7.47 (b s, 5H), 4.00 – 3.44 (b m, 8H); LC/MS: (ES<sup>+</sup>) m/z (M+H)<sup>+</sup> = 475, 477; HPLC R<sub>t</sub> = 1.833.


20

### Example 145



The title compound was prepared following the procedure described before for example 146 and intermediate 4k. <sup>1</sup>H NMR: 8.35-8.33(m, 2H); 8.11(s, 1H); 7.89(s, 1H); 7.43(bs, 5H); 3.89-3.49(m, 8H). LC/MS: (ES<sup>+</sup>) m/z (M+H)<sup>+</sup> = 448. Rt = 1.18min.


## Example 146



5 Intermediate 4m (0.26 mmol) was dissolved in DMF (1 mL) and treated with *N*-benzoylpiperazine hydrochloride (59 mg, 0.26 mmol), DEBPT (79 mg, 0.26 mmol) and Hunig's base (90  $\mu$ L, 0.52 mmol) and the reaction mixture was stirred at rt for 18h. The solvent was removed in vacuum and the residue was purified by reverse phase preparative HPLC. The fractions showing the right LC/MS:(ES<sup>+</sup>) m/z (M+H)<sup>+</sup>  
10 = 449 were collected, concentrated and purified again using a preparative TLC (5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to afford the title compound as a white solid. <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  10.7 (s, 1H), 9.00 (s, 1H), 8.54 (s, 1H), 8.39 (s, 1H), 7.45 (m, 5H), 3.9-3.5 (bm, 8H).


15

## Example 148



20 The title compound was prepared from intermediate 4n using the same coupling conditions described for the last step of the preparation of intermediate 5i. <sup>1</sup>H NMR: 8.82(m, 1H); 8.48-8.45(m, 1H); 8.37-8.33(m, 1H); 8.26-8.23(m, 1H); 7.47(bs, 5H); 3.97-3.54(m, 8H). LC/MS: (ES<sup>+</sup>) m/z(m+H)<sup>+</sup> = 447 Rt = 0.94min.


## Example 151



5            Example 151, was prepared from Intermediate 51 and the thiazol-5-yl stannane to provide 1-picolinoyl-4-[(4-methoxy-7-(thiazol-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{23}H_{21}N_6O_4S$ : 477.13; found 477.13. HPLC retention time: 0.94 minutes (column G).

10


## Example 154



15            The title compound was prepared according to general procedures described before (Sn-coupling).  $^1H$ -NMR: 9.23-9.22 (m, 1H); 8.83-8.81(m, 1H); 8.43 (m, 1H); 8.36 (m, 1H); 7.75-7.73 (m,1H), 7.44 (bs, 5H); 3.85-3.49 (m, 8H). LC/MS:  $(ES^+)$   $m/z$   $(M+H)^+ = 459$ .  $R_t = 1.39$ min.


20

## Example 155




Example 155, 1-(benzoyl)-4-[(4-methoxy-7-(2-hydroxyl-pyrazin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{25}H_{23}N_6O_5$ : 487.17; found 487.14. HPLC retention time: 1.30 minutes (column G).

5

**Example 157**

Example 157, (R)-1-(benzoyl)-3-methyl-4-[(4-methoxy-7-(5-hydroxyl-pyrazin-2-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{25}H_{23}N_6O_4$ : 471.18; found 471.16. HPLC retention time: 1.09 minutes (column G).


**Example 161**

**A**  
 $C_{28}H_{23}N_7O_3$   
 Exact Mass: 505.19  
 Mol. Wt.: 505.53

15 **Procedure as usual to yield A:**  $^1H$  NMR (500 MHz, DMSO)  $\delta$  9.67 (s, 1H), 8.81 (s, 1H), 8.72 (d,  $J$  = 5.4 Hz, 1H), 8.25 (d,  $J$  = 6.1 Hz, 1H), 8.00 (dd,  $J$  = 8.2, 1.8 Hz, 1H), 7.68 (dd,  $J$  = 8.2, 7.4 Hz, 2H), 7.60 (tt,  $J$  = 7.4, 1.8 Hz, 2H), 7.48 (br s, 5H), 4.04-3.46 (m, 8H). MS  $m/z$ :  $(M+H)^+$  calcd for  $C_{28}H_{24}N_7O_3$ : 506.19; found 506.15. HPLC retention time: 1.21 minutes (XTERRA C18 S7 3.0 x 50 mm)).

20


## Example 162




- 5        Example 162, (R)-1-(benzoyl)-3-methyl-4-[(4-methoxy-7-(2-hydroxyl-pyrimidin-5-yl)-6-azaindol-3-yl)-oxoacetyl]piperazine. MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{25}H_{23}N_6O_4$ : 471.18; found 471.13. HPLC retention time: 0.95 minutes (column G).

## Example 163

10




- To a solution of intermediate 5q (50 mg, 0.11 mmol) in DMF (1 ml) was added CuCN (30 mg, 0.335 mmol). The reaction mixture was heated at 170°C for 30 min.
- 15    After cooling to ambient temperature, the reaction mixture was diluted with MeOH (15 ml), filtered under gravity, and the filtrate evaporated *in vacuo* to afford a brownish residue which is a cyanointermediate. To the residue in DMF (1 ml) was added sodium azide (61 mg, 0.95 mmol) and ammonium chloride (50 mg, 0.95 mmol). The mixture was heated at 90°C for one hour. The reaction mixture was then
- 20    diluted with MeOH (4 ml), filtered, and the filtrate purified by preparative reverse phase HPLC using the method: Start %B = 20, Final %B = 80, Gradient time = 15 min, Flow Rate = 40 ml/min, Column : XTERRA C18 5  $\mu$ m 30 x 100 mm, Fraction Collection: 11.26 – 11.71 min. The material was homogenous by  $^1H$  NMR and HPLC, although the mass spectrum indicated an extra peak at  $(M+H)^+ = 431$ ;  $^1H$
- 25    NMR: ( $CD_3OD$ ) 8.41 (s, 1H), 8.12 (s, 1H), 7.47 (b s, 5H), 3.97 – 3.47 (b m, 8H); LC/MS: (ES+)  $m/z$   $(M+H)^+ = 465, 467$ ; HPLC  $R_t = 1.937$

**Example 164**


5            Example 164, was prepared from Intermediate 5a and the 4-hydroxycarbonylphenyl boronic acid to provide 1-benzoyl-4-[7-(4-hydroxycarbonylphenyl)-4-azaindol-3-yl]-oxoacetyl]piperazine. MS  $m/z$ :  $(M+H)^+$  Calc'd for  $C_{28}H_{25}N_4O_5$ : 497.18; found 497.22. HPLC retention time: 1.20 minutes (column C).

10

**Example 165**

Compound of Example 165 was prepared in a similar manner to compound of  
 15    Example 143 starting with intermediate 5r, but at 125°C for 22 hours and purification by preparative thin layer chromatography (4% MeOH/ $CH_2Cl_2$ ).  $^1H$  NMR: ( $CDCl_3$ )  $\delta$  11.85 (s, 1H), 9.91 (d,  $J = 1.6$  Hz, 1H), 8.70 (d,  $J = 2.6$  Hz, 1H), 8.65 (dd,  $J = 1.6, 2.6$  Hz, 1H), 8.52 (s, 1H), 8.35 (d,  $J = 3.1$  Hz, 1H), 3.73 (b m, 2H), 3.56 (b m, 4H), 3.53 (b m, 2H), 1.48 (s, 9H); LC/MS: (ES+)  $m/z$   $(M+H)^+ = 471, 473$ ; HPLC  $R_t = 1.690$ .


## Example 167



5 Intermediate 4m (0.098 mmol) was dissolved in DMF (1 mL) and treated with *N*-[5-(2-Bromofuroyl)]piperazine hydrochloride (30 mg, 0.098 mmol), DEBPT (60 mg, 0.19 mmol ) and Hunig's base (70  $\mu$ L, 0.19 mmol) and the reaction mixture was stirred at rt for 18h. The solvent was removed in vacuum and the residue was purified by reverse phase preparative HPLC. The fractions showing the right LC/MS:(ES<sup>+</sup>)  
 10  $m/z$  (M+H)<sup>+</sup> = 518,520 were collected, concentrated and purified again using a preparative TLC (5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to afford the title compound as a white solid. <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  10.7 (s, 1H), 9.00 (s, 1H), 8.54 (s, 1 H), 8.40 (s, 1H), 7.06 (d, *J*=3.4 Hz, 1H), 6.46 06 (d, *J*=3.4 Hz, 1H), 3.90-3.66 (bm, 8H).

15

## Example 168



Example 168, 1-benzoyl-3-(R)-methyl-4-[(7-(2-thienylcarbonyl)-4-azaindol-3-yl)-oxoacetyl]piperazine, was prepared from a reaction 1-benzoyl-3-(R)-methyl-4-[(7-(methoxymethylamino)carbonyl)-4-azaindol-3-yl)-oxoacetyl]piperazine and 2-thienyl lithium by using the same procedure for the preapartion of I-64, 1-benzoyl-3-(R)-methyl-4-[(7-(2-propynyl)carbonyl)-4-azaindol-3-yl)-oxoacetyl]piperazine. MS  
 20  $m/z$ : (M+H)<sup>+</sup> Calc'd for C<sub>26</sub>H<sub>23</sub>N<sub>4</sub>O<sub>4</sub>S: 487.14; found 487.11. HPLC retention time:  
 25 1.31 minutes (column A).  
 6 and hereafter, the following definitions apply.



**Biology**

- “ $\mu$ M” means micromolar;
- “mL” means milliliter;
- 5 • “ $\mu$ l” means microliter;
- “mg” means milligram;

The materials and experimental procedures used to obtain the results reported in Tables 1-5 are described below.

10

**Cells:**

- Virus production-Human embryonic Kidney cell line, 293, propagated in Dulbecco's Modified Eagle Medium (Life Technologies, Gaithersburg, MD) containing 10% fetal Bovine serum (FBS, Sigma, St. Louis, MO).
- 15 • Virus infection- Human epithelial cell line, HeLa, expressing the HIV-1 receptors CD4 and CCR5 was propagated in Dulbecco's Modified Eagle Medium (Life Technologies, Gaithersburg, MD) containing 10% fetal Bovine serum (FBS, Sigma, St. Louis, MO) and supplemented with 0.2 mg/mL Geneticin (Life Technologies, Gaithersburg, MD) and 0.4 mg/mL Zeocin (Invitrogen, Carlsbad, CA).
- 20

**Virus**-Single-round infectious reporter virus was produced by co-transfecting human embryonic Kidney 293 cells with an HIV-1 envelope DNA expression vector and a proviral cDNA containing an envelope deletion mutation and the luciferase reporter gene inserted in place of HIV-1 nef sequences (Chen et al, Ref. 41). Transfections were performed using lipofectAMINE PLUS reagent as described by the manufacturer (Life Technologies, Gaithersburg, MD).

25

**Experiment**

30

1. Compound was added to HeLa CD4 CCR5 cells plated in 96 well plates at a cell density of  $5 \times 10^4$  cells per well in 100  $\mu$ l Dulbecco's Modified Eagle Medium containing 10 % fetal Bovine serum at a concentration of  $<20 \mu$ M.

35

2. 100  $\mu$ l of single-round infectious reporter virus in Dulbecco's Modified Eagle Medium was then added to the plated cells and compound at an approximate multiplicity of infection (MOI) of 0.01, resulting in a final volume of 200  $\mu$ l per well and a final compound concentration of  $<10 \mu$ M.
- 5 3. Samples were harvested 72 h after infection.
4. Viral infection was monitored by measuring luciferase expression from viral DNA in the infected cells using a luciferase reporter gene assay kit (Roche Molecular Biochemicals, Indianapolis, IN). Infected cell supernatants were  
10 removed and 50  $\mu$ l of Dulbecco's Modified Eagle Medium (without phenol red) and 50  $\mu$ l of luciferase assay reagent reconstituted as described by the manufacturer (Roche Molecular Biochemicals, Indianapolis, IN) was added per well. Luciferase activity was then quantified by measuring luminescence using a  
15 Wallac microbeta scintillation counter.
5. The percent inhibition for each compound was calculated by quantifying the level of luciferase expression in cells infected in the presence of each compound as a percentage of that observed for cells infected in the absence of compound and  
20 subtracting such a determined value from 100.
6. An  $EC_{50}$  provides a method for comparing the antiviral potency of the compounds of this invention. The effective concentration for fifty percent inhibition ( $EC_{50}$ ) was calculated with the Microsoft Excel Xlfit curve fitting software. For each  
25 compound, curves were generated from percent inhibition calculated at 10 different concentrations by using a four parameter logistic model (model 205). The  $EC_{50}$  data for the compounds is shown in Tables 2-4. Table 1 is the key for the data in Tables 2-4.

**Results**Table 1. Biological Data Key for EC<sub>50</sub>s

Compounds* with EC <sub>50</sub> s >5μM	Compounds with EC <sub>50</sub> s >1 μM but <5μM	Compounds with EC <sub>50</sub> >50nM but not yet tested at higher concentrations	Compounds with EC <sub>50</sub> < 1 μM
Group C	Group B	Group A'	Group A

\*Some of these compounds may have been tested at a concentration lower than their EC<sub>50</sub> but showed some ability to cause inhibition and thus should be evaluated at a higher concentration to determine the exact EC<sub>50</sub>.

In Tables 2-5, X<sub>2</sub>, X<sub>4</sub> etc. indicates the point of attachment.






















































































10 **Table 2****Examples**























Table Entry (Example Number.)	R2	R3	R4	R9	A	EC <sub>50</sub> Group from Table 1
1 (Example 1)	H	H		CH <sub>3</sub>		A
2 (Example 2)	H	H		CH <sub>3</sub>		A
4 (Example 4)	H	H		H		A

5 (Example 5)	H	H		CH <sub>3</sub>		A
6 (Example 6)	H	H		CH <sub>3</sub>		A
7 (Example 7)	H	H		H		A
8 (Example 8)	H	H		H		A
9 (Example 9)	H	H		CH <sub>3</sub>		A
10 (Example 16)	H	H		CH <sub>3</sub>		A
11 (Example 17)	H	H		H		A
12 (Example 18)	H	H		CH <sub>3</sub>		A
13 (Example 10)	H	H		H		A
14 (Example 19)	H	H		H		A
15 (Example 11)	H	H		H		A'
16 (Example 20)	H	H		CH <sub>3</sub>		A



















17 (Example 21)	H	H		H		A
18 (Example 22)	OMe	H		H		A
19 (Example 23)	OMe	H		H		A
20 (Example 24)	OMe	H		H		A
21 (Example 25)	OMe	H		H		A
22 (Example 26)	OMe	H		H		A
23 (Example 27)	OMe	H		H		A
24 (Example 28)	OMe	H		H		A
25 (Example 29)	F	H		H		A
26 (Example 30)	F	H		H		A



















27 (Example 15)	OMe	H		H		A
28 (Example 32)	OMe	H		H		A
29 (Example 33)	H	H		Me		A
30 (Example 34)	H	H		H		A
31 (Example 35)	OMe	H		H		A
32 (Example 36)	OMe	H		H		A
33 (Example 37)	F	H		Me		A
34 (Example 38)	F	H		H		A
35 (Example 39)	OMe	H		H		A
36 (Example 40)	OMe	H		H		A
37 (Example 41)	F	H		Me		A



















38 (Example 42)	F	H		H		A
41 (Example 45)	OMe	H		H		A
42 (Example 46)	OMe	H		H		A
43 (Example 47)	OMe	H		H		A
45 (Example 49)	OMe	H		H		A
46 (Example 13)	OMe	H		H		A
47 (Example 55)	OMe	H		H		A
48 (Example 50)	OMe	H		H		A
49 (Example 14)	OMe	H		H		A





















50 (Example 68)	OMe	H		H		A
51 (Example 69)	OMe	H		H		A
52 (Example 70)	OMe	H		H		A
53 (Example 71)	OMe	H		H		A
54 (Example 72)	OMe	H		H		A
55 (Example 82)	OMe	H		H		A
56 (Example 73)	OMe	H		H		A
57 (Example 83)	OMe	H		H		A
58 (Example 84)	OMe	H		H		A
59 (Example 85)	OMe	H		H		A
60 (Example 74)	OMe	H		H		A













































61 (Example 75)	OMe	H		H		A
62 (Example 76)	OMe	H		H		A
63 (Example 77)	OMe	H		H		A
64 (Example 78)	OMe	H		H		A
65 (Example 80)	OMe	H		H		A
66 (Example 79)	OMe	H		H		A
67 (Example 87)	OMe	H		H		A
68 (Example 81)	OMe	H		H		A
69 (Example 88)	OMe	H		H		A





















70 (Example 89)	H	H		H		A
71 (Example 90)	OMe	H		H		A
72 (Example 91)	OMe	H		H		A
72 (Example 92)	OMe	H		H		A
73 (Example 93)	OMe	H		H		A
74 (Example 94)	OMe	H		H		A
75 (Example 95)	F	H		H		A
76 (Example 96)	Cl	H		H		A
77 (Example 97)	OMe	H		H		A

78 (Example 98)	OMe	H		H		A
79 (Example 99)	OMe	H		H		A
80 (Example 100)	OMe	H		H		A
81 (Example 101)	OMe	H		H		A
82 (Example 102)	OMe	H		H		A
83 (Example 103)	OMe	H		H		A
84 (Example 104)	OMe	H		H		A
85 (Example 105)	H	H		(R)-Me		A
86 (Example 106)	H	H		(S)-Me		A

87 (Example 107)	H	H		(R)-Me		A
88 (Example 108)	H	H		(S)-Me		A
89 (Example 109)	H	H		(R)-Me		A
90 (Example 110)	H	H		(S)-Me		A
91 (Example 111)	H	H		(R)-Me		A
92 (Example 112)	H	H		(R)-Me		A
93 (Example 113)	H	H		(R)-Me		A
94 (Example 114)	H	H		(R)-Me		A
95 (Example 115)	OMe	H		H		A
96 (Example 116)	OMe	H		H		A

97 (Example 117)	OMe	H		H		A
98 (Example 118)	OMe	H		H		A
99 (Example 119)	OMe	H		H		
100 (Example 120)	OMe	H		H		
101 (Example 121)	H	H		(R)-Me		
102 (Example 121-2)	H	H		(R)-Me		
103 (Example 122)	H	H		(R)-Me		
104 (Example 123)	H	H		(R)-Me		
105 (Example 124)	H	H		(R)-Me		
106 (Example 125)	F	H		H		

107 (Example 138)	OMe	H		H		
108 (Example 139)	Br	H		(R)-Me		
109 (Example 140)	F	H		H		
110 (Example 141)	F	H		(R)-Me		
111						
112 (Example 143)	Cl	H		H		
113 (Example 144)	Cl	H		H		
114 (Example 145)	F	H		H		
115 (Example 146)	F	H		H		
116 (Example 147)	F	H		H		
117 (Example 148)	F	H		H		
118 (Example 149)	Cl	H		H		

119 (Example 150)	H	H		(R)-Me		
120 (Example 151)	OMe	H		H		
121 (Example 152)	Cl	H		H		
122 (Example 153)	H	H		(R)-Me		
123 (Example 154)	F	H		H		
124 (Example 155)	OMe	H		H		
125 (Example 156)	OMe	H		H		
126 (Example 157)	H	H		(R)-Me		
127 (Example 165)	Cl	H		H		
128 (Example 166)	F	H		H		












129 (Example 167)	F	H		H		
130 (Example 162)	H	H		(R)-Me		
131 (Example 163)	Cl	H		H		
132 (Example 164)	H	H		(R)Me		
133 (Example 169)	OMe	H		H		

Table 3



Example 56








































Table Entry (Example number)	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>9</sub>	A	EC <sub>50</sub> Group from Table 1
1 (Example 56)	H	H		CH <sub>3</sub>		B





















Table 4



5

Table Entry (Example No.)	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>9</sub>	A	EC <sub>50</sub> Group from Table1
1 (Example 65)	H	H		H		A
2 (Example 66)	H	H		(S)-CH <sub>3</sub>		A
3 (Example 67)	H	H		(R)-Me		A
4 Example (57)	H	H		(R)-Me		A
6 (Example 64)	Cl	H		(R)-Me		A
7 (Example 58)	Cl	H		(R)-Me		A

8 (Example 60)	Cl	H		(R)-Me		A
9 (Example 61)	Cl	H		(R)-Me		A
10 (Example 62)	Cl	H		(R)-Me		A
11 (Example 63)	Cl	H		(R)-Me		A
12 (Example 59)	Cl	H		(R)-Me		A
13 (Example 51)	H	H		(R)-Me		A
14 (Example 52)	H	H		(R)-Me		A
15 (Example 53)	H	H		(R)-Me		A
16 (Example 54)	H	H		(R)-Me		A
17 (Example 86)	H	H		(R)-Me		A
18 (Example 126)	H	H		(R)-Me		A


19 (Example 127)	H	H		(R)-Me		A
20 (Example 128)	H	H		(R)-Me		A
21 (Example 129)	H	H		(R)-Me		A
22 (Example 130)	H	H		(R)-Me		A
23 (Example 131)	H	H		H		A
24 (Example 132)	H	H		H		A
25 (Example 133)	H	H		H		A
26 (Example 134)	H	H		H		A
27 (Example 135)	H	H		(R)-Me		A
28 (Example 136)	H	H		(R)-Me		A

29 (Example 137)	H	H		(R)-Me		A
30 (Example 158)	H	H		H		A
31 (Example 159)	H	H		H		A
32 (Example 160)	H	H		H		A
33 (Example 161)	H	H		H		A
37 (Example 168)	H	H		H		A









The 5-aza inhibitors shown in Table 5 can be prepared from intermediates 1a or 2s or the corresponding 7-desbromo-7-chloro intermediates which are prepared analogously and the methods herein or by using other methods described herein.

5

Table 5




5-aza inhibitors

Table Entry (Example Number.)	R2	R3	R4	R9	A
1	MeO	H		H	
2	MeO	H		H	
3	MeO	H		H	
4	MeO	H		H	





The compounds in Table 2a exemplify some of the many additional inhibitors which could be prepared by using methodology contained herein or exemplified in the preparation of the compounds in Table 2.

















Table 2a































Additional Inhibitors

10



















	R2	R3	R4	R9	A
	OMe	H		H	
	F	H		H	



















	Cl	H		H	
	F	H		H	
	OMe	H		H	
	OMe	H		H	
	F	H		H	
	F	H		H	
	F	H		H	
	OMe	H		H	



















	F	H		H	
	Cl	H		H	
	Cl	H		(R)-Me	
	F	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	

















	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	



















































	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	H	H		(R)-Me	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	



























	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	





























	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	F	H		H	
	F	H		H	
	F	H		H	
	OMe	H		H	

	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	





















	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	

















	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	

















	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	





















	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	















	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	

	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	


	OM	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	

	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	



















	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	
	OMe	H		H	

The inhibitors in Table 4a could be prepared using analogous procedures which were demonstrated to prepare the examples in Table 4.

## 5 Table 4a



Other inhibitors

Table Entry	R2	R3	R4	R9	A
1	H	H		H <sub>3</sub>	
2	H	H		H	
3	H	H		H	
4	H	H		H	
5	H	H		H	
6	H	H		H	
7	H	H		H	
8	H	H		H	
9	H	H		H	

The compounds of Table 6 below were all found to be very potent in the assay described above using % inhibition as a criteria. In Table 6,  $X_2$ ,  $X_4$  etc. indicates the point of attachment. The vast majority of the compounds exhibited greater than 98% inhibition at a concentration of 10 $\mu$ M. The data at 10 $\mu$ M was calculated in the following manner:


Method for extrapolating % inhibition at 10 $\mu$ M

The compounds of Table 6 below were all found to be very potent in the assay described above using % inhibition as a criteria. In Table 5,  $X_2$ ,  $X_4$  etc. indicates the point of attachment. The vast majority of the compounds exhibited greater than 98% inhibition at a concentration of 10 $\mu$ M. The data at 10 $\mu$ M was calculated in the following manner:

Method for extrapolating % inhibition at 10 $\mu$ M

The data in Table 6 was obtained using the general procedures above and by the following methods. Data is not reported for all compounds since data for all the compounds is reported by the alternate method in Table 2. The percent inhibition for each compound was calculated by quantifying the level of luciferase expression in cells infected in the presence of compound as a percentage of that observed for cells infected in the absence of compound and subtracting such a determined value from 100. For compounds tested at concentrations less than 10  $\mu$ M, the percent inhibition at 10  $\mu$ M was determined by extrapolation using the XLfit curve fitting feature of the Microsoft Excel spreadsheet software. Curves were obtained from 10 data points (% inhibition determined at 10 concentrations of compound) by using a four parameter logistic model (XLfit model 205:  $y = A + ((B-A)/(1+((C/x)^D)))$ ), where, A = minimum y, B = maximum y, C = logEC<sub>50</sub>, D = slope factor, and x and y are known data values. Extrapolations were performed with the A and B parameters unlocked.

Thus the compounds of this invention are all potent antiviral inhibitors based on this assay.

**Table 6**

5

Compound #	Average % inhibition at 10 $\mu$ M
Intermediate 8	85%
Example 1	56%

The compounds of the present invention may be administered orally,  
parenterally (including subcutaneous injections, intravenous, intramuscular,  
10 intrasternal injection or infusion techniques), by inhalation spray, or rectally, in  
dosage unit formulations containing conventional non-toxic pharmaceutically-  
acceptable carriers, adjuvants and vehicles.

Thus, in accordance with the present invention there is further provided a  
method of treating and a pharmaceutical composition for treating viral infections such  
15 as HIV infection and AIDS. The treatment involves administering to a patient in  
need of such treatment a pharmaceutical composition comprising a pharmaceutical  
carrier and a therapeutically-effective amount of a compound of the present  
invention.



The pharmaceutical composition may be in the form of orally-administrable suspensions or tablets; nasal sprays, sterile injectable preparations, for example, as sterile injectable aqueous or oleagenous suspensions or suppositories.

When administered orally as a suspension, these compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may contain microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners/flavoring agents known in the art. As immediate release tablets, these compositions may contain microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants known in the art.

The injectable solutions or suspensions may be formulated according to known art, using suitable non-toxic, parenterally-acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution or isotonic sodium chloride solution, or suitable dispersing or wetting and suspending agents, such as sterile, bland, fixed oils, including synthetic mono- or diglycerides, and fatty acids, including oleic acid.


The compounds of this invention can be administered orally to humans in a dosage range of 1 to 100 mg/kg body weight in divided doses. One preferred dosage range is 1 to 10 mg/kg body weight orally in divided doses. Another preferred dosage range is 1 to 20 mg/kg body weight orally in divided doses. It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

Scheme 41a depicts methodology for converting a carboxylic acid to an alkynyl ketone. The alkynyl ketone intermediates can then be converted to pyrazoles or isoxazoles upon reaction with hydrazines or hydroxyl amines, respectively.

CLAIMS

What is claimed is:


1. A compound of Formula I, including pharmaceutically acceptable salts  
5 thereof,



wherein:

10

Q is selected from the group consisting of:



- 15  $R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$ , are independently selected from the group consisting of hydrogen, halogen, cyano, nitro,  $\text{COOR}^8$ ,  $\text{XR}^{57}$ ,  $\text{C(O)R}^{57}$ ,  $\text{C(O)NR}^{55}\text{R}^{56}$ , B, D, and E with the proviso that at least one of  $R^1$ - $R^4$  is selected from B or E; wherein - - represents a carbon-carbon bond or does not exist;


- 20 m is 1 or 2;

$R^5$  is hydrogen or  $(\text{CH}_2)_n\text{R}^{44}$  wherein n is 0-6;

$R^6$  is O or does not exist;

25

- A is selected from the group consisting of C<sub>1-6</sub>alkoxy, aryl and heteroaryl; in which said aryl is phenyl or naphthyl; said heteroaryl is selected from the group consisting of pyridinyl, pyrimidinyl, pyrazinyl, triazinyl, furanyl, thienyl, pyrrolyl, imidazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, quinolinyl, isoquinolinyl, benzofuranyl, benzothienyl, benzoimidazolyl and benzothiazolyl; and said aryl or heteroaryl is optionally substituted with one or two of the same or different members selected from the group consisting of amino, nitro, cyano, C<sub>1-6</sub>alkoxy, -C(O)NH<sub>2</sub>, C<sub>1-6</sub>alkyl, -NHC(O)CH<sub>3</sub>, halogen and trifluoromethyl; -W- is



- B is selected from the group consisting of -C(=NR<sup>46</sup>)(R<sup>47</sup>), C(O)NR<sup>40</sup>R<sup>41</sup>, aryl, heteroaryl, heteroalicyclic, S(O)<sub>q</sub>R<sup>8</sup>, P(O)(R<sup>8</sup>)<sub>q</sub>(OR<sup>8</sup>)<sub>2-q</sub>, P(S)(R<sup>8</sup>)<sub>q</sub>(OR<sup>8</sup>)<sub>2-q</sub>, C(O)R<sup>7</sup>, XR<sup>8</sup>, (C<sub>1-6</sub>)alkylNR<sup>40</sup>R<sup>41</sup>, and (C<sub>1-6</sub>)alkylCOOR<sup>8</sup>; wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to three same or different substituents selected from the group F;

q is 0, 1, or 2;

- 20 D is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, and (C<sub>2-6</sub>)alkynyl are optionally substituted with one to three same or different halogens or from one to three same or different substituents selected from the group F;

- 25 E is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, and (C<sub>2-6</sub>)alkynyl are substituted with B;

- 30 F is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, (C<sub>1-6</sub>)alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, thiohydroxy, (C<sub>1-6</sub>)thioalkoxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, cyano, halogen, nitro, carbonyl, thiocarbonyl, benzyl, O-

- thiocarbamyl, N-thiocarbamyl, C-thioamido,  $-\text{NR}^{42}\text{C}(\text{O})-(\text{C}_{1-6})\text{alkyl}$ ,  $-\text{NR}^{42}\text{C}(\text{O})-(\text{C}_{3-6})\text{cycloalkyl}$ ,  $-\text{NR}^{42}\text{C}(\text{O})\text{-aryl}$ ,  $-\text{NR}^{42}\text{C}(\text{O})\text{-heteroaryl}$ ,  $-\text{NR}^{42}\text{C}(\text{O})\text{-heteroalicyclic}$ , a cyclic N-amido,  $-\text{NR}^{42}\text{S}(\text{O})_2-(\text{C}_{1-6})\text{alkyl}$ ,  $-\text{NR}^{42}\text{S}(\text{O})_2-(\text{C}_{3-6})\text{cycloalkyl}$ ,  $-\text{NR}^{42}\text{S}(\text{O})_2\text{-aryl}$ ,  $-\text{NR}^{42}\text{S}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NR}^{42}\text{S}(\text{O})_2\text{-heteroalicyclic}$ , O-carboxy, sulfinyl, sulfonyl,  $-\text{S}(\text{O})_2\text{NR}^{42}\text{R}^{43}$ , phosphonyl,  $\text{NR}^{42}\text{R}^{43}$ ,  $(\text{C}_{1-6})\text{alkylC}(\text{O})\text{NR}^{42}\text{R}^{43}$ ,  $\text{C}(\text{O})\text{NR}^{42}\text{R}^{43}$ ,  $\text{NHC}(\text{O})\text{NR}^{42}\text{R}^{43}$ ,  $\text{OC}(\text{O})\text{NR}^{42}\text{R}^{43}$ ,  $\text{NHC}(\text{O})\text{OR}^{54}$ ,  $(\text{C}_{1-6})\text{alkylNR}^{42}\text{R}^{43}$ ,  $\text{COOR}^{54}$ , and  $(\text{C}_{1-6})\text{alkylCOOR}^{54}$ ; wherein said  $(\text{C}_{1-6})\text{alkyl}$ ,  $(\text{C}_{3-7})\text{cycloalkyl}$ , aryl, heteroaryl, heteroalicyclic,  $(\text{C}_{1-6})\text{alkoxy}$ , aryloxy, heteroaryloxy, heteroalicycloxy,  $(\text{C}_{1-6})\text{thioalkoxy}$ , thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, are optionally substituted with one to nine same or different halogens or from one to five same or different substituents selected from the group G;

- G is selected from the group consisting of  $(\text{C}_{1-6})\text{alkyl}$ ,  $(\text{C}_{3-7})\text{cycloalkyl}$ , aryl, heteroaryl, heteroalicyclic, hydroxy,  $(\text{C}_{1-6})\text{alkoxy}$ , aryloxy, heteroaryloxy, heteroalicycloxy, thiohydroxy,  $(\text{C}_{1-6})\text{thioalkoxy}$ , thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, cyano, halogen, nitro, carbonyl, thiocarbonyl, benzyl, O-thiocarbamyl, N-thiocarbamyl, C-thioamido,  $-\text{NR}^{48}\text{C}(\text{O})-(\text{C}_{1-6})\text{alkyl}$ ,  $-\text{NR}^{48}\text{C}(\text{O})-(\text{C}_{3-6})\text{cycloalkyl}$ ,  $-\text{NR}^{48}\text{C}(\text{O})\text{-aryl}$ ,  $-\text{NR}^{48}\text{C}(\text{O})\text{-heteroaryl}$ ,  $-\text{NR}^{48}\text{C}(\text{O})\text{-heteroalicyclic}$ , a cyclic N-amido,  $-\text{NR}^{48}\text{S}(\text{O})_2-(\text{C}_{1-6})\text{alkyl}$ ,  $-\text{NR}^{48}\text{S}(\text{O})_2-(\text{C}_{3-6})\text{cycloalkyl}$ ,  $-\text{NR}^{48}\text{S}(\text{O})_2\text{-aryl}$ ,  $-\text{NR}^{48}\text{S}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NR}^{48}\text{S}(\text{O})_2\text{-heteroalicyclic}$ , O-carboxy, sulfinyl, sulfonyl, sulfonamide, phosphonyl,  $\text{NR}^{48}\text{R}^{49}$ ,  $(\text{C}_{1-6})\text{alkylC}(\text{O})\text{NR}^{48}\text{R}^{49}$ ,  $\text{C}(\text{O})\text{NR}^{48}\text{R}^{49}$ ,  $\text{NHC}(\text{O})\text{NR}^{48}\text{R}^{49}$ ,  $\text{OC}(\text{O})\text{NR}^{48}\text{R}^{49}$ ,  $\text{NHC}(\text{O})\text{OR}^{54}$ ,  $(\text{C}_{1-6})\text{alkylNR}^{48}\text{R}^{49}$ ,  $\text{COOR}^{54}$ , and  $(\text{C}_{1-6})\text{alkylCOOR}^{54}$ ;
- $\text{R}^7$  is selected from the group consisting of aryl, heteroaryl, and heteroalicyclic wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or with from one to three same or different substituents selected from the group F;
- $\text{R}^8$  is selected from the group consisting of hydrogen,  $(\text{C}_{1-6})\text{alkyl}$ ,  $(\text{C}_{3-7})\text{cycloalkyl}$ ,  $(\text{C}_{2-6})\text{alkenyl}$ ,  $(\text{C}_{3-7})\text{cycloalkenyl}$ ,  $(\text{C}_{2-6})\text{alkynyl}$ , aryl, heteroaryl, and heteroalicyclic wherein said  $(\text{C}_{1-6})\text{alkyl}$ ,  $(\text{C}_{3-7})\text{cycloalkyl}$ ,  $(\text{C}_{2-6})\text{alkenyl}$ ,  $(\text{C}_{3-7})\text{cycloalkenyl}$ ,  $(\text{C}_{2-6})\text{alkynyl}$ , aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to six same or different halogens or from one to five same or different substituents selected from the group F;

$R^9, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}$ , are each independently selected from the group consisting of hydrogen, or  $(C_{1-6})$ alkyl wherein each of said  $(C_{1-6})$ alkyl being optionally substituted with one to three same or different halogens;

5 X is selected from the group consisting of  $NR^5$ , O, and S;

$R^{40}$  and  $R^{41}$  are independently selected from the group consisting of  
 (a) hydrogen; (b)  $(C_{1-6})$ alkyl or  $(C_{3-7})$ cycloalkyl substituted with one to three same or different halogens or from one to two same or different substituents selected from the  
 10 group F; and (c)  $(C_{1-6})$ alkoxy, aryl, heteroaryl, heteroalicyclic or  $R^{40}$  and  $R^{41}$  taken together with the nitrogen to which they are attached form a heteroalicyclic ring which may contain up to 5 additional heteroatoms selected from N, O,  $S(O)_m$ , wherein  $m'$  is 0, 1, or 2; and wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or  
 15 different substituents selected from the group F;  
 with the proviso that only one of  $R^{40}$  and  $R^{41}$  may be hydrogen.

$R^{42}$  and  $R^{43}$  are independently selected from the group consisting of hydrogen,  
 $(C_{1-6})$ alkyl,  $(C_{1-6})$ alkoxy,  $(C_{3-7})$ cycloalkyl,  $(C_{2-6})$ alkenyl,  $(C_{3-7})$ cycloalkenyl,  
 20  $(C_{2-6})$ alkynyl, aryl, heteroaryl and heteroalicyclic; or  $R^{42}$  and  $R^{43}$  taken together with the nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to 5 additional heteroatoms selected from N, O,  $S(O)_m$ , wherein  $m'$  is 0, 1, or 2; and wherein said  $(C_{1-6})$ alkyl,  
 $(C_{1-6})$ alkoxy,  $(C_{3-7})$ cycloalkyl,  $(C_{2-6})$ alkenyl,  $(C_{3-7})$ cycloalkenyl,  $(C_{2-6})$ alkynyl, aryl,  
 25 heteroaryl, and heteroalicyclic are optionally substituted with one to nine same or different halogens or from one to five same or different substituents selected from the group G;

$R^{44}$  is selected from the group consisting of:

30

(1) H,  $(C_{1-6})$ alkyl,  $(C_{3-6})$ cycloalkyl,  $(C_{2-6})$ alkenyl,  $(C_{3-6})$ cycloalkenyl,  
 $(C_{2-6})$ alkynyl, halogen, CN, nitro, Ar,  $COOR^{50}$ ,  $COOAr$ ,  $-CONR_aR_b$ ,  $TR^{50}$ ,  $NR_aR_b$ ,  
 $-NC(O)NR_aR_b$ ,  $-OC(O)R^{50}$ ,  $-C[N(R_a)_2] = N-T-R_b$ ,  $YR^{50}$ ,  $-C(O)R^{50}$ ,  $-C(O)Ar$ ,  
 $-S(O)R_a$  and  $-S(O)_2R_a$ ; provided when  $R^{44}$  is  $-S(O)R_a$  or  $-S(O)_2R_a$  then  $R_a$  is not H; and

35

(2) a 4-7 membered heterocyclic ring, optionally substituted with  $R^{50}$ , which may contain 1-3 heteroatoms selected from the group consisting of O, S, SO,  $SO_2$ , N, and  $NR^{52}$ , wherein  $R^{52}$  is selected from the group consisting of hydrogen,  $(C_{1-4})$ alkyl,  $(C_{2-4})$ alkenyl and  $(C_{2-4})$ alkynyl;

5

T is S or O;

Ar is phenyl or heteroaryl; wherein said phenyl or heteroaryl is optionally substituted with one to three of the same or different halogens,  $C_{1-6}$  alkoxy,  $C_{1-6}$  alkyl or amino;

10

$R_a$  and  $R_b$  are each independently H,  $(C_{1-6})$ alkyl or phenyl;

$R^{46}$  is selected from the group consisting of H,  $OR^8$ , and  $NR^{40}R^{41}$ ;

15  $R^{47}$  is selected from the group consisting of H, amino, halogen, and  $(C_{1-6})$ alkyl;

$R^{48}$  and  $R^{49}$  are independently selected from the group consisting of hydrogen,  $(C_{1-6})$ alkyl,  $(C_{1-6})$ alkoxy,  $(C_{3-7})$ cycloalkyl,  $(C_{2-6})$ alkenyl,  $(C_{3-7})$ cycloalkenyl,  $(C_{2-6})$ alkynyl, aryl, heteroaryl and heteroalicyclic; or  $R^{48}$  and  $R^{49}$  taken together with the nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to 5 additional heteroatoms selected from N, O,  $S(O)_m$ , wherein  $m'$  is 0, 1, or 2;

20

$R^{50}$  is selected from the group consisting of H,  $(C_{1-6})$ alkyl,  $(C_{3-6})$ cycloalkyl, and benzyl; each of said alkyl, cycloalkyl and benzyl being optionally substituted with one to three same or different halogen, amino, OH, CN or  $NO_2$ ;

25

$R^{51}$  is selected from the group consisting of H,  $(C_{1-6})$ alkyl,  $(C_{3-6})$ cycloalkyl,  $(C_{2-6})$ alkenyl,  $(C_{3-6})$ cycloalkenyl,  $(C_{2-6})$ alkynyl and  $C(O)R^{53}$ ; wherein  $R^{53}$  is H,  $(C_{1-6})$ alkyl, or  $(C_{3-6})$ cycloalkyl and each of said  $(C_{1-6})$ alkyl and  $(C_{3-6})$ cycloalkyl being optionally substituted with one to three same or different halogen, amino, OH, CN or  $NO_2$ ;

30

Y is O, S or  $NR^{50}R^{51}$ ;

35

$R^{54}$  is selected from the group consisting of hydrogen,  $(C_{1-6})$ alkyl,


(C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic; wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to six same or different halogens or from one to five same or  
 5 different substituents selected from the group consisting of amino, OH, CN and NO<sub>2</sub>;

R<sup>54'</sup> is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic; wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl,  
 10 (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to six same or different halogens or from one to five same or different substituents selected from the group consisting of amino, OH, CN and NO<sub>2</sub>;

R<sup>55</sup> and R<sup>56</sup> are independently selected from the group consisting of hydrogen,  
 15 (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl and (C<sub>2-6</sub>)alkynyl; and

R<sup>57</sup> is selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl and (C<sub>2-6</sub>)alkynyl;  
 with the proviso that in the formulas above the carbon atoms which comprise the  
 20 carbon-carbon double bond of any alkenyl or the carbon-carbon triple bond of said alkynyl are not the point of attachment to the oxygen, nitrogen, or sulfur to which it is said to be attached;

2. A compound of claim 1, including pharmaceutically acceptable salts thereof  
 25 wherein,



I

wherein:

30

R<sup>1</sup> is hydrogen;

R<sup>2</sup> and R<sup>3</sup>, are each independently selected from the group (a)-(k) consisting of:

- (a) hydrogen,  
 (b) halogen,  
 (c) cyano,  
 (d) nitro,  
 5 (e) amino,  
 (f) C<sub>1-4</sub>alkylamino,  
 (g) di(C<sub>1-2</sub>alkyl)amino,  
 (h) hydroxy,  
 (i) C<sub>1-3</sub>alkyl optionally substituted with one to three same or different  
 10 halogen, hydroxy, C<sub>1-2</sub>alkoxy, amino, C<sub>1-4</sub>alkylamino, di  
 (C<sub>1-4</sub>alkyl)amino, cyano,  
 (j) C<sub>1-6</sub>alkoxy,  
 (k) heteroaryl, said heteroaryl is selected from the group consisting of  
 pyridinyl, pyrazinyl, pyridazinyl, pyrimidinyl, furanyl, thienyl, benzothienyl,  
 15 thiazolyl, isothiazolyl, oxazolyl, benzooxazolyl, isoxazolyl, imidazolyl,  
 benzoimidazolyl, 1H-imidazo[4,5-b]pyridin-2-yl, 1H-imidazo[4,5-c]pyridin-2-yl,  
 oxadiazolyl, thiadiazolyl, pyrazolyl, tetrazolyl, tetrazinyl, triazinyl and triazolyl, and  
 said heteroaryl is optionally substituted with C<sub>1-6</sub> alkyl groups  
 (l) phenyl which is independently substituted with one to three same or different  
 20 halogen, hydroxy, C<sub>1-2</sub>alkoxy, amino, C<sub>1-4</sub>alkylamino, di  
 (C<sub>1-4</sub>alkyl)amino, cyano,

R<sup>4</sup> is selected from the group consisting of hydrogen, halogen, cyano, nitro, COOR<sup>8</sup>,  
 XR<sup>57</sup>, C(O)R<sup>57</sup>, C(O)NR<sup>55</sup>R<sup>56</sup>, B, D, and E with the proviso that when at least one of  
 25 R<sup>2</sup> or R<sup>3</sup> is not either heteroaryl or substituted phenyl, then R<sub>4</sub> is selected from B or E;

m is 2;

R<sup>5</sup> is hydrogen;

30

R<sup>6</sup> does not exist;

A is selected from the group consisting of C<sub>1-6</sub>alkoxy, aryl and heteroaryl; in  
 which said aryl is phenyl or said heteroaryl is selected from the group consisting of  
 35 pyridinyl, pyrimidinyl, pyrazinyl, triazinyl, furanyl, thienyl, pyrrolyl, imidazolyl,  
 thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, quinolinyl, isoquinolinyl, benzofuranyl,  
 benzothienyl, benzoimidazolyl and benzothiazolyl; and said aryl or heteroaryl is



optionally substituted with one or two of the same or different amino, cyano, C<sub>1-6</sub>alkoxy, C<sub>1-6</sub>alkyl, -NHC(O)CH<sub>3</sub>, halogen and trifluoromethyl;

5 B is selected from the group consisting of -C(=NR<sup>46</sup>)(R<sup>47</sup>), C(O)NR<sup>40</sup>R<sup>41</sup>, aryl, heteroaryl, heteroalicyclic, S(O)<sub>q</sub>R<sup>8</sup>, P(O)(R<sup>8</sup>)<sub>q</sub>(OR<sup>8</sup>)<sub>2-q</sub>, P(S)(R<sup>8</sup>)<sub>q</sub>(OR<sup>8</sup>)<sub>2-q</sub>, C(O)R<sup>8</sup>, XR<sup>8</sup>, (C<sub>1-6</sub>)alkylNR<sup>40</sup>R<sup>41</sup>, and (C<sub>1-6</sub>)alkylCOOR<sup>8</sup> wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

10 q is 0, 1, or 2;

D is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, and (C<sub>2-6</sub>)alkynyl are optionally substituted with one to nine same or different halogens or from one to five same or different substituents selected from the group F;

15

E is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, and (C<sub>2-6</sub>)alkynyl are substituted with B;

20

F is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, (C<sub>1-6</sub>)alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, thiohydroxy, (C<sub>1-6</sub>)thioalkoxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, cyano, halogen, nitro, carbonyl, thiocarbonyl, benzyl, O-thiocarbamyl, N-thiocarbamyl, C-thioamido, -NR<sup>42</sup>C(O)-(C<sub>1-6</sub>)alkyl, -NR<sup>42</sup>C(O)-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>42</sup>C(O)-aryl, -NR<sup>42</sup>C(O)-heteroaryl, -NR<sup>42</sup>C(O)-heteroalicyclic, a cyclic N-amido, -NR<sup>42</sup>S(O)<sub>2</sub>-(C<sub>1-6</sub>)alkyl, -NR<sup>42</sup>S(O)<sub>2</sub>-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>42</sup>S(O)<sub>2</sub>-aryl, -NR<sup>42</sup>S(O)<sub>2</sub>-heteroaryl, -NR<sup>42</sup>S(O)<sub>2</sub>-heteroalicyclic, sulfinyl, sulfonyl, -S(O)<sub>2</sub>NR<sup>42</sup>R<sup>43</sup>, phosphonyl, NR<sup>42</sup>R<sup>43</sup>, (C<sub>1-6</sub>)alkylC(O)NR<sup>42</sup>R<sup>43</sup>, C(O)NR<sup>42</sup>R<sup>43</sup>, NHC(O)NR<sup>42</sup>R<sup>43</sup>, OC(O)NR<sup>42</sup>R<sup>43</sup>, NHC(O)OR<sup>54</sup>, (C<sub>1-6</sub>)alkylNR<sup>42</sup>R<sup>43</sup>, COOR<sup>54</sup>, and (C<sub>1-6</sub>)alkylCOOR<sup>54</sup> wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, (C<sub>1-6</sub>)alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, (C<sub>1-6</sub>)thioalkoxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;

25

30

35

G is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, (C<sub>1-6</sub>)alkoxy, aryloxy, heteroaryloxy, heteroalicycloxy, thiohydroxy, (C<sub>1-6</sub>)thioalkoxy, thioaryloxy, thioheteroaryloxy, thioheteroalicycloxy, cyano, halogen, nitro, carbonyl, thiocarbonyl, benzyl, O-thiocarbamyl, N-thiocarbamyl, C-thioamido, -NR<sup>48</sup>C(O)-(C<sub>1-6</sub>)alkyl, -NR<sup>48</sup>C(O)-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>48</sup>C(O)-aryl, -NR<sup>48</sup>C(O)-heteroaryl, -NR<sup>48</sup>C(O)-heteroalicyclic, a cyclic N-amido, -NR<sup>48</sup>S(O)<sub>2</sub>-(C<sub>1-6</sub>)alkyl, -NR<sup>48</sup>S(O)<sub>2</sub>-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>48</sup>S(O)<sub>2</sub>-aryl, -NR<sup>48</sup>S(O)<sub>2</sub>-heteroaryl, -NR<sup>48</sup>S(O)<sub>2</sub>-heteroalicyclic, sulfinyl, sulfonyl, -S(O)<sub>2</sub>NR<sup>48</sup>R<sup>49</sup>, NR<sup>48</sup>R<sup>49</sup>, (C<sub>1-6</sub>)alkyl C(O)NR<sup>48</sup>R<sup>49</sup>, C(O)NR<sup>48</sup>R<sup>49</sup>, NHC(O)NR<sup>48</sup>R<sup>49</sup>, OC(O)NR<sup>48</sup>R<sup>49</sup>, NHC(O)OR<sup>54</sup>, (C<sub>1-6</sub>)alkylNR<sup>48</sup>R<sup>49</sup>, COOR<sup>54</sup>, and (C<sub>1-6</sub>)alkylCOOR<sup>54</sup>;

R<sup>7</sup> is selected from the group consisting of aryl, heteroaryl, and heteroalicyclic wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or with from one to two same or different substituents selected from the group F;

R<sup>8</sup> is selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, R<sup>18</sup>, R<sup>19</sup> and R<sup>20</sup> are each independently selected from hydrogen or C<sub>1-3</sub>alkyl being optionally substituted with one to three fluorines;

R<sup>23</sup>, R<sup>24</sup>, R<sup>25</sup>, R<sup>26</sup>, R<sup>27</sup>, R<sup>28</sup>, R<sup>29</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, wherein each of said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl being optionally substituted with one to three same or different substituents selected from the group consisting of halogen, hydroxy, cyano, amino and nitro;

X is selected from the group consisting of NR<sup>5</sup>, O, and S;

R<sup>40</sup> and R<sup>41</sup> are independently selected from the group consisting of

Hydrogen;

- or (C<sub>1-6</sub>)alkyl or (C<sub>3-7</sub>)cycloalkyl substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F; or (C<sub>1-6</sub>)alkoxy, aryl, heteroaryl, heteroalicyclic or R<sup>40</sup> and R<sup>41</sup> taken together with the nitrogen to which they are attached form a heteroalicyclic ring which may contain up to 2 additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2; and wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;
- 10 with the proviso that only one of R<sup>40</sup> and R<sup>41</sup> may be hydrogen.

- R<sup>42</sup> and R<sup>43</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic or R<sup>42</sup> and R<sup>43</sup> taken together with the nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2; and wherein said (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;
- 15
- 20

- R<sup>44</sup> is selected from the group consisting of -H
- 25

R<sub>a</sub> and R<sub>b</sub> are each independently H, (C<sub>1-6</sub>)alkyl or phenyl;

R<sup>46</sup> is selected from the group consisting of H, OR<sup>8</sup>, and NR<sup>40</sup>R<sup>41</sup>;

- 30 R<sup>47</sup> is selected from the group consisting of H, amino, halogen, and (C<sub>1-6</sub>)alkyl;

- R<sup>48</sup> and R<sup>49</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, allyl, aryl, heteroaryl, heteroalicyclic or R<sup>48</sup> and R<sup>49</sup> taken together with the nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2;
- 35

$R^{50}$  is selected from the group consisting of H,  $(C_{1-6})$ alkyl,  $(C_{3-6})$ cycloalkyl, and benzyl, each of said alkyl, cycloalkyl and benzyl being optionally substituted with one to three same or different halogen, amino, OH, CN or  $NO_2$ ;

- 5  $R^{51}$  is selected from the group consisting of H,  $(C_{1-6})$ alkyl,  $(C_{3-6})$ cycloalkyl,  $(C_{2-6})$ alkenyl,  $(C_{3-6})$ cycloalkenyl,  $(C_{2-6})$ alkynyl or  $C(O)R^{53}$ , wherein  $R^{53}$  is H,  $(C_{1-6})$ alkyl, or  $(C_{3-6})$ cycloalkyl and each of said  $(C_{1-6})$ alkyl and  $(C_{3-6})$ cycloalkyl being optionally substituted with one to three same or different halogen, amino, OH, CN or  $NO_2$ ;

10

Y is O, S or  $NR^{50}R^{51}$ ;

- $R^{54}$  is selected from the group consisting of hydrogen,  $(C_{1-6})$ alkyl,  $(C_{3-7})$ cycloalkyl, allyl, aryl, heteroaryl, and heteroalicyclic wherein said
- 15  $(C_{1-6})$ alkyl,  $(C_{3-7})$ cycloalkyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group consisting of: amino, OH, and  $NR^{55}R^{56}$ ;

- $R^{54'}$  is selected from the group consisting of  $(C_{1-6})$ alkyl,  $(C_{3-7})$ cycloalkyl, allyl, aryl, heteroaryl, and heteroalicyclic wherein said
- 20  $(C_{1-6})$ alkyl,  $(C_{3-7})$ cycloalkyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group consisting of: amino, OH, and  $NR^{55}R^{56}$ ;

- 25  $R^{55}$  and  $R^{56}$  are independently selected from the group consisting of hydrogen,  $(C_{1-6})$ alkyl, allyl, or  $(C_{3-7})$ cycloalkyl; and

$R^{57}$  is selected from the group consisting of hydrogen,  $(C_{1-6})$ alkyl,  $(C_{3-7})$ cycloalkyl,  $(C_{2-6})$ alkenyl,  $(C_{3-7})$ cycloalkenyl,  $(C_{2-6})$ alkynyl.

30

3. A compound of claim 2, including pharmaceutically acceptable salts thereof, wherein:

- A is selected from the group consisting of phenyl and heteroaryl in which said
- 35 heteroaryl is selected from pyridinyl, furanyl and thienyl, and said phenyl or said

heteroaryl is optionally substituted with one to two of the same or different amino, C<sub>1-6</sub>alkyl, or halogen;

-- represents a carbon-carbon bond;


5

R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup>, and R<sup>14</sup> are each hydrogen; and

R<sup>15</sup> and R<sup>16</sup> are each independently hydrogen or methyl with the proviso that only one is methyl.

10

Q is either




and then R<sup>2</sup> is selected from the group consisting of hydrogen, halogen and methoxy; and

15

R<sub>3</sub> is hydrogen;

Or Q is:



20

and R<sup>2</sup> is halogen or hydrogen and R<sup>3</sup> is hydrogen;

25 R<sup>4</sup> is selected from the group consisting of B or E;

B is selected from the group consisting of -C(O)NR<sup>40</sup>R<sup>41</sup>, substituted phenyl, heteroaryl, and C(O)R<sup>7</sup> wherein said heteroaryl is optionally substituted and phenyl is substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

30

E is selected from the group consisting of (C<sub>2</sub>)alkenyl, or (C<sub>2</sub>)alkynyl, wherein (C<sub>2</sub>)alkenyl or (C<sub>2</sub>)alkynyl are substituted with B;

F is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-6</sub>)cycloalkyl, aryl,  
 5 heteroaryl, heteroalicyclic, hydroxy, (C<sub>1-6</sub>)alkoxy, (C<sub>1-6</sub>)thioalkoxy, cyano, halogen, carbonyl, benzyl, -NR<sup>42</sup>C(O)-(C<sub>1-6</sub>)alkyl, -NR<sup>42</sup>C(O)-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>42</sup>C(O)-aryl, -NR<sup>42</sup>C(O)-heteroaryl, -NR<sup>42</sup>C(O)-heteroalicyclic, a cyclic N-amido, -NR<sup>42</sup>S(O)<sub>2</sub>-(C<sub>1-6</sub>)alkyl, -NR<sup>42</sup>S(O)<sub>2</sub>-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>42</sup>S(O)<sub>2</sub>-aryl, -NR<sup>42</sup>S(O)<sub>2</sub>-heteroaryl, -NR<sup>42</sup>S(O)<sub>2</sub>-heteroalicyclic, -S(O)<sub>2</sub>NR<sup>42</sup>R<sup>43</sup>, NR<sup>42</sup>R<sup>43</sup>,  
 10 (C<sub>1-6</sub>)alkylC(O)NR<sup>42</sup>R<sup>43</sup>, C(O)NR<sup>42</sup>R<sup>43</sup>, NHC(O)NR<sup>42</sup>R<sup>43</sup>, OC(O)NR<sup>42</sup>R<sup>43</sup>, NHC(O)OR<sup>54</sup>, (C<sub>1-6</sub>)alkylNR<sup>42</sup>R<sup>43</sup>, COOR<sup>54</sup>, and (C<sub>1-6</sub>)alkylCOOR<sup>54</sup> wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-6</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, (C<sub>1-6</sub>)alkoxy, are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;

15 G is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy, (C<sub>1-6</sub>)alkoxy, (C<sub>1-6</sub>)thioalkoxy, thioaryloxy, cyano, halogen, nitro, carbonyl, thiocarbonyl, benzyl, -NR<sup>48</sup>C(O)-(C<sub>1-6</sub>)alkyl, -NR<sup>48</sup>C(O)-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>48</sup>C(O)-aryl, -NR<sup>48</sup>C(O)-heteroaryl, -NR<sup>48</sup>C(O)-heteroalicyclic, a cyclic N-amido, -NR<sup>48</sup>S(O)<sub>2</sub>-(C<sub>1-6</sub>)alkyl, -NR<sup>48</sup>S(O)<sub>2</sub>-(C<sub>3-6</sub>)cycloalkyl, -NR<sup>48</sup>S(O)<sub>2</sub>-aryl, -NR<sup>48</sup>S(O)<sub>2</sub>-heteroaryl, -NR<sup>48</sup>S(O)<sub>2</sub>-heteroalicyclic, sulfonyl, -S(O)<sub>2</sub>NR<sup>48</sup>R<sup>49</sup>, NR<sup>48</sup>R<sup>49</sup>, (C<sub>1-6</sub>)alkyl C(O)NR<sup>48</sup>R<sup>49</sup>, C(O)NR<sup>48</sup>R<sup>49</sup>, NHC(O)NR<sup>48</sup>R<sup>49</sup>, OC(O)NR<sup>48</sup>R<sup>49</sup>, NHC(O)OR<sup>54</sup>, (C<sub>1-6</sub>)alkylNR<sup>48</sup>R<sup>49</sup>, COOR<sup>54</sup>, and (C<sub>1-6</sub>)alkylCOOR<sup>54</sup>;

25 R<sup>7</sup> is selected from the group consisting of aryl, heteroaryl, and heteroalicyclic wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or with from one to two same or different substituents selected from the group F;

30 R<sup>8</sup> is selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, and (C<sub>3-7</sub>)cycloalkyl, wherein (C<sub>1-6</sub>)alkyl, and (C<sub>3-7</sub>)cycloalkyl are optionally substituted with one to six same or different halogens or from one to two same or different substituents selected from the group F;

35 R<sup>40</sup> and R<sup>41</sup> are independently selected from the group consisting of Hydrogen;

- or (C<sub>1-6</sub>)alkyl or (C<sub>3-7</sub>)cycloalkyl substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F; or (C<sub>1-6</sub>)alkoxy, aryl, heteroaryl, heteroalicyclic or R<sup>40</sup> and R<sup>41</sup> taken together with the nitrogen to which they are attached form a heteroalicyclic ring which may contain up to 2 additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2; and wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F; with the proviso that only one of R<sup>40</sup> and R<sup>41</sup> may be hydrogen.
- R<sup>42</sup> and R<sup>43</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic or R<sup>42</sup> and R<sup>43</sup> taken together with the nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2; and wherein said (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;
- R<sup>44</sup> is selected from the group consisting of -H
- R<sup>48</sup> and R<sup>49</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, heteroalicyclic or R<sup>48</sup> and R<sup>49</sup> taken together with the nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2;
- R<sup>54</sup> is selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, and heteroalicyclic wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group consisting of: amino, OH, and NR<sup>55</sup>R<sup>56</sup>; R<sup>54</sup> is selected from the group consisting of (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, and heteroalicyclic wherein said (C<sub>1-6</sub>)alkyl, (C<sub>3-7</sub>)cycloalkyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group consisting of: amino, OH, and NR<sup>55</sup>R<sup>56</sup>;

$R^{55}$  and  $R^{56}$  are independently selected from the group consisting of hydrogen,  $(C_{1-6})$ alkyl, or  $(C_{3-7})$ cycloalkyl

- 5 4. A compound of claim 3, including pharmaceutically acceptable salts thereof, wherein:

$R^4$  is selected from the group consisting of B;

- 10 B is selected from the group consisting of  $-C(O)NR^{40}R^{41}$ , substituted phenyl, or heteroaryl, wherein said phenyl is substituted and heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

15

F is selected from the group consisting of  $(C_{1-6})$ alkyl,  $(C_{3-6})$ cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxy,  $(C_{1-6})$ alkoxy,  $(C_{1-6})$ thioalkoxy, cyano, halogen, carbonyl, benzyl,  $-NR^{42}C(O)-(C_{1-6})$ alkyl,  $-NR^{42}C(O)-(C_{3-6})$ cycloalkyl,  $-NR^{42}C(O)$ -aryl,  $-NR^{42}C(O)$ -heteroaryl,  $-NR^{42}C(O)$ -heteroalicyclic, a cyclic N-amido,  $-NR^{42}S(O)_2-(C_{1-6})$ alkyl,  $-NR^{42}R^{43}$ ,  $C(O)NR^{42}R^{43}$ ,  $COOR^{54}$ , and wherein said  $(C_{1-6})$ alkyl,  $(C_{3-6})$ cycloalkyl, aryl, heteroaryl, heteroalicyclic,  $(C_{1-6})$ alkoxy, are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;

20

- 25 G is selected from the group consisting of  $(C_{1-6})$ alkyl, hydroxy,  $(C_{1-6})$ alkoxy, halogen,  $-NR^{48}C(O)-(C_{1-6})$ alkyl,  $-NR^{48}C(O)-(C_{3-6})$ cycloalkyl, a cyclic N-amido,  $-NR^{48}S(O)_2-(C_{1-6})$ alkyl,  $NR^{48}R^{49}$ ,  $(C_{1-6})$ alkyl  $C(O)NR^{48}R^{49}$ ,  $C(O)NR^{48}R^{49}$ ,  $(C_{1-6})$ alkyl  $NR^{48}R^{49}$ ;

- 30  $R^{40}$  is Hydrogen;

$R^{41}$  is  $(C_{1-3})$ alkoxy, heteroaryl, or aryl, wherein said aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;

35

$R^{42}$  and  $R^{43}$  are independently selected from the group consisting of hydrogen,  $(C_{1-6})$ alkyl,  $(C_{1-6})$ alkoxy,  $(C_{3-7})$ cycloalkyl, aryl, heteroaryl, heteroalicyclic or  $R^{42}$  and  $R^{43}$  taken together with the nitrogen to which they are attached form a heteroaryl ring




- or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, O, S(O)<sub>m</sub>, wherein m' is 0, 1, or 2; and wherein said (C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>3-7</sub>)cycloalkyl, (C<sub>2-6</sub>)alkenyl, (C<sub>3-7</sub>)cycloalkenyl, (C<sub>2-6</sub>)alkynyl, aryl, heteroaryl, and heteroalicyclic are optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group G;

- R<sup>48</sup> and R<sup>49</sup> are independently selected from the group consisting of hydrogen, (C<sub>1-6</sub>)alkyl or R<sup>48</sup> and R<sup>49</sup> taken together with the nitrogen to which they are attached form a heteroaryl ring or a heteroalicyclic ring which may contain up to two additional heteroatoms selected from N, or O;

5. A compound of claim 3, including pharmaceutically acceptable salts thereof, wherein:

Q is



20


R<sup>4</sup> is B;

A is Phenyl or 2-pyridyl;

- 25 B is selected from the group consisting of -C(O)NR<sup>40</sup>R<sup>41</sup> or heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

6. A compound of claim 4, including pharmaceutically acceptable salts thereof, wherein:

30 Q is



R<sup>4</sup> is B;

5 A is Phenyl or 2-pyridyl;

B is selected from the group consisting of -C(O)NR<sup>40</sup>R<sup>41</sup> or heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

10

7. A compound of claim 5, including pharmaceutically acceptable salts thereof, wherein:

B is heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

15


8. A compound of claim 6, including pharmaceutically acceptable salts thereof, wherein:

B is heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

20

9. A compound of claim 3, including pharmaceutically acceptable salts thereof, wherein:

25 Q is




R<sup>2</sup> is selected from the group consisting of hydrogen, halogen, and methoxy;

R<sup>4</sup> is B;

B is selected from the group consisting of -C(O)NR<sup>40</sup>R<sup>41</sup> or heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or  
5 from one to two same or different substituents selected from the group F;

10. A compound of claim 4, including pharmaceutically acceptable salts thereof, wherein:

Q is



10

R<sup>2</sup> is selected from the group consisting of hydrogen, halogen, and methoxy;

R<sup>4</sup> is B;

15 B is selected from the group consisting of -C(O)NR<sup>40</sup>R<sup>41</sup> or heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

20 11. A compound of claim 9, including pharmaceutically acceptable salts thereof, wherein:

A is Phenyl or 2-pyridyl;

12. A compound of claim 10, including pharmaceutically acceptable salts thereof, wherein:

25 A is Phenyl or 2-pyridyl;

13. A compound of claim 11 including pharmaceutically acceptable salts thereof, wherein:

B is -C(O)NR<sup>40</sup>R<sup>41</sup>;

30

14. A compound of claim 12 including pharmaceutically acceptable salts thereof, wherein:

B is  $-\text{C}(\text{O})\text{NR}^{40}\text{R}^{41}$

15. A compound of claim 11 including pharmaceutically acceptable salts thereof, wherein:

- 5 B is heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

16. A compound of claim 12 including pharmaceutically acceptable salts thereof, wherein:

B is heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

15

17. A compound of claim 3 in which:

$\text{R}^4$  is B; and

- 20 F is selected from the group consisting of  $(\text{C}_{1-6})$ alkyl, hydroxy, heteroaryl, heteroalicyclic, methoxy, methylthioalkoxy, halogen, carbonyl,  $\text{C}(\text{O})\text{NR}^{42}\text{R}^{43}$ ,  $-\text{NR}^{42}\text{C}(\text{O})-(\text{C}_{1-6})$ alkyl,  $-\text{NR}^{42}\text{C}(\text{O})-(\text{C}_{3-6})$ cycloalkyl,  $-\text{NR}^{42}\text{C}(\text{O})$ -aryl,  $-\text{NR}^{42}\text{C}(\text{O})$ -heteroaryl,  $-\text{NR}^{42}\text{C}(\text{O})$ -heteroalicyclic, a cyclic N-amido,  $-\text{NR}^{42}\text{S}(\text{O})_2-(\text{C}_{1-6})$ alkyl,  $-\text{NR}^{42}\text{S}(\text{O})_2-(\text{C}_{3-6})$ cycloalkyl,  $-\text{NR}^{42}\text{S}(\text{O})_2$ -aryl,  $-\text{NR}^{42}\text{S}(\text{O})_2$ -heteroaryl,  $-\text{NR}^{42}\text{S}(\text{O})_2$ -heteroalicyclic,  $\text{NR}^{42}\text{R}^{43}$ ,  $\text{COOH}$ ;
- 25

18. A compound of claim 4 in which:

$\text{R}^4$  is B; and

30

- F is selected from the group consisting of  $(\text{C}_{1-6})$ alkyl, hydroxy, heteroaryl, heteroalicyclic, methoxy, methylthioalkoxy, halogen, carbonyl,  $\text{C}(\text{O})\text{NR}^{42}\text{R}^{43}$ ,  $-\text{NR}^{42}\text{C}(\text{O})-(\text{C}_{1-6})$ alkyl,  $-\text{NR}^{42}\text{C}(\text{O})-(\text{C}_{3-6})$ cycloalkyl,  $-\text{NR}^{42}\text{C}(\text{O})$ -aryl,  $-\text{NR}^{42}\text{C}(\text{O})$ -heteroaryl,  $-\text{NR}^{42}\text{C}(\text{O})$ -heteroalicyclic, a cyclic N-amido,  $-\text{NR}^{42}\text{S}(\text{O})_2-(\text{C}_{1-6})$ alkyl,  $-\text{NR}^{42}\text{S}(\text{O})_2-(\text{C}_{3-6})$ cycloalkyl,  $-\text{NR}^{42}\text{S}(\text{O})_2$ -aryl,  $-\text{NR}^{42}\text{S}(\text{O})_2$ -heteroaryl,  $-\text{NR}^{42}\text{S}(\text{O})_2$ -heteroalicyclic,  $\text{NR}^{42}\text{R}^{43}$ ,  $\text{COOH}$ ;
- 35

19. A compound of claim 17 in which:


A is Phenyl or 2-pyridyl;

20. A compound of claim 18 in which:

A is Phenyl or 2-pyridyl;

- 5 21. A compound of claim 3, including pharmaceutically acceptable salts thereof, wherein:

Q is



10

$R^2$  is selected from the group consisting of hydrogen or methoxy;


$R^3$  is hydrogen;

- 15  $R^4$  is B;

B is selected from the group consisting of  $-C(O)NR^{40}R^{41}$  or heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

- 20 22. A compound of claim 4, including pharmaceutically acceptable salts thereof, wherein:

Q is



25

$R^2$  is selected from the group consisting of hydrogen or methoxy;

R<sup>3</sup> is hydrogen;

R<sup>4</sup> is B;

- 5 B is selected from the group consisting of -C(O)NR<sup>40</sup>R<sup>41</sup> or heteroaryl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

23. A compound of claim 11 wherein R<sup>2</sup> is Fluoro;
- 10 24. A compound of claim 12 wherein R<sup>2</sup> is Fluoro;
25. A compound of claim 11 wherein R<sup>2</sup> is Methoxy;
26. A compound of claim 12 wherein R<sup>2</sup> is Methoxy;
- 15 27. A compound of claim 11 wherein  
B is selected from the group consisting of thiazole, pyridazine, pyrazine, pyrazole, isoxazole, isothiazole, imidazole, furyl, thienyl, oxazole, oxadiazole, thiadiazole, pyrimidine, pyrazole, triazine, triazole, tetrazole, pyridyl, wherein said heteroaryl is  
20 optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;
28. A compound of claim 12 wherein  
B is selected from the group consisting of thiazole, pyridazine, pyrazine, pyrazole,  
25 isoxazole, isothiazole, imidazole, furyl, thienyl, oxazole, oxadiazole, thiadiazole, pyrimidine, pyrazole, triazine, triazole, tetrazole, pyridyl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;
- 30 29. A compound of claim 5 wherein  
B is selected from the group consisting of thiazole, pyridazine, pyrazine, pyrazole, isoxazole, isothiazole, imidazole, furyl, thienyl, oxazole, oxadiazole, thiadiazole, pyrimidine, pyrazole, triazine, triazole, tetrazole, pyridyl, wherein said heteroaryl is  
35 optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;
30. A compound of claim 6 wherein

B is selected from the group consisting of thiazole, pyridazine, pyrazine, pyrazole, isoxazole, isothiazole, imidazole, furyl, thienyl, oxazole, oxadiazole, thiadiazole, pyrimidine, pyrazole, triazine, triazole, tetrazole, pyridyl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two  
5 same or different substituents selected from the group F;

31. A compound of claim 21 wherein  
B is selected from the group consisting of thiazole, pyridazine, pyrazine, pyrazole, isoxazole, isothiazole, imidazole, furyl, thienyl, oxazole, oxadiazole, thiadiazole,  
10 pyrimidine, pyrazole, triazine, triazole, tetrazole, pyridyl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

32. A compound of claim 22 wherein  
15 B is selected from the group consisting of thiazole, pyridazine, pyrazine, pyrazole, isoxazole, isothiazole, imidazole, furyl, thienyl, oxazole, oxadiazole, thiadiazole, pyrimidine, pyrazole, triazine, triazole, tetrazole, pyridyl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

20  
33. A compound of claim 9 wherein  
B is selected from the group consisting of thiazole, pyridazine, pyrazine, pyrazole, isoxazole, isothiazole, imidazole, furyl, thienyl, oxazole, oxadiazole, thiadiazole, pyrimidine, pyrazole, triazine, triazole, tetrazole, pyridyl, wherein said heteroaryl is  
25 optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

34. A compound of claim 10 wherein  
B is selected from the group consisting of thiazole, pyridazine, pyrazine, pyrazole,  
30 isoxazole, isothiazole, imidazole, furyl, thienyl, oxazole, oxadiazole, thiadiazole, pyrimidine, pyrazole, triazine, triazole, tetrazole, pyridyl, wherein said heteroaryl is optionally substituted with one to three same or different halogens or from one to two same or different substituents selected from the group F;

35. A compound of claim 8 wherein  
B is is heteroaryl wherein said heteroaryl is optionally substituted with one to three same or different halogens or a substituent selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NHS(O)<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), methoxy, -C(O)-NH<sub>2</sub>,

C(O)NHMe, C(O)NMe<sub>2</sub>, trifluoromethyl, -NHC(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -heteroaryl, cyclic N-amido;

36. A compound of claim 8 wherein
- 5 B is -C(O)NH-heteroaryl wherein said heteroaryl is optionally substituted with one to three same or different halogens or a substituent selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -methoxy, -NHC(C<sub>1</sub>-C<sub>6</sub> alkyl), or -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>;
- 10 37. A compound of claim 16 wherein
- B is heteroaryl wherein said heteroaryl is optionally substituted with one to three same or different halogens or a substituent selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NHS(O)<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), methoxy, -C(O)-NH<sub>2</sub>, C(O)NHMe, C(O)NMe<sub>2</sub>, trifluoromethyl, -NHC(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -heteroaryl, cyclic N-amido;
- 15 38. A compound of claim 14 wherein
- B is -C(O)NH-heteroaryl wherein said heteroaryl is optionally substituted with one to three same or different halogens or a substituent selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -methoxy, -NHC(C<sub>1</sub>-C<sub>6</sub> alkyl), or -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>;
- 20 39. A compound of claim 28 wherein
- B is wherein said heteroaryl is optionally substituted with one to three same or different halogens or a substituent selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NHS(O)<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), methoxy, -C(O)-NH<sub>2</sub>, C(O)NHMe, C(O)NMe<sub>2</sub>, trifluoromethyl, -NHC(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -heteroaryl, cyclic N-amido;
- 25 40. A compound of claim 27 wherein
- B is -C(O)NH-heteroaryl wherein said heteroaryl is optionally substituted with one to three same or different halogens or a substituent selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -methoxy, -NHC(C<sub>1</sub>-C<sub>6</sub> alkyl), or -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>;
- 30 41. A compound of claim 6 wherein
- B is thienyl;
- 35



42. A compound of claim 40 wherein  
B is thienyl which is optionally substituted with one to three same or different  
halogens or a substituent selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-  
C<sub>6</sub> alkyl), -NHS(O)<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), methoxy, -C(O)-NH<sub>2</sub>, C(O)NHMe, C(O)NMe<sub>2</sub>,  
5 trifluoromethyl, -NHC(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -heteroaryl, cyclic N-amido;
43. A compound of claim 16 wherein  
B is thienyl;
- 10 44. A compound of claim 42 wherein  
B is thienyl which is optionally substituted with one to three same or different  
halogens or a substituent selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-  
C<sub>6</sub> alkyl), -NHS(O)<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), methoxy, -C(O)-NH<sub>2</sub>, C(O)NHMe, C(O)NMe<sub>2</sub>,  
trifluoromethyl, -NHC(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -heteroaryl, cyclic N-amido;  
15
45. A compound of claim 16 wherein  
B is thienyl;
- 20 46. A compound of claim 42 wherein  
B is thienyl which is optionally substituted with one to three same or different  
halogens or a substituent selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-  
C<sub>6</sub> alkyl), -NHS(O)<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), methoxy, -C(O)-NH<sub>2</sub>, C(O)NHMe, C(O)NMe<sub>2</sub>,  
trifluoromethyl, -NHC(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -heteroaryl, cyclic N-amido;
- 25 47. A compound of claim 28 wherein  
B is selected from the group consisting of thiazole, pyridazine, pyrazine, pyrazole,  
isoxazole, isothiazole, imidazole, furyl, thienyl, oxazole, oxadiazole, thiadiazole,  
pyrimidine, pyrazole, triazine, triazole, tetrazole, pyridyl, wherein said heteroaryl is  
optionally substituted with one to three same or different halogens or a substituent  
30 selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NHS(O)<sub>2</sub>-(C<sub>1</sub>-  
C<sub>6</sub> alkyl), methoxy, -C(O)-NH<sub>2</sub>, C(O)NHMe, C(O)NMe<sub>2</sub>, trifluoromethyl, -NHC(C<sub>1</sub>-  
C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -heteroaryl, cyclic N-amido;
48. A compound of claim 30 wherein  
35 B is selected from the group consisting of thiazole, pyridazine, pyrazine, pyrazole,  
isoxazole, isothiazole, imidazole, furyl, thienyl, oxazole, oxadiazole, thiadiazole,  
pyrimidine, pyrazole, triazine, triazole, tetrazole, pyridyl, wherein said heteroaryl is  
optionally substituted with one to three same or different halogens or a substituent

selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NHS(O)<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub> alkyl), methoxy, -C(O)-NH<sub>2</sub>, C(O)NHMe, C(O)NMe<sub>2</sub>, trifluoromethyl, -NHC(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -heteroaryl, cyclic N-amido;

- 5 49. A compound of claim 32 wherein  
B is selected from the group consisting of thiazole, pyridazine, pyrazine, pyrazole,  
isoxazole, isothiazole, imidazole, furyl, thienyl, oxazole, oxadiazole, thiadiazole,  
pyrimidine, pyrazole, triazine, triazole, tetrazole, pyridyl, wherein said heteroaryl is  
optionally substituted with one to three same or different halogens or a substituent  
10 selected from the group (C<sub>1</sub>-C<sub>6</sub> alkyl), amino, -NHC(O)-(C<sub>1</sub>-C<sub>6</sub> alkyl), -NHS(O)<sub>2</sub>-(C<sub>1</sub>-  
C<sub>6</sub> alkyl), methoxy, -C(O)-NH<sub>2</sub>, C(O)NHMe, C(O)NMe<sub>2</sub>, trifluoromethyl, -NHC(C<sub>1</sub>-  
C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)<sub>2</sub>, -heteroaryl, cyclic N-amido;
- 15 50. A compound of claim 3 which is depicted in Table 2;
51. A compound of claim 3 which is depicted in Table 3;
52. A compound of claim 3 which is depicted in Table 4
- 20 53. A compound of claim 3 which is depicted in Table 5
54. A pharmaceutical formulation which comprises an antiviral effective amount  
of a compound of Formula I, including pharmaceutically acceptable salts thereof, as  
claimed in any of claims 1-50, and a pharmaceutically acceptable carrier.
- 25 55. The pharmaceutical formulation of claim 51, useful for treating infection by  
HIV, which additionally comprises an antiviral effective amount of an AIDS  
treatment agent selected from the group consisting of:
- (a) an AIDS antiviral agent;
  - 30 (b) an anti-infective agent;
  - (c) an immunomodulator; and
  - (d) HIV entry inhibitors.

56. A method for treating mammals infected with a virus, comprising administering to said mammal an antiviral effective amount of a compound of Formula I, including pharmaceutically acceptable salts thereof, as claimed in any of claims 1-50.

5

57. The method of claim 53, comprising administering to said mammal an antiviral effective amount of a compound of Formula I in combination with an antiviral effective amount of an AIDS treatment agent selected from the group consisting of: an AIDS antiviral agent; an anti-infective agent; an immunomodulator;  
10 and HIV entry inhibitors.

58. The method of claim 54 wherein the virus is HIV.

59. The method of claim 53 wherein the virus is HIV.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/00455

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : A61P 31/18, 31/12; A61K 31/496, 31/501, 31/53, 31/506; C07D 401/14, 403/14, 241/02, 253/08, 251/02, 241/36, 471/02

US CL : 514/241, 242, 243, 249, 252.02, 252.11, 252.18, 253.04, 253.06, 257

**B. FIELDS SEARCHED**Minimum documentation searched (classification system followed by classification symbols)  
U.S. : 514/241, 242, 243, 249, 252.02, 252.11, 252.18, 253.04, 253.06, 257

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
CAS ONLINE**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96/11929 A1 (SMITHKLINE BEECHAM PLC) 25 April 1996.	1-59

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

06 May 2002 (06.05.2002)

Date of mailing of the international search report

31 MAY 2002

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Brenda L. Coleman

Telephone No. 703-308-1235